

Species composition and distribution of mayflies (Ephemeroptera) in relation to land use systems along the Thika River, Kenya

C Maina, BM Mwangi & JJ Jumbe

To cite this article: C Maina, BM Mwangi & JJ Jumbe (2021): Species composition and distribution of mayflies (Ephemeroptera) in relation to land use systems along the Thika River, Kenya, African Journal of Aquatic Science, DOI: [10.2989/16085914.2020.1842169](https://doi.org/10.2989/16085914.2020.1842169)

To link to this article: <https://doi.org/10.2989/16085914.2020.1842169>



Published online: 26 Feb 2021.



Submit your article to this journal [↗](#)



View related articles [↗](#)



View Crossmark data [↗](#)

Species composition and distribution of mayflies (Ephemeroptera) in relation to land use systems along the Thika River, Kenya

C Maina¹, BM Mwangi^{2*} and JJ Jumbe¹ 

¹ Department of Zoological sciences, Kenyatta University, Nairobi, Kenya

² Department of Physical and Biological Sciences, Murang'a University of Technology, Murang'a, Kenya

*Correspondence: bmmwangi@mut.ac.ke

Mayflies (order Ephemeroptera) are widely distributed and commonly used in biological assessment of water quality because of their sensitivity. Their application as bio-indicators in tropical streams and rivers, however, is hampered by lack of adequate information about their status in many such ecosystems. The current study investigated their species composition, distribution and abundance in relation to land use and water quality characteristics of the Thika River, Kenya. The Thika River emanates from the Aberdare Ridges and flows through forest, agricultural and urban catchments. Sampling was carried out at three sites representing forest, agricultural and urban land use systems. Dissolved oxygen, conductivity, total dissolved solids, temperature and transparency were measured *in situ* at each sampling station. Macroinvertebrates were sampled using a modified Hess sampler and the Ephemeroptera were later identified and sorted in the laboratory. Thirteen Ephemeroptera genera belonging to six families were obtained. Mean larvae abundance was 1 238 ind. m⁻². *Acanthiops* (70.4%) and *Afronurus* (13.3%) dominated the community at all sites. The current study revealed that decreasing Ephemeroptera taxa abundance along the river channel indicates decreasing water quality.

Keywords: abundance, *Acanthiops*, *Afronurus*, macroinvertebrates, water quality

Introduction

Good water quality is essential because it maintains the ecological processes that support aquatic biodiversity. Rivers emanating from forested catchment areas are generally of good water quality, as a result of minimal human activities (Garcia et al. 2014). However, declining water quality as a result of environmental perturbation affects the stability of the biodiversity, hence hindering the ecosystems services and functions of aquatic ecosystems (Ndung'u 2014). Water quality deterioration is, therefore, a significant threat in many rivers (Ndaruga 2004; Arimoro et al. 2007), which is as a result of growth in human population (DeSA 2013). In particular, industrialization, urbanization and agricultural activities have led to increased nutrient levels in the water bodies (Heathwaite et al. 1996). The Thika River emanates from a heavily forested catchment and then flows through various land use types including large coffee plantations, horticultural and animal farms, where it is heavily impacted by agricultural, industrial and domestic pollution.

The method of water quality assessments using macroinvertebrates as indicators has been widely exploited in temperate areas. Ephemeroptera is a widely distributed order of insects with >2 500 described species worldwide (Hubbard 1990). There are 23 families and 371 genera worldwide (Hubbard 1990). In Lake Erie, Ephemeroptera is successfully utilised in bio-monitoring (Schloesser and Nalepa 2002). A recent study showed peak emergence of burrowing mayfly nymphs (*Hexagenia* spp.) that was associated with an improvement of the ecosystem health

(Schloesser and Nalepa 2002). Mayflies play an essential role in almost all undisturbed freshwater communities, and their larvae frequently form a considerable part of the material sampled during bio-monitoring procedures (Bauernfeind and Moog 2000). In tropical areas, the method has been given less attention while most water quality assessments are done using the physical and chemical processes. Aschalew and Moog (2015) found that until present, conventional physico-chemical techniques are used in some streams for monitoring the river water quality. The current study therefore aimed at investigating the community structure of mayflies in relation to water quality changes along the Thika River, Kenya. This is by first assessing the water quality status of the Thika River as it flows downstream through the various land use systems, and relating this to the composition, distribution and abundance of Ephemeroptera in relation to water quality changes.

Materials and methods

Study area

The centroid for the Thika River catchment lies at 36°47'31.3" E, 0°46'52.2" S (Figure 1). Annual precipitation and temperature are relatively stable with the coolest months occurring from June to August. The hottest air temperatures are normally from December to March. The average annual rainfall in Thika Town and its surrounds ranges between 900 and 1 250 mm per year. Lake Ol-borosat influences the hydrology of Aberdare




Figure 1: Map of study area showing sampling sites 1, 2 and 3 within the Thika catchment in Murang'a county, Kenya. Site 1 was surrounded by natural forest in the Aberdare ranges, Site 2 by agricultural farmlands and Site 3 by industrial and urban settlements

ranges region, and the natural forest vegetation, both of which impact on the process of evapotranspiration. This eventually produces precipitation in the region. The precipitation replenishes the underground water content that feeds the river (Kotchoni et al. 2019). The river is also fed by surface runoff that is often contaminated with agrarian, municipal and industrial effluents (Kimani et al. 2016). The river flows through the border of

Murang'a-Kiambu and empties in River Tana. The river substratum consists mainly of a mixture of red volcanic, black cotton soils and murrum. Decomposing macrophytes and plant remains also form part of the river substrate. Three sites were selected along this river system from the forest to the downstream reaches.

The Thika River is impacted heavily by industrial and domestic wastes as it flows downstream from its source

(Karuri et al. 2003). For instance, the levels of heavy metals at a downstream site near Blue Post hotel were found to be higher than levels at upstream sites (Karuri et al. 2003). Inoti et al. (2012) found out that the concentration of heavy metals like iron, cadmium, chromium, and lead in the soil along the Thika River catchment areas was above the maximum permitted levels.

Site 1 is located approximately 1 km from the river's source (Figure 1). The shallower river channel is heavily shaded by the thick forest canopy, with more exposed pools at various points along its reaches. The dominant water flora was sub-merged and floating macrophytes, such as *Azolla africana*, *Salvania molesta* and *Nymphaea lotus*. Emergent macrophytes were also present, for instance, *Vossia cuspidate*, *Pycreus lanceolatus* and *Scirpus jacobi*. The streambed is granite with fallen leaves and huge rocks. The water in the site was clear with no visible siltation. The river margin was colonized by terrestrial vegetation. Trees dominating were *Elaeis guineensis*, *Cedrus deodara* and *Bambusa* spp. in addition to several grass species, like *Zostera marina* and *Ruppia maritima*. There were minimal human activities adjacent to the waters' edge, and the site was therefore selected to be used as a control point.

Site 2 (cultivated) was 29 km from site 1 (Figure 1). The riparian vegetation consists of *Lantana camara*, *Maranta arundinacea*, farming plants such as *Zea mays*, and napier grass. Riparian zones were heavily infested with non-native alien trees including *Eucalyptus* species and *Grevillae robusta*. The site was devoid of aquatic vegetation thus water was exposed to sunlight. Algal growth was extensive, which made the water turn green with a bad smell emanating from the water. The area had a low abundance of macrophytes such as *N. lotus* and *Salvania* sp. The substratum in this section is sandy loam. The location is downstream from extensive coffee plantations and horticultural farms, growing crops such as vegetables, tomatoes, flowers and French beans. There are also settlements, and livestock farming that have free access to the river.

Site 3 (urban) was 3 km from site 2 down the confluence of the river with Chania River (Figure 1). Although no vegetation occurred in the riparian zone, there were more aquatic plants at site 3 compared to site 2, with *Salvinia* sp., *N. lotus* and *M. arundinacea* predominating. The riparian zone was dominated by non-native alien trees such as *Acacia* spp. and *Eucalyptus globulus*. The substratum is principally clay and silt, and the water is exposed to direct sun. Thika Town dumps its sewage and industrial waste into the river after treatment at the site.

Sampling and laboratory analysis

Sampling was done monthly at each site for six months (September 2015–February 2016). Water depth was measured using a metre ruler, according to Arimoro and Muller 2010. River width was measured across the river using a tape measure. During the wet season of 2015, sites 2 and 3 became heavily flooded as a result of extensive rainfall requiring a boat to cross the river. Water transparency was measured in the field using a white and black Secchi disc (20 mm diameter). The Secchi disc was gently lowered into the water and the depth

of disappearance and reappearance recorded (Wetzel 2000). Water temperature was measured using a pocket thermometer. Water conductivity, pH and total dissolved solutes (TDS) were measured using a multi-probe meter (Hanna Instruments Model H199130, Romania). DO was measured using a portable dissolved oxygen meter (MRC, model DO-5510, UK). Three readings were taken per sampling event.

Macroinvertebrates were collected using a disturbance removal sampling technique (DRST) (Kage 2003) and a Hess sampler was used in shallow sites with a rocky substrate. An Eckman grab sampler was used on deeper sites with a muddy substratum. At each sampling site, three samples of macroinvertebrates were collected. These were from the middle and approximately 1 m from the left and right banks. The three samples were then pooled, representing a single sample for each site and fixed with 4% formalin. In the laboratory, the macroinvertebrates sampled, were filtered and washed free of sediments through a 250 µm sieve. All Ephemeroptera were identified to the lowest identifiable taxonomic level, using a dissecting microscope ($\times 10$ magnification) and relevant taxonomic keys at the National Museums of Kenya (Merit and cummins 1996; Barber-James and Lugo-Ortoz 2003; Huxley 2003). The unidentified individuals were taken to South Africa for further identification.

Data analysis

The data were checked for homogeneity of variance using Levene's *F*-test. A one-way ANOVA was used to determine differences in water quality parameters between sites. A Kruskal–Wallis test was used for determining the differences in Ephemeroptera abundances among the three sampling sites. The difference between community structure for dry and wet sampling seasons was tested using a *t*-test, because two means had to be compared. Correlation coefficient (*r*) analyses were done on the individual water quality parameters to identify the magnitude of relationships between them. It was also used to test the relationship between water quality and abundance of Ephemeroptera larvae along the Thika River. All the statistical analyses were done using SPSS program (version 22), and rejection of Null hypothesis set at $p \leq 0.05$.

Results

Physico-chemical parameters

Water depths increased significantly from site 1 downstream to site 3 ($F_{(2, 51)} = 40.3$; $p < 0.001$; Table 1). Water depths did not vary significantly between dry and wet seasons irrespective of sites ($t = 3.325$; $df = 52$; $p > 0.5$) (Table 2). River widths were significantly different among the sites ($F_{(2, 51)} = 83.71$, $p < 0.001$). Site 1 had the lowest mean 8.6 ± 0.54 m while site 3 had the highest mean of 17.1 ± 0.49 m (Table 1). Seasonal variation in river width was statistically different between the dry and wet seasons ($t = 2.556$; $df = 52$; $p < 0.5$) (Table 2). Spatial water transparency was not significantly different ($F_{(2, 51)} = 2.785$, $p > 0.5$). The seasonal variation in water transparencies was significantly different ($t = 3.663$; $df = 52$; $p < 0.5$) (Table 2). Water temperatures were significantly different among

Table 1: Physico-chemical characteristics (mean \pm SE) of the three sampling sites along the Thika River (September 2015–February 2016)

Parameter	Site			p-value
	Forested	Cultivated	Urban	
Temperature ($^{\circ}$ C)	14.3 \pm 0.28 ^a	21.9 \pm 0.48 ^b	21.9 \pm 0.36 ^b	<0.001*
DO (mg l ⁻¹)	12.1 \pm 0.26 ^b	8.7 \pm 0.11 ^a	8.8 \pm 0.09 ^a	<0.001*
Conductivity (μ S cm ⁻¹)	16.9 \pm 0.69 ^a	79.5 \pm 2.58 ^b	93.5 \pm 5.37 ^c	<0.001*
pH	6.1 \pm 0.60	6.3 \pm 0.10	6.4 \pm 0.90	0.112
TDS (mg l ⁻¹)	10.1 \pm 0.53 ^a	40.5 \pm 1.49 ^b	47.9 \pm 2.93 ^c	<0.001*
Transparency (m)	0.7 \pm 0.30	0.5 \pm 0.10	0.4 \pm 0.50	0.071
Width (m)	8.6 \pm 0.54 ^a	12.7 \pm 0.33 ^b	17.1 \pm 0.49 ^c	<0.001*
Depth (m)	0.5 \pm 0.05 ^a	1.1 \pm 0.08 ^b	1.4 \pm 0.09 ^c	<0.001*

Note: The same small letter within the same row indicates that means do not differ significantly from one another (One-way ANOVA, SNK-test, $\alpha = 0.5$). * Statistically significant: $p \leq 0.05$

Table 2: Seasonal variation of physico- chemical parameters of the Thika River in September 2015 to February 2016, mean \pm SE

Parameters	Dry	Wet	p-value
Temperature ($^{\circ}$ C)	20.56 \pm 0.77	18.19 \pm 0.70	0.027*
DO (mg l ⁻¹)	10.21 \pm 0.38	9.44 \pm 0.26	0.106
Conductivity (μ S cm ⁻¹)	60.46 \pm 7.41	66.10 \pm 6.74	0.576
pH	6.25 \pm 0.40	6.31 \pm 0.90	0.531
TDS (mg l ⁻¹)	30.53 \pm 3.70	35.11 \pm 3.35	0.364
Transparency (m)	0.67 \pm 0.55	0.41 \pm 0.50	<0.001*
Width (m)	11.48 \pm 0.67	14.11 \pm 0.78	0.014*
Depth (m)	0.81 \pm 0.80	1.23 \pm 0.10	0.380

* Statistically significant: $p \leq 0.05$

Table 3: Spatial variations in abundance (ind. m⁻²) of Ephemeroptera of the Thika River at the sites 1, 2 and 3 during the study period, September 2015 to February 2016

Family	Ephemeroptera taxa		Sites		
	Species		Forested	Cultivated	Urban
Oligoneuriidae	<i>Oligoneuriopsis</i>	<i>dobbsi</i>	143	0	0
Heptageniidae	<i>Afronurus</i>	sp. 1	295	42	72
Heptageniidae	<i>Afronurus</i>	sp. 2	2	2	0
Baetidae	<i>Acanthiops</i>	<i>cooperi</i>	0	5	0
Baetidae	<i>Acanthiops</i>	sp. 1	2 231	111	407
Baetidae	<i>Acanthiops</i>	sp. 2	138	25	37
Baetidae	<i>Acanthiops</i>	sp. 3	0	0	4
Baetidae	<i>Tanzaniops</i>	sp.	2	0	5
Baetidae	<i>Afroptilum</i>	sudafricanum	0	37	0
Baetidae	<i>Dabulamanzia</i>	sp.	0	0	7
Caenidae	<i>Caenis</i>	sp.	41	9	23
Leptophlebiidae	<i>Euthraulus</i>	sp.	0	11	7
Ephemerythidae	<i>Ephemerythus</i>	sp.	0	25	27
Total density (ind. m ⁻²)			2 852	267	595
Relative abundance (%)			76.8	7.2	16.0

sites ($F_{(2, 51)} = 134.37$, $p < 0.001$). Site 1 had the lowest mean 14.3 ± 0.28 $^{\circ}$ C, while site 2 had a mean of 21.9 ± 0.48 $^{\circ}$ C, which was the highest. Student Newman's Keul's multiple range test further showed that site 1 varied from sites 2 and 3 (Table 1). Water temperature was significantly different ($t = 0.027$; $df = 52$; $p < 0.5$) between the dry and wet season (Table 2).

Spatial DO levels were significantly different between sites ($F_{(2, 51)} = 131.14$, $p < 0.001$). Student Newman's Keul's multiple range test also revealed that DO at site 1 varied from sites 2 and 3 (Table 1). Seasonal variation in DO, however, was not significant ($t = 1.647$, $df = 46$, $p > 0.5$) (Table 2). Water pH was not significantly different among sites ($F_{(2, 51)} = 2.28$, $p > 0.5$).

Water conductivity was highly significant among the sites ($F_{(2, 51)} = 138.73$, $p < 0.001$). Site 1 had lowest conductivity $16.9 \pm 0.69 \mu\text{S cm}^{-1}$. It further increased to $93.5 \pm 5.37 \mu\text{S cm}^{-1}$ in site 3. Student Newman-Keuls multiple range test also revealed that the three sites were varied (Table 1). However, seasonal variation was not significant ($t = 0.563$; $df = 52$; $p > 0.5$) (Table 2). Spatial TDS differed significantly ($F_{(2, 51)} = 107.91$, $p < 0.001$). It was lowest in site 1 ($10.1 \pm 0.53 \text{ mg l}^{-1}$) and sharply increased to 47.9 mg l^{-1} at site 3 (Table 2). It was not statistically different ($t = 0.916$; $df = 52$, $p > 0.5$) (Table 2) between the dry and wet season.

Mayflies were present along the entire system, with lower abundances in the middle reaches. In total, 13 taxa of Ephemeroptera in six families were found during the entire sampling season (Table 3).

The major mayfly families in decreasing relative abundance and abundance were Baetidae and Heptageniidae. *Acanthiops* was the preponderant genus of Ephemeroptera in all the sampling sites. Oligoneuriidae was represented by *Oligoneuriopsis dobbsi* and occurred sporadically only in site 1. Leptophlebiidae family was poorly represented in the Thika River system. Families Caenidae, Heptageniidae, and Baetidae were distributed in all the sites. Interestingly, *Caenis*, *Euthraulus* and *Ephemerythus* sp. occurred together and were all absent in October and November when the rainy season was at its peak.

Distribution of mayflies differed significantly among sites ($H_{(2)} = 10.294$, $p < 0.5$, $\alpha = 0.5$; Table 3). The distribution of mayflies were ten times greater in site 1 (2852 ind. m^{-2}) compared to site 2 (267 ind. m^{-2}). *Acanthiops* dominated the mayfly community at all three sampling sites with 2369 ind. m^{-2} at site 1, 141 ind. m^{-2} at site 2 and 448 ind. m^{-2} at site 3. Other Ephemeroptera species were rare in the studied sites (Table 3). Seasonal variation in abundance was not significantly different ($H_{(5)} = 3.58$, $p > 0.5$, $\alpha = 0.5$), irrespective of sites (Figure 2). With respect to relative abundance, some of the correlations were weak while none of the correlations were significant (Table 4).

Discussion

Variations observed in background physico-chemical characteristics of the river are primarily governed by the local climatic conditions and catchment characteristics (Gibbs 1970; Arimoro and Muller 2010). Water temperature, DO, and the presence of appropriate mesohabitat structures are critical factors determining the occurrence and distribution of mayfly larvae (Bauernfeind and Moog 2000). The elevated levels of conductivity, TDS, temperature and low values of DO and transparency observed at sites 2 and 3 indicate deterioration of water quality impacted by several human activities taking place in these sites. Aura et al. (2010) recorded a similar trend in River Nzoia, Kenya, which receives sewage and urban effluents. Ideally, nutrient measures (PO_4^{3-} , NO_3^- and TP), although not measured in the current study, should be included in such a study.

Site 1 exhibited relatively clean water status, as was shown by the physico-chemical parameters in the section.




Figure 2: Relative abundance of Ephemeroptera genera of the Thika River at the forested site (a), cultivated site (b) and urban site (c) from September 2015 to February 2016

Sites 2 and 3 affected by discharges from agricultural and urban lands respectively, showed high values of conductivity, TDS, temperature, and is accompanied by

Table 4: Pearson correlation coefficient (r) of water parameters and abundance of Ephemeroptera of the Thika River during the study period, September 2015 to February 2016

	Temp (°C)	DO (mg l ⁻¹)	EC (µcm)	pH	TDS (mg l ⁻¹)	Transparency (m)	Depth (m)	Velocity (m s ⁻¹)	Abundance
Temp (°C)	1	-0.867**	0.400*	0.425**	0.497**	-0.787**	-0.804**	-0.865**	-0.233
DO (mg l ⁻¹)		1	-0.650**	-0.669**	-0.690**	0.425**	-0.404*	-0.583**	0.218
EC (µS cm ⁻¹)			1	0.385*	0.989**	0.156	-0.021	0.344*	-0.136
pH				1	0.326	-0.083	-0.054	0.233	-0.13
TDS (mg l ⁻¹)					1	0.037	0.113	0.446**	-0.154
Transparency (m)						1	-0.948**	-0.760**	0.159
Depth (m)							1	0.877**	-0.167
Velocity (m s ⁻¹)								1	-0.187
Abundance									1

* $p \leq 0.05$, ** $p \leq 0.001$

reduced values of DO and transparency. These values indicate that the water in the sites was impacted. The degree of silt-sand deposition was another factor that differentiated the sites. Human activities at sites 2 and 3 resulted in increased siltation that ultimately destroyed the microhabitats for the mayfly larvae. Arimoro and Muller (2010) found parallel results in River Orogodo (Nigeria) in sections exposed to anthropogenic disturbance. Studies carried out in most freshwater streams, show that mayflies are among the insect groups with the highest abundance in the macrobenthic community (Ogbeibu and Oribhabor 2002; Bonzemo 2013).

The total number of Ephemeropteran taxa obtained in the current study was high (13) when compared with other studies undertaken in different rivers in Kenya. Kage (2003) and Bonzemo (2013) reported three and nine taxa in Nairobi and Kisibi rivers, respectively. The occurrence may be as a result of the more extensive and more diverse meso-habitats in the Thika River, resulting in a higher mayfly abundance and diversity. A study with coinciding results was that of Arimoro and Muller (2010), who reported 13 taxa in River Orogodo in Nigeria.

The distribution of Ephemeroptera species along the Thika River was reasonably differentiated. The restricted presence of *Oligoneuriopsis dobbsi* in the forested site was predictable, because species in the genus had been recorded as dwellers of fast flowing streams and at high elevations (Day and Moor 2002). Bauernfeind and Moog (2000) found that Oligoneuriidae larvae are one of the families that indicate an ecologically intact environment. Site 1 showed a wide variety of microhabitats appropriate for mayflies. These may explain the high abundance of Ephemeroptera at the site. Baetidae family from the current study may be a potential indicator of water quality and ecological integrity primarily as a result of its presence in both the impacted and un-impacted reaches of the river. However, it appears to be sensitive to pollution as the numbers were remarkably reduced at the impacted sites (sites 2 and 3). Genus *Caenis* from several studies has been documented to be tolerant to organic contamination (Menetrey et al. 2008). The current study agrees with the findings because, *Caenis* was found in both impacted (sites 2 and 3) and unimpacted (site 1) sites, though in low relative abundance.

Ephemeroptera larvae are well represented in the Thika River, although they were considerably reduced in impacted sites. The highest abundances of mayfly was found in site 1. It has been found that surface waters bounded by agrarian lands have a higher conductivity when compared with other land uses (Detenbeck et al. 1996). Despite the high conductivity and TDS in site 3, signs of recovery were evidenced by the slightly reduced temperature, increased DO and the rise in mayfly abundance than in site 2. Lock and Goethals (2013) found that mayflies are always present in water with a low conductivity, but if the conductivity is high, they are only present when both the phosphate concentration and BOD are low.

Ephemeroptera abundance varied spatially, but not temporally, in response to physico-chemical factors of the water. The significantly lower abundance in sites 2 and 3 can be ascribed to combined influences of changes in substrate composition as a result of human activities at the sites, as well as deteriorated water quality at these sites. Change of sampling method from using a Hess sampler on the shallow rocky substrates to a grab sampler in the deeper soft muddy substrates may also account for the reduced abundances. Land use at Sites 2 and 3 were different from Site 1. At Sites 2 and 3, land use was heavily impacted by agricultural and urban activities, respectively, contributing to the change in species. The resulting food shortage led to a significant decrease in numbers of Ephemeroptera in the wet season. In the current study, wet season coincided with *El Niño* rains in 2015, which reached a climax of November. This is the reason why the month had the lowest abundance. Macroinvertebrates were utterly absent at sites 2 and 3, as a result of floods. The results were similar to those found by Arimoro and Muler (2010), who observed that the wet season had the least mean abundance of mayflies in river Orogodo in Nigeria.

The abundance of mayflies was highest in the upper reaches of the Thika River (site 1), but reduced drastically in site 2 and 3. The principal ecological stresses being land use and anthropogenic activities. However, considering the abundance of sites 2 and 3, the later had slightly higher abundance. The higher abundance on site 3 showed that Ephemeropterans could recover from environmental

Table 5: Levene's test for equality of variances for temperature, dissolved oxygen, conductivity, pH, total dissolved solids, transparency and width at the forested, cultivated and urban sampling sites

Site	t-test for equality of means									
	Levene's test for equality of variances					t-test for equality of means				
	F	Sig.	t	df	Sig. (two-tailed)	Mean difference	Std. error difference	95% confidence interval of the difference		
								Lower	Upper	
Forested	Temperature (°C)	0.138	0.715	5.226	16	0.000	1.81111	0.34659	1.07638	2.54585
				5.226	15.641	0.000	1.81111	0.34659	1.07500	2.54722
	Dissolved oxygen (mg l ⁻¹)	29.049	0.000	6.158	16	0.000	1.77111	0.28763	1.16137	2.38085
				6.158	8.289	0.000	1.77111	0.28763	1.11185	2.43037
	Conductivity (µS cm ⁻¹)	0.206	0.656	-4.239	16	0.001	-4.16667	0.98300	-6.25053	-2.08280
				-4.239	15.599	0.001	-4.16667	0.98300	-6.25489	-2.07844
	pH	4.881	0.042	1.562	16	0.138	0.19222	0.12308	-0.06869	0.45313
				1.562	13.614	0.141	0.19222	0.12308	-0.07245	0.45690
	Total dissolved solids (mg l ⁻¹)	0.172	0.684	-4.662	16	0.000	-3.29778	0.70740	-4.79739	-1.79816
				-4.662	15.925	0.000	-3.29778	0.70740	-4.79796	-1.79759
Transparency (m)	2.534	0.131	-2.573	16	0.020	-0.11444	0.04449	-0.20875	-0.02014	
			-2.573	11.676	0.025	-0.11444	0.04449	-0.21167	-0.01722	
Width (m)	4.580	0.048	-2.739	16	0.015	-2.50889	0.91605	-4.45083	-0.56695	
			-2.739	12.308	0.018	-2.50889	0.91605	-4.49927	-0.51850	
Depth (m)	0.181	0.676	-4.265	16	0.001	-3.04889	0.71482	-4.56423	-1.53354	
			-4.265	15.763	0.001	-3.04889	0.71482	-4.56609	-1.53169	
Cultivated	Temperature (°C)	5.143	0.038	5.008	16	0.000	3.06667	0.61235	1.76855	4.36478
				5.008	12.835	0.000	3.06667	0.61235	1.74204	4.39129
	Dissolved oxygen (mg l ⁻¹)	3.173	0.094	1.138	16	0.272	0.24889	0.21876	-0.21487	0.71264
				1.138	14.112	0.274	0.24889	0.21876	-0.21996	0.71774
	Conductivity (µS cm ⁻¹)	31.701	0.000	-0.655	16	0.522	-3.44444	5.25782	-14.59052	7.70163
				-0.655	8.920	0.529	-3.44444	5.25782	-15.35473	8.46585
	pH	5.344	0.034	-0.353	16	0.729	-0.07556	0.21416	-0.52955	0.37843
				-0.353	13.464	0.730	-0.07556	0.21416	-0.53659	0.38548
	Total dissolved solids (mg l ⁻¹)	12.408	0.003	-1.336	16	0.200	-3.89222	2.91295	-10.06739	2.28295
				-1.336	11.442	0.207	-3.89222	2.91295	-10.27345	2.48901
Transparency (m)	17.329	0.001	4.547	16	0.000	0.64111	0.14100	0.34221	0.94002	
			4.547	9.259	0.001	0.64111	0.14100	0.32350	0.95872	
Width (m)	7.582	0.014	-4.978	16	0.000	-2.12667	0.42722	-3.03233	-1.22101	
			-4.978	11.947	0.000	-2.12667	0.42722	-3.05795	-1.19538	
Depth (m)	4.680	0.046	-4.141	16	0.001	-0.46000	0.11108	-0.69548	-0.22452	
			-4.141	13.129	0.001	-0.46000	0.11108	-0.69973	-0.22027	

Table 5: (cont.)

Site	Levene's test for equality of variances		t-test for equality of means						
	F	Sig.	t	df	Sig. (two-tailed)	Mean difference	Std. error difference	95% confidence interval of the difference	
								Lower	Upper
Temperature (°C)	12.691	0.003	4.487	16	0.000	2.21111	0.49279	1.16645	3.25577
			4.487	9.147	0.001	2.21111	0.49279	1.09908	3.32314
Dissolved oxygen (mg l ⁻¹)	5.298	0.035	1.709	16	0.107	0.28556	0.16707	-0.06862	0.63973
			1.709	13.118	0.111	0.28556	0.16707	-0.07505	0.64616
Conductivity (µS cm ⁻¹)	32.544	0.000	-0.862	16	0.402	-9.32000	10.81788	-32.25288	13.61288
			-0.862	8.945	0.411	-9.32000	10.81788	-33.81490	15.17490
pH	34.728	0.000	-1.977	16	0.066	-0.31111	0.15737	-0.64472	0.02249
			-1.977	8.391	0.082	-0.31111	0.15737	-0.67108	0.04886
Total dissolved solids (mg l ⁻¹)	39.924	0.000	-1.122	16	0.278	-6.53889	5.82538	-18.88814	5.81036
			-1.122	8.491	0.292	-6.53889	5.82538	-19.83816	6.76039
Transparency (m)	1.988	0.178	3.592	16	0.002	0.27667	0.07703	0.11337	0.43997
			3.592	14.759	0.003	0.27667	0.07703	0.11224	0.44109
Width (m)	2.142	0.163	-5.470	16	0.000	-3.26000	0.59594	-4.52334	-1.99666
			-5.470	14.410	0.000	-3.26000	0.59594	-4.53476	-1.98524
Depth (m)	0.373	0.550	-2.869	16	0.011	-0.45111	0.15725	-0.78446	-0.11776
			-2.869	15.900	0.011	-0.45111	0.15725	-0.78463	-0.11759

stressors. The increase in abundance of mayflies at site 3 is in agreement to the intermediate disturbance hypothesis, where higher abundance and species diversity is expected in middle reaches, where disturbance is neither lacking nor too severe (Townsend et al. 1997).

Correlation between the abundance of mayfly larvae and water quality parameters measured (temperature, dissolved oxygen, electrical conductivity, TDS and transparency) was not statistically significant (Table 4), owing to the weak relationship between abundance and all the water parameters. This weak association mayfly abundance and the water quality parameters suggests that other external factors not measured may be controlling the abundance of mayfly larvae in the Thika River. Site 2 had slightly higher temperature, lower DO and the highly reduced abundance of mayflies, which suggest that the site was the most impacted. Water temperature, conductivity, pH, TDS and depth were negatively related to the abundance of mayflies. Many studies in different rivers have found a similar result. For instance, Arimoro and Muller (2010) found that Ephemeroptera abundance was negatively correlated with conductivity in River Orogodo in Nigeria.

In conclusion, the current study revealed that decreasing Ephemeroptera taxa abundance along the river channel indicates decreasing water quality. The Oligoneuriidae family was unique for site 1, which had good quality, natural water. Sites 2 and 3 showed a sharp decline in abundance of mayfly larvae, with species of the genus *Acanthiops* dominating in numbers. The dominance of *Acanthiops* highlights that this genus is tolerant to the forms of disturbances in the two sites. It was further revealed that Site 3 increased in DO and temperature that was accompanied by a greater increase in abundance of mayfly than at Site 2. More importantly, the site aquatic vegetation improved which promoted increased mayfly abundances. Kage (2003) also noted that increased abundance of aquatic vegetation in downstream sites favours higher mayfly abundance, as a result of increased habitat heterogeneity. Moreover, organic load dilution is known to occur downstream, generating a species composition and abundance similar to the upstream stations (Garcia-Amisen et al. 2014). Because the abundance of mayfly larvae decreased with a decline in water quality, it is evident that they prefer moderate to excellent water quality. Although the quantitative method is a good indicator of the water quality status at different sites, the South African Scoring System (SASS) may also be a good tool for the rapid bioassessment of water quality (Dallas 1995).

Acknowledgments — We acknowledge the Department of Zoological Sciences, Kenyatta University, Kenya for providing logistical, technical and laboratory support during the course of the current study. We also thank the Department of Zoology of the National Museums of Kenya, Insect Division for assisting in the identification of Ephemeroptera specimens.

ORCID

JJ Jumbe: <http://orcid.org/0000-0002-4993-2955>

References

- Arimoro FO, Muller WJ. 2010. Macroinvertebrates community and diversity relation to water quality status of river Ase, Niger Delta, Nigeria. *Journal of Fisheries and Aquatic Science* 2: 338–340.
- Arimoro FO, Ikomi RB, Iwegbue CM. 2007. Ecology and abundance of *Oligochaetes* as indicators of organic pollution in an urban stream in southern Nigeria. *Pakistan Journal of Biological Sciences* 10: 446–453.
- Aschalew L, Moog O. 2015. Benthic macroinvertebrates based new biotic score 'ETHbios' for assessing ecological conditions of highland streams and rivers in Ethiopia. *Limnologia* 52: 11–19. <https://doi.org/10.1016/j.limno.2015.02.002>.
- Aura CM, Raburu PO, Herrmann J. 2010. A preliminary macroinvertebrate Index of Biotic Integrity for bioassessment of the Kipkaren and Sosiani Rivers, Nzoia River basin, Kenya. *Lakes & Reservoirs. Research and Management* 15: 119–128.
- Barber-James HM, Lugo-Ortiz CR. 2003. Ephemeroptera. In: de Moor IJ, Day JA, de Moor FC (Eds.), *Guide to the Freshwater Invertebrates of Southern Africa*. Pretoria: Water Resource Commission. pp. 16–159
- Bauernfeind E, Moog O. 2000. Mayflies (Insecta: Ephemeroptera) and the assessment of ecological integrity: a methodological approach (pp 71–83). In: Jungwirth M, Muhar S, Schmutz S (Eds.). *Assessing the Ecological Integrity of Running Waters*. Amsterdam: Kluwer Academic Publishers. pp 71–83.
- Bonzemo SW. 2013. Assessment of water quality status of River Kibisi, Kenya using the Ephemeroptera, Plecoptera, and Trichoptera (EPT) Index. Msc thesis, Kenyatta University, Kenya. <http://library.ku.ac.ke/handle/123456789/9206>.
- Dallas HF. 1995. An evaluation of SASS (South African Scoring System) as a tool for the rapid bioassessment of water quality. MSc thesis, University of Cape Town, South Africa.
- Day JA, De Moor IJ. 2002. Guides to the freshwater invertebrates of southern Africa: Volume 5 Non-Arthropods–The Protozoans. Porifera, Cnidaria, Platyhelminthes, Nemertea, Rotifera, Nematoda, Nematomorpha, Gastrotrichia, Bryozoa, Tardigrada, Polychaeta, Oligochaeta and Hirudinea. WRC Report No. TT. 167: 13–49. Pretoria: Water Research Commission.
- DeSA UN. 2013. World population prospects: the 2012 revision. New York. Population Division of the Department of Economic and Social Affairs of the United Nations Secretariat. New York. <https://www.un.org/en/development/desa/publications/world-population-prospects-the-2012-revision.html>. [Accessed 14 February 2020]
- Detenbeck NE, Taylor DL, Lima A, Hagley C. 1996. Temporal and spatial variability in water quality of wetlands in the Minneapolis/St. Paul, MN metropolitan area: implications for monitoring strategies and designs. *Environmental Monitoring and Assessment* 40: 11–40.
- Garcia L, Pardo I, Delgado C. 2014. Macroinvertebrate indicators of ecological status in Mediterranean temporary stream types of the Balearic Islands. *Ecological Indicators* 45: 650–663. <https://doi.org/10.1016/j.ecolind.2014.05.030>.
- García-Armisen T, İnceoğlu Ö, Ouattara NK, Anzil A, Verbanck MA, Brion N, et al. 2014. Seasonal variations and resilience of bacterial communities in sewage polluted urban river. *PLoS ONE* 9: e92579. <https://doi.org/10.1371/journal.pone.0092579>.
- Gibbs RJ. 1970. Mechanisms controlling world water chemistry. *Science* 170: 1088–1090.
- Heathwaite AL, Johnes PJ, Peters NE. 1996. Trends in nutrients. *Hydrological Processes* 10: 263–293.
- Hubbard MD. 1990. *Mayflies of the World: a Catalog of the Family and Genus Group Taxa (Insecta: Ephemeroptera)* (No. 8). London: CRC Press, Inc.
- Huxley T. 2003. Provisional atlas of the British aquatic bugs (Hemiptera, Heteroptera). Centre for Ecology and Hydrology

- Biological Records Centre. *Cambridge Journal* 23: 118–128.
- Inoti KJ, Fanuel K, George O, Paul O. 2012. Assessment of heavy metal concentrations in urban-grown vegetables in Thika Town, Kenya. *African Journal of Food Science* 6: 41–46.
- Kage FG. 2003. Impact of heavy metal pollution on the composition and abundance of benthic macroinvertebrates along Nairobi River, Kenya. MSc thesis, Kenyatta University, Kenya.
- Karuri AW, Wamicha W, Maina D, Bartilol SK. 2003. Studies on the influences of land use on soil and water resources in Thika district, Kenya (pp 8). In: Mureithi JG, Macharia PN. Proceedings of the 18th Conference and End of the Millennium Celebrations 4-8 December 2000, Mombasa (Kenya), 4–8 December 2000. Kenya: Soil Science Society of East Africa.
- Kimani PK, Thiong'o GT, Mwangi JK. 2016. Spatial and seasonal variation of selected water quality parameters in Chania River Catchment, Kenya. *British Journal of Applied Science & Technology* 18: 1–16.
- Kotchoni DOV, Vouillamoz J-M, Lawson FMA, Adjomayi P, Boukari M, Taylor RG. 2019. Relationships between rainfall and groundwater recharge in seasonally humid Benin: a comparative analysis of long-term hydrographs in sedimentary and crystalline aquifers. *Hydrogeology Journal* 27: 447–457. <https://doi.org/10.1007/s10040-018-1806-2>.
- Lock, K, Goethals, PL. 2013. Habitat suitability modeling for mayflies (Ephemeroptera) in Flanders (Belgium). *Ecological Informatics* 17: 30–35. <https://doi.org/10.1016/j.ecoinf.2011.12.004>.
- Menetrey N, Oertli B, Sartori M, Wagner A, Lachavanne JB. 2008. Eutrophication: Are mayflies (Ephemeroptera) good bioindicators for ponds? *Hydrobiologia* 597: 125–135. <https://doi.org/10.1007/s10750-007-9223-x>.
- Merritt RW, Cummins KW. 1996. *An Introduction to the Aquatic Insects of North America*. 3rd edn. Dubuque: Kendall/Hunt Pub. Co.
- Ndaruga AM, Ndiritu GG, Gichuki NN, Wamicha WN. 2004. Impact of water quality on macroinvertebrate assemblages along a tropical stream in Kenya. *African Journal of Ecology* 42: 208–216. <https://doi.org/10.1111/j.1365-2028.2004.00516.x>.
- Ndungu JN. 2014. Assessing Water Quality in Lake Naivasha. PhD thesis, University of Twente, The Netherlands. <https://doi.org/10.3990/1.9789036537001>.
- Ogbeibu AE, Oribhabor BJ. 2002. Ecological impact of river impoundment using benthic macro-invertebrates as indicators. *Water Research* 36: 2427–2436. [https://doi.org/10.1016/S0043-1354\(01\)00489-4](https://doi.org/10.1016/S0043-1354(01)00489-4).
- Schloesser DW, Nalepa TF. 2002. Comparison of 5 benthic samplers to collect burrowing mayfly nymphs (*Hexagenia* spp.: Ephemeroptera: Ephemeridae) in sediments of the Laurentian Great Lakes. *Journal of the North American Benthological Society* 21: 487–501. <https://doi.org/10.2307/1468485>.
- Townsend, CR, Scarsbrook, MR and Doledec, S. 1997. The intermediate disturbance hypothesis, refugia and biodiversity in streams. *Limnology and Oceanography* 42: 938–949.
- Wetzel RG, Likens G. 2013. *Limnological Analysis*. New York: Springer Science & Business Media.