



## **Antibiotic Susceptibility Profile of Bacteria Isolated from Kenyan Bank Notes Circulating in Nyeri Town**

**S. G. Muguongo<sup>1\*</sup>, A. K. Nyamache<sup>1</sup> and J. M. Maingi<sup>1</sup>**

<sup>1</sup>Department of Microbiology, Kenyatta University, P.O. Box 43844-00100, Nairobi, Kenya.

### **Authors' contributions**

This work was carried out in collaboration among all authors. Author SGM did the sample collection, isolation, biochemical identification, drug susceptibility testing of different isolates from Kenyan bank notes, analyzed data and prepared the first draft, author AKN provided the working protocols for biochemical identification (Supervised) the activity, author JMM provided general guidance on the experiment and edited the manuscript. All the authors read and approved the manuscript.

### **Article Information**

DOI: 10.9734/MRJI/2019/v28i230125

#### Editor(s):

(1) Dr. Essam Hussein Abdel-Shakour, Professor, Department of Botany and Microbiology, Faculty of Science, Al-Azhar University, Cairo, Egypt.

#### Reviewers:

(1) Oshim, Ifeanyi Onyema, Nnamdi Azikiwe University, Nigeria.

(2) Mabel Aworh, Ahmadu Bello University, Nigeria.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/49596>

**Original Research Article**

**Received 05 April 2019**

**Accepted 20 June 2019**

**Published 25 June 2019**

### **ABSTRACT**

**Aims:** The aim of this study was to characterize bacteria isolated from circulating Kenyan banknotes and also antibiotic susceptibility profiles within Nyeri County.

**Study Design:** This was a cross-sectional study and simple random sampling was used to collect 25 of each paper currency denomination.

**Place and Duration of Study:** Samples analyses were done at Outspan Teaching and Referral Hospital (OTRH) laboratory, between March, 2019 and April, 2019.

**Methodology:** Total of 125 currencies of five different denominations were collected from different marketing sources such as Butcheries, Restaurants, Health facilities, Mpesa outlets and Transport Saccos and dropped in sterile bags. The bacterial isolates were characterized on the basis of their morphology, staining and biochemical tests. Antibiotic sensitivity tests were done by Kirby Bauer disc diffusion technique.

**Results:** Total of 19 different bacterial species were isolated from five Kenyan Bank note currencies. Of these, 37 (52.2%) was *Staphylococcus aureus* followed by *Staphylococcus sciuri* ssp.*lentus* 7 (9.9%), *Staphylococcus gallinarum* 2 (2.8%), *Staphylococcus intermedius* 6 (8.5%),

\*Corresponding author: E-mail: [sgmuguongo@gmail.com](mailto:sgmuguongo@gmail.com);

*Micrococcus sp.* 1 (1.4%), *Staphylococcus schleiferi ssp.coagulans* 2 (2.8%), *Staphylococcus sciuri ssp.rodentium* 1 (1.4%), *Kluyvera ascorbata* 1 (1.4%), *Proteus penneri* 1 (1.4%), *Aeromonas media* 3 (4.2%), *Burkholderia cepacia ssp.komplex* (1.4%), *Aeromonas enteropelogenes* 1 (1.4%), *Enterobacter cloacae* 1 (1.4%), *Klebsiella oxytoca* 2 (2.8%), *Leclercia adecarboxylata* 1 (1.4%), *Raoultella ornithinolytica* 1 (1.4%), *Vibrio metschnikovii* 1 (1.4%), *Myroides odoratus* 1 (1.4%) and *Yersinia pestis* 1 (1.4%). Overall gram positive and gram negative bacterial isolates exhibited resistance to vancomycin, clindamycin and amoxicillin with percentages 40 (71%), 28 (50%), and 37 (66%) and 9 (64%), 8 (57%) and 6 (43%) respectively.

**Conclusion:** This study revealed that Kenyan banknote currencies circulating in Nyeri County were contaminated with different pathogenic and potential pathogenic bacteria including multi drug resistant strains. Hence, great care must be taken while handling money during the preparation and handling of food to avoid cross contamination.

**Keywords:** Antibiotic; susceptibility; Kenyan; banknotes; contamination.

## ABBREVIATIONS

OTRH	: Outspan Teaching and Referral Hospital
Shs	: shillings
S	: Sensitive
R	: Resistant
CRO	: Ceftriaxone
TE	: Tetracycline
AML	: Amoxicillin
CIP	: Ciprofloxacin
CN	: Gentamycin
CD	: Clindamycin
VA	: Vancomycin
E	: Erthromycin.

## 1. INTRODUCTION

Money is any medium of exchange that is widely accepted in payment for goods, services and in settlements of debts. Paper currency is widely exchanged for goods and services in countries worldwide [1]. It also serves as a standard of value for measuring the relative worth of goods and services [1]. Modern scientific techniques have confirmed these theories and have shown that viable pathogenic organisms (viruses, bacteria, and fungi) can be isolated on the surfaces of both paper and coin currency [2].

Contamination of materials by pathogenic microorganisms is of public health concern as contaminated materials could be a source of transmitting microbial pathogens [3]. Movement of materials from hand to hand makes it more prone to contamination by pathogenic organisms. Polymer currency therefore, poses a serious threat to public health since communicable diseases could also be contracted through fomites [2]. Currency is handled by all

categories of people and may be contaminated during coughing, sneezing, touching with hands and placement on dirty surfaces. Many people tongue-wet their fingers when counting money and contaminate their fingers as well as currency notes. So, it is obvious that gets on hands may be transferred to money and vice-versa [4]. Paper banknotes have a large surface area for bacterial attachment and would be a vector for transmission of potentially pathogenic microorganisms between populations [5]. The risk of microbial transfer by paper currency is influenced by factors such as paper value and duration of usage [6].

Studies on the persistence of pathogens on paper currency showed that currency notes are considered as a potential cause of food-borne diseases [7]. There are evidences of isolation of food-borne pathogens including *Salmonella spp*, *E. coli*, *S. aureus*, *P. aeruginosa*, *enterococci spp* and *Serratia marcescens* from the banknotes of different countries [8]. Various pathogens which may cause throat infection, pneumonia, peptic ulcers, tonsillitis, urino-genital tract infections, gastro enteritis and lung abscess had been reported [9]. According to a study 100% notes were contaminated with *E. coli*, *Klebsiella pneumonia*, *Pseudomonas aeruginosa* and *Staphylococcus aureus* [2] and similar bacteria also found on the currency notes of Coimbatore city, Tamil Nadu [10]. Orukotan and Yabaya [11] also surveyed naira notes, comprising of all the denominations for microbial contamination in Kaduna metropolis. The microorganisms recovered from these notes included *Escherichia coli*, *Bacillus*, *Salmonella*, *Streptococcus*, *Staphylococcus aureus*, *Proteus*, *Klebsiella*, *Micrococcus*, *Fusarium*, *Penicillium*, *Aspergillus* and *Rhizopus*.

Antimicrobial resistance capabilities of microorganisms have become a major public health concern in many regions of the world [12, 13]. So investigation of the situation of antibiotic resistance capabilities in bacteria is essential to gauge the level of threat. Knowledge of the microbial diversity of currency notes in circulation can provide the basis for raise health consciousness in people during currency handling and effective control of infection transmission. The aim of this study was to characterize bacteria isolated from circulating Kenyan banknotes and also antibiotic susceptibility profiles within Nyeri County.

## 2. MATERIALS AND METHODS


### 2.1 Study Area

Nyeri County is a county in the central region of Kenya. Nyeri town is the capital and largest town

is Nyeri County. It has a population of 661,156 and an area of 3,356 km<sup>2</sup> making it one of the most densely populated areas in Kenya.

### 2.2 Sample Collection and Transport

The study currency notes were collected during period between March, 2019 and April, 2019. The control sample bank notes were collected at random from the tellers in the Central Bank Nyeri. The experimental sample notes were collected from different marketing sources such as Butcheries, Restaurants, Health facilities, Mpesa outlets and Transport Saccos. To collect the currency notes, the individuals were asked to drop the currency into a sterile zipped plastic packet, which were sealed and immediately transported to the Outspan Teaching and Referral Hospital (OTRH) laboratory for microbial analysis [9].



Map 1. Map showing Nyeri municipality (highlighted in black) in Nyeri County, Kenya

## 2.3 Study Design

This was a cross-sectional study and simple random sampling was used to collect 25 of each paper currency denomination.

## 2.4 Sample Size

The currency notes studied were fifty, one hundred, two hundred, five hundred and one thousand Kenyan shillings notes. The study had a total sample size of 125 bank notes and five control bank notes, one from every denomination.

## 2.5 Isolation of Microbes

The currency notes were dipped in sterile normal saline and vigorously shaken for 3 minutes. A sterile cotton swab was dipped and inoculated in blood agar and Mac Conkey agar for each note. The plates were incubated at 37°C for 18-24 hours. After 18-24 hours the plates were observed for bacterial colonies.

## 2.6 Morphological and Biochemical Characterization of the Isolates

The bacterial isolates were characterized on the basis of their morphology, staining and biochemical tests. Gram staining was done as described by Barrow and Feltham [14]. All isolated microorganisms were subjected to microscopic examination and the shape, arrangement and Gram's reaction were detected and recorded. For biochemical test, the study used cypress diagnostic Bacterial Identification System for both gram positive and Gram negative consisting of 24 miniaturized biochemical tests.

## 2.7 Susceptibility Studies on the Bacteria Isolates

The Kirby-Bauer Disc Diffusion Method (Struve et al. [41]) was used to test the in vitro susceptibility of the identified isolates to Ceftriaxone 30 µg, Tetracycline 30 µg, Amoxicillin 30 µg, Ciprofloxacin 5 µg, Gentamycin 10 µg, Clindamycin 2 µg, Vancomycin 30 µg and Erythromycin 15 µg. A sterile platinum loop was used to pick overnight bacterial colonies from the culture plate and emulsified in 4 ml of sterile peptone water to match with 0.5 McFarland turbidity standards (1.0x10<sup>8</sup> cfu/ml). Using a sterile swab, the

surface of Mueller Hinton agar (Oxoid, Basingstoke, UK) was evenly inoculated with the suspension and allowed to air dry for 10 minutes. Using multichannel disc dispenser (Oxoid, Basingstoke, UK) antibiotics discs were deposited onto the surface of the inoculated medium and plates incubated at 37°C for 24 hours. The exercise was replicated and the results compared with chart provided by the Clinical and Laboratories Standards Institute. *E. coli* (ATCC 25922) and *S. aureus* (ATCC 25923) were used as control. Susceptibility of bacterial isolates was recorded as 'sensitive' or 'resistant' according to Clinical and Laboratory Standards Institute guideline [15].

## 2.8 Data Analysis

Descriptive statistics was used to analyze various data from the laboratory. These included averages, percentages and frequencies. Continuous data were expressed as means and categorical data expressed as proportions. Statistical analysis was performed using statistical package for social sciences (SPSS) software for Windows, ver. 21 (SPSS, IBM, USA).

## 2.9 Ethical Considerations

The research proposal was submitted to Kenyatta University and Outspan Teaching and Referral Hospital (OTRH) for approval to conduct the research.

## 3. RESULTS AND DISCUSSION

### 3.1 Types of Bacterial Contaminants in Bank Notes

From the study 69 (55.2%) were dirty and 56 (44.8%) were clean. 100% of the currency notes used as control from Nyeri Central Bank were mint. It is worth noting that bacterial growth was not detected in 5 samples of mint "newly printed" banknotes. Lack of growth in these notes might be attributed to the fact that they had not been in circulation that exposed them to usage and handling. However, some researchers believed that uncirculated notes are contaminated with fastidious organisms and the media or culture conditions employed were inappropriate for their isolation [16].

Out of the 125 currency notes of five different denominations obtained from the five sources, 110 (88%) in blood agar and 95 (76%) in

MacConkey showed growth in plates; whereas all (5) notes obtained from the bank were sterile. A total of 71 isolates were obtained from contaminated currency notes with 56 (78.9%) being Gram's positive and 15 (21.1%) being Gram's negative. In the present study, the isolation of Gram's positive as well as Gram's negative bacteria from currency notes confirmed that currency might be playing an important role as a vector in the transmission of pathogenic bacteria in the community. In the current study, the identification and enumeration of various types of pathogenic microorganisms that were obtained from the Kenyan banknotes were contaminated with some strains of the pathogen bacteria. These results were compatible with previous researchers from other countries which elucidated that currency banknotes are usually contaminated by pathogenic microorganisms [17, 18].

Of the 71 isolates, 56 (78.9%) were Gram positive bacteria isolated from banknotes: *Staphylococcus aureus* 37 (66.1%), *Staphylococcus sciuri ssp.lentus* 7 (12.5%), *Staphylococcus intermedius* 6 (10.6%), *Staphylococcus schleiferi ssp.coagulans* 2 (3.6%), *Micrococcus sp.* 1 (1.8%), *Staphylococcus gallinarum* 2 (3.6%) and *Staphylococcus sciuri ssp.rodentium* 1 (1.8%) as shown in Fig. 1. Different species of bacteria isolated in this current study are almost similar to the studies done in Saudi Arabia, Pakistan, Ghana, Nigeria, and US [19,20,21,22]. All these studies established gram positive bacteria as the

major isolates from the contaminated currencies which agrees with our current study.

The most common Gram positive bacterial isolates from this study was *Staphylococcus aureus* 37 (66.1%). This occurrence is fully supported by similar researched work on microbiological evaluation of naira notes handled by fish sellers in umuahia metropolis and bacterial contamination of Nigerian currency notes and associated risk factors [23].

*Staphylococcus aureus* is commonly present on the skin and in the nasal passage of human and its presence in paper currency is also abundant [22]. The coagulase-negative staphylococci are part of the normal human flora and sometimes cause infections such as food poisoning [22] and other diseases often associated with implanted appliances and devices [22], especially in very young, old, and immunocompromised patients. Though *Staphylococcus aureus* are part of the normal flora of the skin and mucous membrane, their high incidence has clinical significance and they are considered well-recognized pathogen. A number of studies have documented the clinical significance of *S. aureus* as a causative agent of urinary tract infections [22]. *S. aureus* is also associated with toxic shock syndrome, skin infections e.g. frunculosis and respiratory tract infections. From this study, the bacterial isolates that were isolated from this study were associated with oral, nasal and skin contamination. Contamination of different objects by potential pathogenic microorganisms is a





Fig. 1. Gram positive bacteria isolated from Kenyan bank notes circulating in Nyeri Town between March, 2019 and April, 2019



**Fig. 2. Gram negative bacteria isolated from Kenyan bank notes circulating in Nyeri Town between March, 2019 and April, 2019**

serious concern of public health because items that pass from one hand to another gives the opportunity of contamination with wide range of pathogenic microorganisms [24].

Of the 71 isolates, 15 (21.1%) Gram negative bacteria isolated from Kenyan bank notes include: *Burkholderia cepacia* ssp.komplex 1 (6.67%), *Aeromonas enteropelogenes* 1 (6.67%), *Kluyvera ascorbata* 1 (6.67%), *Aeromonas media* 3 (20.0%), *Raoultella ornithinolytica* 1 (6.67%), *Enterobacter cloacae* 1 (6.67%), *Klebsiella oxytoca* 2 (13.3%), *Leclercia adecarboxylata* 1 (6.67%), *Vibrio metschnikovii* 1 (6.67%), *Proteus penneri* 1 (6.67%), *Myroides odoratus* 1 (6.67%) and *Yersinia pestis* 1 (6.67%) as shown in Fig. 2.

The dominant gram negative bacterial isolates from this study was *Aeromonas media* 3 (20.0%) followed by *Klebsiella oxytoca* (13.3%) as shown by Fig. 2. A study by Elumalai et al. [25] isolated eight different types of bacterial species *E. coli*, *Proteus mirabilis*, *Vibrio spp.*, *S. aureus*, *Pseudomonas spp.*, *Salmonella spp.*, *Bacillus spp.*, and *Klebsiella spp.* from 30 Indian currency notes consisting of five notes each of Indian Rupee  $\neq$  5 and  $\neq$  10 denominations. The current study agrees with a study done by Ahmed et al. [26] in India that, found *Proteus* sp. to be one of the predominant organisms isolated from contaminated currency.

Table 1, Shows that total of 19 different bacterial species were isolated from five Kenyan bank note currencies. Of these, 37 (52.2%) was *Staphylococcus aureus* followed by *Staphylococcus sciuri* ssp.lentus 7 (9.9%), *Staphylococcus gallinarum* 2 (2.8%), *Staphylococcus intermedius* 6 (8.5%), *Micrococcus* sp. 1 (1.4%), *Staphylococcus schleiferi* ssp.coagulans 2 (2.8%), *Staphylococcus sciuri* ssp.rodentium 1 (1.4%), *Kluyvera ascorbata* 1 (1.4%), *Proteus penneri* 1 (1.4%), *Aeromonas media* 3 (4.2%), *Burkholderia cepacia* ssp.komplex (1.4%), *Aeromonas enteropelogenes* 1 (1.4%), *Enterobacter cloacae* 1 (1.4%), *Klebsiella oxytoca* 2 (2.8%), *Leclercia adecarboxylata* 1 (1.4%), *Raoultella ornithinolytica* 1 (1.4%), *Vibrio metschnikovii* 1 (1.4%), *Myroides odoratus* 1 (1.4%) and *Yersinia pestis* 1 (1.4%). This current study agrees with the study by Tagoe et al. [20], that *staphylococcus* is the most observed isolate from currency notes. Studies in different parts of India show that predominant organisms isolated from contaminated currency were *Bacillus* sp followed by Coagulase negative Staphylococci (CNS) and *Micrococcus* sp [27]. Orukotan and Yabaya [11] also surveyed naira notes, comprising of all the denominations for microbial contamination in Kaduna metropolis. The microorganisms recovered from these notes included *Escherichia coli*, *Bacillus*, *Salmonella*, *Streptococcus*, *Staphylococcus aureus*, *Proteus*, *Klebsiella*,

**Table 1. Relative occurrence of bacterial species on bank notes currency of different denominations circulating in Nyeri Town between March, 2019 and April, 2019**

Currency Denomination (Ksh)	50	100	200	500	1000	Number (%)
No. of currency	N = 25					
<i>Staphylococcus aureus</i>	10	6	5	8	8	37 (52.2%)
<i>Staphylococcus sciuri ssp.lentus</i>	2	2	2	1	0	7 (9.9%)
<i>Staphylococcus gallinarum</i>	1	0	1	0	0	2 (2.8%)
<i>Staphylococcus intermedius</i>	0	2	0	3	1	6 (8.5%)
<i>Micrococcus sp.</i>	0	0	1	0	0	1 (1.4%)
<i>Staphylococcus schleiferi ssp.coagulans</i>	1	0	0	0	1	2 (2.8%)
<i>Staphylococcus sciuri ssp.rodentium</i>	0	0	1	0	0	1(1.4%)
<i>Kluyvera ascorbata</i>	0	1	0	0	0	1(1.4%)
<i>Proteus penneri</i>	1	0	0	0	0	1(1.4%)
<i>Aeromonas media</i>	1	0	1	0	1	3 (4.2%)
<i>Burkholderia cepacia ssp.komplex</i>	1	0	0	0	0	1(1.4%)
<i>Aeromonas enteropelogenes</i>	0	1	0	0	0	1(1.4%)
<i>Enterobacter cloacae</i>	1	0	0	0	0	1(1.4%)
<i>Klebsiella oxytoca</i>	0	1	0	1	0	2 (2.8%)
<i>Leclercia adecarboxylata</i>	0	1	0	0	0	1(1.4%)
<i>Raoultella ornithinolytica</i>	0	0	1	0	0	1(1.4%)
<i>Vibrio metschnikovii</i>	0	0	0	1	0	1(1.4%)
<i>Myroides odoratus</i>	0	0	0	0	1	1(1.4%)
<i>Yersinia pestis</i>	1	0	0	0	0	1(1.4%)
Total	19	14	12	14	12	71(100%)

*Micrococcus*, *Fusarium*, *Penicillium*, *Aspergillus* and *Rhizopus*. Ahmed et al. [26] suggested that the Bangladesh paper currency commonly contaminated with pathogenic microorganisms and this contamination may play a significant role in the transmission of potentially harmful microorganisms or different diseases such as cholera, diarrhea, skin infections and also poses antibiotic resistant. The presence of these bacterial contaminant may cause a wide variety of diseases from food poisoning, wound and skin infections, respiratory and gastrointestinal problems to life threatening diseases such as meningitis and septicemia [28]. Paper notes of currency which is handled by a large number of people increase the possibility of acting as environmental vehicle for the transmission of potential pathogenic microorganisms [4].

Mohamed et al. [29] in a study of bacterial contamination on Libyan paper banknotes in circulation found *Enterobacter cloacae* (11%), *Klebsiella pneumonia* and *Enterobacter, Kluyvera spp* (4%) which is a lower percentage from our current study that found *Enterobacter cloacae* (1.4%), *Klebsiella oxytoca* (2.8%), *Kluyvera ascorbata* (1.4%). The presence of these pathogenic bacteria in this current study reveals that the majority of people are exposed to contaminated currency notes. Keeping money in dirty places and as a habit, wetting fingers with

saliva while counting currency notes suggests that humans are the major source of microorganisms on currency. As damaged or soiled notes are contaminated, they are particularly dangerous to health. Additionally, unwashed fingers contained many microorganisms, of which could be transient or resident [4]. These practices, including indiscriminate coughing, sneezing and defecation with indecent handling of currency notes were the most common sources of contamination [3, 9]. Furthermore, the materials of which the currency was manufactured are probably a factor that affects the survival of microorganisms on the banknotes [2].

Generally, lower value denomination currencies 50sh and 100sh were more contaminated with bacterial species than higher value denomination ones like 500sh and 1000sh. The current study agrees with a study by Yakubu et al. [30] that, currency notes of lower denominations were the most contaminated, presumably because lower denomination notes pass through more hands in their lifetime than the higher denomination notes. A study by Pavani and Srividya [31] established that most prevalent contamination (100%) was found among the Rupees 10 notes and coins and least prevalent contamination was found in Rupees 50 and 100. The denomination notes which receive most handling and exchanged

**Table 2. Antibiotic susceptibility patterns of gram positive bacterial isolates (No. of R or S isolates / n)**

Bacteria	CRO	TE	AML	CIP	CN	CD	VA	E
<i>Staphylococcus aureus</i> (n=37)	S (36/37)	S (35/37)	R (20/37)	S (36/37)	S (37/37)	R (24/37)	R (26/37)	S (21/37)
<i>Staphylococcus intermedius</i>	S (5/6)	S (6/6)	R (4/6)	S (6/6)	S (6/6)	R (4/6)	R (5/6)	R (3/6)
<i>Staphylococcus sciuri</i> ssp. <i>lentus</i>	S (7/7)	S (7/7)	S (6/7)	S (7/7)	S (6/7)	R (4/7)	S (5/7)	S=(5/7)
<i>Staphylococcus gallinarum</i>	S (2/2)	S (2/2)	R (2/2)	R (2/2)	S (2/2)	R (2/2)	R (2/2)	S (2/2)
<i>Staphylococcus sciuri</i> ssp. <i>rodentium</i>	S (1/1)	R (1/1)	R (1/1)	S (1/1)				
<i>Micrococcus</i> sp.	S (1/1)	S (1/1)	R (1/1)	S (1/1)	S (1/1)	R (1/1)	R (1/1)	R (1/1)
<i>Staphylococcus schleiferi</i> ssp. <i>coagulans</i>	S (2/2)	S (2/2)	S (2/2)	S (2/2)	R (1/2)	R (1/2)	R (1/2)	S (2/2)

S=Sensitive, R=Resistant, CRO= Ceftriaxone, TE= Tetracycline, AML= Amoxycilin, CIP= Ciprofloxacin, CN= Gentamycin, CD= Clindamycin, VA= Vancomycin, E= Erthromycin. (Clinical and Laboratory Standards Institute guideline, 2014)

**Table 3. Antibiotic susceptibility patterns of Gram negative bacterial isolates (No. of R or S isolates/ n)**

Bacteria	CRO	TE	AML	CIP	CN	CD	VA	E
<i>Kluyvera ascorbata</i>	S (1/1)	R (1/1)	S (1/1)	S (1/1)				
<i>Proteus penneri</i>	S (1/1)							
<i>Aeromonas media</i>	S (3/3)	S (3/3)	S (2/3)	S (3/3)	S (3/3)	S (2/3)	R (2/3)	S (3/3)
<i>Burkholderia cepacia</i> ssp. <i>komplex</i>	S (1/1)	R (1/1)	S (1/1)					
<i>Aeromonas enteropelogenes</i>	S (1/1)	S (1/1)	R (1/1)	S (1/1)	S (1/1)	R (1/1)	R (1/1)	R (1/1)
<i>Enterobacter cloacae</i>	S (1/1)	S (1/1)	R (1/1)	S (1/1)	S (1/1)	R (1/1)	S (1/1)	R (1/1)
<i>Klebsiella oxytoca</i>	S (2/2)	S (2/2)	R (2/2)	S (2/2)	S (2/2)	R (2/2)	R (2/2)	R (1/2)
<i>Leclercia adecarboxylata</i>	S (1/1)	S (1/1)	R (1/1)	S (1/1)	S (1/1)	R (1/1)	R (1/1)	S (1/1)
<i>Raoultella ornithinolytica</i>	R (1/1)	S (1/1)	R (1/1)	S (1/1)				
<i>Vibrio metschnikovii</i>	S (1/1)	R (1/1)	R (1/1)	S (1/1)				
<i>Myroides odoratus</i>	S (1/1)	R (1/1)	S (1/1)					

S=Sensitive, R=Resistant, CRO= Ceftriaxone, TE= Tetracycline, AML= Amoxycilin, CIP= Ciprofloxacin, CN= Gentamycin, CD= Clindamycin, VA= Vancomycin, E= Erthromycin. (Clinical and Laboratory Standards Institute guideline, 2014)

many times are more prone for contamination than other notes. Similar results were stated by Azza et al. [32] that found large denominations for their savings either at home or in banks which may keep them away from hand contamination for a period of time.

### 3.2 Antibiotic Susceptibility Testing

Table 2, shows antibiotic susceptibility patterns of gram positive bacterial isolates. The current studies reveal many multidrug resistant bacteria like *Staphylococcus aureus*, *Staphylococcus intermedius* and *Micrococcus sp.* to Amoxycillin, Clindamycin and Vancomycin. This current study agrees with a study done by Felgo and Nkansah [33] who found multidrug resistant bacteria prevalent in the currency that included methicillin resistant *Staphylococcus aureus*, methicillin resistant coagulase negative staphylococci, multi drug resistant *Escherichia coli* and *Klebsiella sp.* Srinu et al. [34] also reported that *S. aureus* was sensitive to Streptomycin, cotrimoxazole and Ciprofloxacin which concur with the current study that *S. aureus* is sensitive to Ciprofloxacin.

The study found that 54 (96.0%), 55 (98.0%) and 55 (98.0%) of gram positive isolates were susceptible to ceftriaxone, tetracycline and gentamycin respectively. Table 2 shows resistance rates of all bacterial isolates; overall isolates exhibited resistance to vancomycin, amoxycillin and clindamycin with percentages 40 (71%), 28 (50%), and 37 (66%) respectively. On another hand, isolates of all bacterial species showed absent or little resistance rates against antibiotics like ceftriaxone, tetracycline, gentamycin, ciprofloxacin and Erthromycin that were 2 (3.5%), 2 (3.5%), 2 (3.5%), 3 (7%) and 20 (35.7%) respectively. It is known that infection by multidrug-resistant bacteria limit therapeutic options and subsequently facilitate the dissemination of these strains. Paper currency notes collected from meat sellers in market places of Tanga city of Tanzania shows that 28.125% *S. aureus* isolates were multidrug resistant. *S. aureus* isolates resistant to vancomycin also resistant to Methicillin [35]. This study supported by study conducted in Lusaka, Zambia, A total of 205 paper currency notes were collected from restaurants and hotels and the prevalence of *S. aureus* is 25.85% [36].

According to Table 3, the study found that all 14 (100.0%) of the gram negative bacteria isolates were susceptible to Gentamycin, Ciprofloxacin and Tetracycline which concurs with a study by

Vriesekoop et al. [37] who also found gram negative bacteria like *Klebsiella sp.*, *Enterobacter sp.* and *Proteus sp.* to be sensitive to Amoxoftine, Gentamicin, Nalidixic acid and Ofloxacin. The development of antimicrobial resistance in bacteria renders some infections untreatable today and antimicrobial resistance is now a major health concern [12].

This study revealed that many multidrug resistant strains of different isolates were prevalent in the Kenyan bank note currencies that further emphasize the public health significance of the notes and clearly indicates a marked resistance to the commonly used antibiotics. For example; isolates of various gram negative bacterial species recorded high rates of resistance collectively as 9 (64%), 8 (57%) and 6 (43%) against vancomycin, clindamycin and amoxycillin respectively. This result agree with [38,39,40] presence of multidrug-resistant strains poses a big challenge to human survival and continued existence in relation to bacterial infection and diseases that is highly consequential when contracted by the debilitated individuals. The observed high antibiotic resistances could be attributed to the abuse of antibiotics which showed that majority of the populace sampled purchases antibiotics in the open market without any medical prescription and use them for the wrong diseases and infections [40]. Antibiotics like ciprofloxacin, gentamicin, ceftriaxone and tetracycline; collectively expressed absent and little resistance rates. This latter observation goes with [39,40]. It is therefore suggested that individuals should improve upon their personal health consciousness by washing hands after handling of currency notes [28]. Babies must be prevented from handling currency notes and adults should avoid using saliva during counting of paper.

### 4. CONCLUSION

This study revealed that Kenyan banknote currencies circulating in Nyeri County were contaminated with different pathogenic and potential pathogenic bacteria including multi drug resistant strains. Hence, great care must be taken while handling money during the preparation and handling of food to avoid cross contamination. So, awareness related to the improvement of personal hygiene and good money handling practice such as washing hands properly with soap and water after handling currency before eating and avoiding using saliva during counting money are strongly

recommended as the main pillar to reduce the risk of infection.

## ACKNOWLEDGEMENTS

We are very grateful to the Department of Microbiology, Outspan Teaching and Referral Hospital (OTRH), for allowing us to use their laboratory facility for the study. We are grateful to Nicholas Gituma Bundi for the unlimited help and support during our project, data analysis and manuscript proofreading.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Alwakeel SS, Naseer AL. Bacterial and fungal contamination of Saudi Arabian paper currency and cell phones. *Asian Journal of Biological Sciences*. 2011; 4:556-562.
2. Pradeep NV, Marulasiddaiah BS, Chetana M. Microbial contamination of Indian currency notes in circulation. *Journal of Research in Biology*. 2012;2(4):377-382.
3. Al-Ghamdi A, Abdelmalek S, Bamaga M, Azhar E, Wakid M, Alsaied Z. Bacterial contamination of Saudi "One" riyal paper notes. *Southeast Asian Journal of Tropical Medicine and Public Health*. 2011;42:711-6.
4. Pal K, Das NS, Bhattacharya S. Bacteriological profile of Indian currency circulating in a tertiary care hospital in rural Bengal. *IJRRMS*. 2013;3(2):23-30.
5. Girma G, Ketema T, Bacha K. Microbial load and safety of paper currencies from some food vendors in Jimma Town, Southwest Ethiopia. *BMC Res Notes*. 2014;7:843. DOI:10.1186/1756-0500-7-843.
6. Angelakis E, Esam IA, Bibi F, Yasir M, Al-Ghamdi AK, Ashshi AM, Adel G, Elshemi AG, Raoult D. Paper money and coins as potential vectors of transmissible disease. *Future Microbiol*. 2014;9(2):249-261. DOI:10.2217/fmb.13.161.4
7. Girma G, Ketema T, Bacha K. Microbial load and safety of paper currencies from some food vendors in Jimma Town, Southwest Ethiopia. *BMC Res Notes*. 2014;7:843
8. Alemu A. Microbial contamination of currency notes and coins in circulation: A potential public health hazard. *Biomed Biotechnol*. 2014;2(3):46-53.
9. Saeed S, Rasheed H. Evaluation of bacterial contamination of Pakistani paper currency notes (Rupee) in circulation in Karachi. *European J. Biological Sci*. 2011;3(3):94-98.
10. Ali R, Abbas SZ, Hussain Z, Hussain K, Hayat A, Khan A. Bacteriological analysis and antibiogram of Pakistani paper currency notes in circulation in Karachi, Sindh, Pakistan. *International Journal of Scientific Research in Environmental Sciences*. 2015;3(10):370-376. View at Publisher· View at Google Scholar.
11. Orukotan AA, Yabaya A. Microbial contamination of Naira notes in circulation with in Kaduna Metropolis. *Journal of Medical and Applied Biosciences*. 2011;2: 20-7.
12. Uddin GMN, Larsen MH, Guardabassi L, Dalsgaard A. Bacterial flora and antimicrobial resistance in raw frozen cultured seafood imported to Denmark. *J Food Prot*. 2013;76:490-499.
13. Darehabi HK, Naseri MH, Menbari S, Mobaleghi J, Kalantar E. Antibiotic resistance pattern of *Escherichia coli* groups A, B1, B2 and D isolated from frozen foods and children with diarrhea in Sanandaj, Iran. *Int J Enteric Pathog*. 2013; 1:1-4.
14. Barrow GI, Feltham RKA. Cowan and steel's manual for the identification of medical bacteria. 3. ed. Cambridge: Cambridge University Press. 1993;216. Available:<http://dx.doi.org/10.1017/CBO9780511527104>
15. Clinical and laboratory standards institute. Performance standards for antimicrobial susceptibility testing; twenty-fourth informational supplement. Wayne, USA: CLSI; 2014.
16. Akoachere JFT, Gaele N, Dlonga H, Nkuo- Akenji TK. Public health implications of contamination of Franc CFA (XAF) circulating in Buea (Cameroon) with drug resistant pathogens. *BMC Research Notes*. 2014;7(1):16.
17. Alemu A. Microbial contamination of currency notes and coins in circulation: A potential public health hazard. *Biomedicine and Biotechnology*. 2014;2:46-43.

18. Gedik H, V OSS TA, Voss A. Money and transmission of bacteria. *Antimicrob Resist Infect Control*. 2013;2:22.
19. Sabahat S, Humaira R. Evaluation of bacterial contamination of Pakistani paper currency notes (rupee) in circulation in Karachi. *European Journal of Biological Sciences*. 2011;3(3):94-8.
20. Tagoe DN, Baidoo SE, Dadzie I, Ahator D. A study of Bacterial Contamination of Ghanaian Currency Notes in Circulation. *The Internet J Microbiology*; 2014.
21. Tagoe DN, Adams A, Land VG. Antibiotic resistant bacterial contamination of the Ghanaian currency note: A potential Health problem. *J Microbiol Biotech Res*. 2011;1(4):37-44.
22. Yazah AJ, Yusuf J, Agbo AJ. Bacterial contaminants of Nigerian currency notes and associated risk factors. *Research Journal of Medical Sciences*. 2012;6(1):1-6.
23. Mbajuka CS, Obeagu EI, Nwosu DC, Agbo CE. Microbiological evaluation of naira notes handled by fish sellers in Umuahia metropolis. *World Engineering and Applied Science Journal*. 2014;5(2): 44-52.
24. Agarwal G, Ingle N, Kaur N, Ingle E, Charania Z. Assessment of microbial contamination of Indian currency notes in Mathra city, India: A cross sectional study. *Journal of Advanced Oral Respiration*. 2015;6(2):43-48.
25. Elumalai EK, David E, Hemachandran J. Bacterial contamination of Indian currency notes (rupee). *The International Journal of Occupational and Environmental Medicine*. 2012;3:204-205.
26. Ahmed MSU, Parveen S, Nasreen T, Feroza B. Evaluation of microbial contamination of Bangladesh paper currency notes (Taka) in circulation. *Adv Biol Res*. 2010;4:266-271.
27. Jane-Francis Tatah Kihla Akoachere<sup>1</sup>, 2\*, Nana Gaelle<sup>1</sup>, Henry Meriki Dilonga<sup>1</sup>, Theresa K Nkuo-Akenji<sup>1</sup>. Public health implications of contamination of Franc CFA (XAF) circulating in Buea (Cameroon) with drug resistant pathogens. *BMC Research Notes*. 2014;7:16.
28. Adamu JY, Jairus Y, Ameh JA. Bacteria contaminants of Nigerian currency notes and associated risks factors. *Research Journal of Medical Science*. 2012;6(1):1-6.
29. Mohamed M. Elemam AD, Mahmoud BS, Khaled D. A study of bacterial contamination on Libyan paper banknotes in circulation. *American Journal of Microbiology and Biotechnology*. 2016; 3(1):1-6.
30. Yakubu J, Ehiowemwenguan G, Inetianbor JE. Microorganisms associated with mutilated Naira notes in Benin-city, Nigeria. *International Journal of Basic and Applied Science*. 2014;3:9-15.
31. Pavani G, Srividya Y. Bacterial analysis of currency in circulation in a rural teaching hospital in India. *Journal Microbiol. Biotech. Res*. 2014;4:34-38.
32. Azza SM, Abuelnaga AA, Samy MA, Bakry AS. Bacteriological assay for the Egyptian currency collected from veterinary field. *International Journal of Microbiological Research*. 2014;5(1):48-53.
33. Felgo P, Nkansah M. Bacterial load on Ghanaian currency notes. *Afr J Microbiology Research*. 2010;4(22):2375-80.
34. Srinu B, Vijaya Kumar A, Kumar E, Madhava Rao T. Antimicrobial resistance of bacterial foodborne pathogens. *J. Chemical and Pharmaceutical Res*. 2012; 4(7):3734–3736.
35. Neel R. Multidrug resistance of isolates of methicillin resistant *Staphylococcus aureus* (MRSA) in paper currency notes from meat sellers in Tanga, Tanzania. *International Journal LifeSc Bt & Pharm Res*. 2012;1(4):9-13.
36. Neel R. Multidrug resistance of isolates of methicillin resistant *Staphylococcus aureus* (MRSA) in paper currency notes from restaurants and hotels in Lusaka in Zambia. *International Journal of Pharmacy and Pharmaceutical Sciences*. 2013;5(1): 363-6.
37. Vriesekoop F, Russell C, Alvarez MB. Dirty money. An investigation into Hygiene status of some of the world's currencies as obtained from food outlets. *Foodborne Pathogens and Disease*. 2010;1497-1502.
38. Ogunleye AG, Omoya FO, Ayantola KJ. Bacterial antibiogram and physicochemical parameters of well water in Iworoko-Ekiti, Nigeria. *J. Appl. Life Sci. Int*. 2016;4(4):1-10.
39. Samie A, Makonto TE, Odiyo J, Ouaboi-Egbenni PO, Mojapelo P, Bessong PO. Microbial quality, diversity and antibiotic susceptibility profiles of bacterial isolates from borehole water used by schools in Greater Giyani Municipality,

- Mopani District, South Africa. Afr. J. Microbiol. Res. 2011;5(3):198-210.
40. Ayandiran TA, Ayandele AA, Dahunsi SO, Ajala OO. Microbial assessment and prevalence of antibiotic resistance in polluted Oluwa River, Nigeria. Egypt. J. Aquat. Res. 2014;40:291–299.
41. Struve C, Bojer M, Krogfelt KA. Characterization of klebsiellapneumoniaetype 1 fimbriae by detection of phase variation during colonization and infection and impact on virulence. Infect Immun. 2008;76(9):4055–4065.

© 2019 Muguongo et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<http://www.sdiarticle3.com/review-history/49596>