

## Research Article

# Phytochemical Screening and *In Vitro* Evaluation of the Antioxidant Potential of Dichloromethane Extracts of *Strychnos henningsii* Gilg. and *Ficus sycomorus* L.

Kenedy Wanjala Wafula <sup>1</sup>, Joseph Kiambi Mworira <sup>2</sup>, and Mathew Piero Ngugi <sup>1</sup>

<sup>1</sup>Department of Biochemistry, Microbiology and Biotechnology, Kenyatta University, P.O. Box 43844-00100, Nairobi, Kenya

<sup>2</sup>Department of Physical Sciences, Meru University of Science and Technology, P.O. Box 972 60200, Meru, Kenya

Correspondence should be addressed to Kenedy Wanjala Wafula; kenwaf@gmail.com

Received 7 December 2022; Revised 22 February 2023; Accepted 10 March 2023; Published 29 March 2023

Academic Editor: Daiana S. Avila

Copyright © 2023 Kenedy Wanjala Wafula et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Medicinal plants are a rich source of antioxidants such as flavonoids, phenols, tannins, and alkaloids among others and are currently used as alternative and complementary drugs in the management of stress-related disorders. *Strychnos henningsii* and *Ficus sycomorus* have been traditionally used by the people of Mbeere, Embu county, Kenya, as medicine for the treatment of various oxidative stress-related disorders such as diabetes and rheumatism; however, no empirical data are available to authenticate the said claim. The aim of this study was to evaluate preliminary phytochemical screening and *in vitro* antioxidant activity of dichloromethane (DCM) leaf extract of *S. henningsii* and stem bark extract of *F. sycomorus* using DPPH, hydrogen peroxide, and ferric reducing power assays; total flavonoids and phenolic compounds were also determined by colorimetric assay and Folin-Ciocalteu reaction, respectively. Phytochemical screening showed that both extracts possessed saponins, flavonoids, phenols, steroids, alkaloids, and cardiac glycosides; however, terpenoids were found to be absent in *S. henningsii*. The total phenolic and flavonoid content of the DCM stem bark extract of *F. sycomorus* was lower than that of the leaf extract of *S. henningsii*. These extracts significantly exhibited strong antioxidant activities at different concentrations tested. The IC<sub>50</sub> values of *S. henningsii* and *F. sycomorus* were 0.325 mg/ml and 0.330 mg/ml for hydrogen peroxide and 0.068 mg/ml and 0.062 mg/ml for DPPH, respectively. Both DCM leaf and stem bark extracts of *S. henningsii* and *F. sycomorus* were found to have strong ferric reducing power. Therefore, both extracts showed significant nonenzyme-based antioxidant activities. The two plants possess phytochemicals that have significant antioxidant properties.

## 1. Introduction

Oxidative stress is the disparity between the production of free radicals and antioxidant defenses in the body [1–5]. Free radicals are defined as compounds with unpaired electrons, making them highly reactive molecules that can attack any stable molecules such as proteins, carbohydrates, and lipids [6, 7]. Reactive oxygen species (ROS) are the most common and widely known free radicals. They include superoxide (O<sub>2</sub><sup>-</sup>), hydroxyl (HO<sup>-</sup>), hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>), and nitric oxide (NO<sup>-</sup>). Most biochemical reactions in the body are known to generate ROS [8], which are potent in damaging important biomolecules such as proteins, nucleic

acids, and lipids if they are not scavenged by antioxidants [9]. Free radicals are well known to be involved in aging and pathogenesis of stress-related disorders such as diabetes, nephrotoxicity, hepatotoxicity, malignancy, cardiovascular disorders, inflammation, and neurological disorders [10–12].

Human cells are well protected by antioxidant defense systems against ROS attack; however, at low concentrations of antioxidant enzymes, some cells have been shown to be sensitive to ROS [13–16]. The cellular antioxidant level is used to determine the susceptibility of tissues to oxidative damage. This level normally changes during oxidative stress [17–19]. A wide variety of antioxidants are naturally

obtained from plants that constitute our daily diet. Commonly known dietary antioxidants are vitamins C and E, carotenoids, and green tea [20, 21]. The consumption of food and fruits rich in antioxidants plays a significant role in augmenting the body's natural resistance to oxidative stress [22, 23]. Plants also have many other nonnutrient antioxidants such as phenols, flavonoids, and alkaloids. These polyphenol compounds have been extensively studied and documented as quenchers of free radicals [24, 25].

Since antioxidants hold a key in preventing oxidative stress-related disorders, many plant extracts and their secondary metabolites are being explored for their antioxidant effects [26, 27]. The use of plant-based antioxidants plays an important role in preventing the activation of the oxidation-induced signaling pathways in our bodies [28]. Therefore, the identification of the antioxidant activities of the DCM leaf extract of *S. henningsii* and stem bark of *F. sycomorus* is an important step in increasing our understanding about their usage in the treatment of various stress-related disorders.

Commercially available antioxidant drugs include butyrate hydroxyanisole, butylated hydroxytoluene, propyl gallate, and fluconazole [29–31]. However, studies have shown that these synthetic antioxidants have toxic effects and show negative health influence [32, 33] and have led to some restrictions being imposed on their use [34]. Researchers now have focused their attention on plant-derived antioxidants among others [35].

Globally, several plants have been traditionally used for their antioxidant activities [36, 37]. Ethno-pharmacological surveys indicate that medicinal plants play a vital role in the management of oxidative stress-related disorders [38, 39]. Plant extracts naturally possess phytochemicals such as flavonoids, tannins, phenols, and alkaloids [40], which are well-known antioxidants and are currently pursued as alternative and complementary remedies against oxidative stress-related disorders [41]. Several efficacy studies conducted on herbal plants have shown that plant-based antioxidants are relatively safe, cost efficient, and effective in disease management [42].

The genus *Ficus* is widely known to have strong antioxidant properties due to its richness in phenols and flavonoids [43–45]. Traditionally, *Ficus sycomorus* fruits, stem barks, and roots have been used as herbal remedies for several ailments such as diarrhea, liver disease, skin infections, stomach disorders, helminthiasis, lactation disorders, epilepsy, tuberculosis, sterility, and diabetes mellitus [46–48]. *S. henningsii* is a widely distributed evergreen herb in East Africa [47] and is used in the management of rheumatism, snake bite, abdominal pain, gastrointestinal pain, gynecological complaints, malaria, and diabetes mellitus [49]. The crude extracts of *Strychnos henningsii* have been documented to possess significant therapeutic effects against stress-related disorders [50]. Based on traditional pharmacology, it has been actively and successfully employed by the Mbeere community in Embu county, Kenya, in the management of diabetes, which is an oxidative stress-related disorder.

In view of this background, the present study seeks to investigate the *in vitro* antioxidant activities of dichloromethanolic (DCM) leaf extract of *S. henningsii* and stem bark extract of *F. sycomorus*. The study aims to explore and provide preliminary information on the *in vitro* antioxidant activities of *S. henningsii* and *F. sycomorus* as possible bioresources for the generation of herbal formulations used in the treatment and management of oxidative stress-related disorders. The study also aims to reveal relevant research gaps that need to be explored further.

## 2. Materials and Methods

**2.1. Collection of Plant Materials.** The authors sought authorization from The National Commission for Science, Technology and Innovation (NACOSTI/P89/6765/9816). Plant materials of *S. henningsii* leaves and stem barks of *F. sycomorus* were collected from their natural habitats from Makunguru village, Mbeere north subcounty, Embu county, Kenya, in October, 2015. The GPS locations for *S. henningsii* and *F. sycomorus* specimens were 0°34'11"S, 37°37'31"E and 0°35'28"S, 36°36'22"E, respectively. The collection of these samples was done based on ethnobotanical information availed by local herbalists in the area. The plant identification was done by an acknowledged authority from Kenyatta University authenticating their botanical identities. *S. henningsii* and *F. sycomorus* were assigned voucher specimen numbers (*S. henningsii*. (Wkw001/10/21015) and *F. sycomorus* (Wkw 002/10/2015), respectively). They were deposited in the Kenyatta University Herbarium for future reference. Sample materials were carefully sorted, packed in sealed bags, and transported to the Department of Biochemistry and Biotechnology, Kenyatta University, where further processing and subsequent study was undertaken.

**2.2. Extract Preparations.** The fresh plant materials were air dried at room temperature under a shade for a week. The dried leaves were milled into fine powder by use of an electric mill. The powdered plant materials were sieved using a mesh pore of 0.5 mm and packed in closed, dry sealed bags and stored awaiting extraction. Two hundred and fifty grams (250 g) of each powdered plant material was soaked in 1 litre of dichloromethane (DCM) and macerated for 24 hours. The resultant extract was poured into a clean dry conical flask and then filtered using Whatman's No. 1 filter papers. The filtrate was extracted using Soxhlet apparatus for 5–6 h and then concentrated under reduced pressure and vacuum using a rotary evaporator at a temperature of 40°C. The concentrates were placed in airtight containers weighed and stored at –4°C awaiting use in bioassays.

**2.3. Qualitative Phytochemical Screening.** Qualitative phytochemical screening of DCM extracts of *S. henningsii* and *F. sycomorus* was performed to determine the presence or absence of selected plant secondary metabolites using standard methods described by Harbone [51] and Kotake [52]. Secondary metabolites screened include flavonoids, cardiac glycosides, saponins, alkaloids, sterols, phenolics,

and terpenoids. These phytochemicals are associated with antioxidant activities.

#### 2.4. Quantitative Phytochemical Determination

**2.4.1. Determination of Total Phenolic Contents.** Folin–Ciocalteu reagent was used to determine the total phenolic levels of the plant extracts as described by Spanos et al. [53] and modified by Lister and Wilson [54]. Briefly, 2 ml of each plant extract, 2.5 ml of 10% dilution of Folin–Ciocalteu reagent and 2 ml of Na<sub>2</sub>CO<sub>3</sub> (7.5%, w/v) were mixed and left to stand for 15 minutes at 45°C. The absorbance of all the treatments was determined at 765 nm spectrophotometrically. Gallic acid was used as the reference to derive the calibration curve. The total phenolic content was determined using the linear equation based on the calibration curve and contents expressed as milligrams of gallic equivalent per gram of dry weight (mg GAE/g dw) [55].

**2.4.2. Determination of Total Flavonoid Contents.** The colorimetric methodology described by Lamaison and Carnet [56] and Nurcholis et al. [57] was used to determine the total flavonoid contents of the extracts. Briefly, a volume of 1.5 ml of the extracts was mixed with an equivalent volume of 2%

AlCl<sub>3</sub>.6H<sub>2</sub>O (2 g in 100 ml methanol) solution. The solution was vigorously shaken to mix and then incubated for 10 minutes, and the absorbance was read at 430 nm using a spectrophotometer. Rutin was used as the reference to generate the calibration curve. The total flavonoid content was expressed as milligrams of rutin equivalent per gram of dry weight (mg RE/g dw) based on the calibration curve [55].

#### 2.5. Determination of In Vitro Antioxidant Activities

**2.5.1. Determination of In Vitro Hydrogen Peroxide Scavenging Activity.** The *in vitro* hydrogen peroxide scavenging potential of DCM extracts of *S. henningsii* and *F. sycomorus* was determined following the protocol described by Ruch et al. [58]. Briefly, 50 Mm, pH 7.4, phosphate buffer solution was used to prepare 250 ml solution of hydrogen peroxide (40 mM). Hydrogen peroxide solution at a volume of 0.6 ml was added to 1 ml of varying concentrations (0.1–0.5 mg/ml) of the plant extract and ascorbic acid (standard). The mixture was left to stand for 10 minutes, after which the absorbance was determined at 560 nm using a UV spectrophotometer. The blank solution containing phosphate buffer was used as the negative control. This was done in three replicates. The percentage radical scavenging activity is as follows:

$$\% \text{ Hydrogen scavenging activity} = \frac{\text{Abs. of control} - \text{Abs. of sample/standard}}{\text{Abs. of control}} \times 100, \quad (1)$$

where Abs. is the absorbance.

**2.5.2. Determination of In Vitro Diphenyl-2-picrylhydrazyl (DPPH) Radical Scavenging Activity.** The abilities of the DCM extracts of *S. henningsii* and *F. sycomorus* to scavenge DPPH radicals *in vitro* were determined based on the method documented by Mehrotra et al. [59–61]. Following this method, 2.66 mg of DPPH was dissolved in 50 ml of ethanol to form a concentration of 0.135 mM. Various dilutions, namely, 0.2, 0.1, 0.05, 0.025, and 0.0125 mg/ml, of the plant extracts and ascorbic acid (standard) were prepared. One milliliter of the DPPH solution dissolved in

methanol was mixed with 1 ml of each diluted plant extract and ascorbic acid (reference drug). The mixtures were then agitated thoroughly and left in a dark room for 30 minutes at room temperature. Three replicates of the assays were prepared. The absorbance of the mixture was then measured at 517 nm using a spectrophotometer. The actual decrease in absorbance was measured against that of the control. The negative control was a blank solution containing ethanol without H<sub>2</sub>O<sub>2</sub>. The percentage DPPH scavenging abilities of the plant extracts were then derived using the following equation:

$$\% \text{ DPPH scavenging activities} = \frac{\text{Abs. of control} - \text{Abs. of sample/standard}}{\text{Abs. of control}} \times 100, \quad (2)$$

where Abs. is the absorbance.

**2.5.3. Calculation of Half Maximal Inhibitory Concentrations (IC<sub>50</sub>) in Hydrogen Peroxide and DPPH Radicals.** The half maximal inhibitory concentration (IC<sub>50</sub>) of DCM extracts of *S. henningsii*, *F. sycomorus*, and ascorbic acid (standard) were analyzed using linear regression analysis in MS Excel.

The IC<sub>50</sub>, which represents the concentration at which 50% of the radicals were scavenged by test samples, was determined from a graph of percentage scavenging activity against the concentration of the test sample.

**2.5.4. In Vitro Ferric Reducing Power Assay.** The *in vitro* ferric reducing power of the DCM extracts of *S. henningsii*

and *F. sycomorus* alongside ascorbic acid (positive control) was established according to the protocol described by Oyaizu [62]. In brief, various concentrations (0.2–1 mg/ml) of 1 ml of the plant extracts and ascorbic acid were added to 2.5 ml of 0.2 M phosphate buffer of pH 7. The resulting solution was then mixed with 2.5 ml of potassium ferricyanide and incubated at 50°C for 20 minutes. Afterwards, 2.5 ml of trichloroacetic acid (10%) was then added to the mixture and centrifuged for 10 minutes at 3000 rpm. Then, 2.5 ml was drawn from the upper layer of the solution and then added to 2.5 ml of distilled water and 0.5 ml freshly prepared ferric chloride ( $\text{FeCl}_3$ ) solution (1%) was added. The assay was done in triplicates. The absorbance of the extracts and ascorbic acid was determined at 700 nm using a spectrophotometer.

**2.6. Statistical Analysis.** The data were subjected to descriptive statistics using Minitab Statistical Software 17.0 (State College, Pennsylvania) and expressed as mean  $\pm$  standard error of mean (SEM). One-way analysis of variance (ANOVA) was performed to determine the statistically significant difference among treatments. Tukey's tests were performed for pairwise comparison of means. Unpaired student's *t*-test was used for the comparison of mean total phenolic and flavonoid contents of DCM leaf extract of *S. henningsii* and stem bark of *F. sycomorus*. The values of  $p \leq 0.05$  were considered to be significantly different. The data obtained were presented in a tabular and graphical form. The phytochemical screening was done qualitatively, and the result obtained (positive/negative) for each test was recorded in a table.

### 3. Results

**3.1. Qualitative Phytochemical Screening.** The phytochemistry of the leaf extract of *S. henningsii* revealed the presence of alkaloids, phenols, saponins, cardiac glycosides, flavonoids, and steroids while terpenoids were absent. On the other hand, *F. sycomorus* contained saponins, flavonoids, alkaloids, steroids, phenols, cardiac glycosides, and terpenoids (Table 1).

#### 3.2. Quantitative Phytochemicals Screening

**3.2.1. Total Phenolic Contents.** The total phenolic concentration in the leaf extract of *S. henningsii* and stem bark extract of *F. sycomorus* were quantified and expressed as milligrams of gallic acid equivalent per gram of dry weight (mg GAE/g dw) (Table 2) using a standard gallic acid calibration curve ( $y = 1.52x + 0.234$ ;  $R^2 = 0.9782$ ). Generally, at all the concentrations tested (0.2, 0.4, 0.6, 0.8, and 1.0 mg/ml), the two extracts had significantly different phenolic contents ( $p \leq 0.05$ ), with *S. henningsii* extract having the highest phenolic content (Table 2).

**3.2.2. Total Flavonoid Contents.** The total flavonoid contents of the DCM leaf and stem bark extracts of *S. henningsii* and *F. sycomorus* were calculated from the standard rutin

TABLE 1: Phytochemical screening of DCM leaf extracts of *S. henningsii* and stem bark of *F. sycomorus*.

Composition	<i>F. sycomorus</i>	<i>S. henningsii</i>
Saponins	+	+
Alkaloids	+	+
Terpenoids	+	-
Flavonoids	+	+
Steroids	+	+
Phenolics	+	+
Cardiac glycosides	+	+

+, present; -, absence.

calibration curve ( $y = 2.535x - 0.047$ ;  $R^2 = 0.9778$ ). The leaf extract of *S. henningsii* had significantly higher flavonoid concentrations than the *F. sycomorus* stem bark extract ( $p \leq 0.05$ ; Table 2). However, the rutin equivalence at the lowest concentration of 0.1 mg/ml for both extracts was not significantly different ( $p > 0.05$ ; Table 3).

**3.3. In Vitro Hydrogen Peroxide Radical Scavenging Activities of DCM Extracts of *S. henningsii* and *F. sycomorus*.** The *in vitro* hydrogen peroxide scavenging potential of the DCM leaf extract of *S. henningsii* and stem bark extract of *F. sycomorus* was analyzed, as shown in Figure 1. Generally, both extracts of *S. henningsii* and *F. sycomorus* showed remarkable *in vitro* hydrogen peroxide scavenging activity at all the concentrations (0.1, 0.2, 0.3, 0.4, and 0.5 mg/ml). The extracts showed  $\text{H}_2\text{O}_2$  scavenging activities in a dose-related manner. The  $\text{H}_2\text{O}_2$  scavenging activities of both extracts were significantly lower than that of the standard (ascorbic acid). However, there was no significant difference in the hydrogen peroxide scavenging activities of *S. henningsii* and *F. sycomorus* at all the tested concentrations ( $p > 0.05$ ; Figure 1 and Table 4).

**3.4. In Vitro DPPH Radical Scavenging Activities of DCM Extracts of *S. henningsii* and *F. sycomorus*.** As shown in Figure 2 and Table 5, both extracts of *S. henningsii* and *F. sycomorus* as well as the ascorbic acid demonstrated dose-dependent scavenging of DPPH radicals. At all the tested doses (0.00125–0.2 mg/ml), the two extracts showed lower DPPH scavenging activities than the standard (ascorbic acid). The extract activities were found not to be significantly different at all the tested concentrations.

**3.5.  $\text{IC}_{50}$  for Hydrogen Peroxide and DPPH.** The half maximal percentage inhibition ( $\text{IC}_{50}$ ) value is widely used as a quantitative measure of extracts, antioxidant potential. The two extracts showed a lower  $\text{IC}_{50}$ , as shown in Table 6.

**3.6. In Vitro Ferric Reducing Power Activities of DCM Extracts of *S. henningsii* and *F. sycomorus*.** In this assay, the extracts were tested for their ability to reduce  $\text{Fe}^{3+}$  to  $\text{Fe}^{2+}$  via electron donation. The results showed a dose-related ferric reduction by the DCM extracts of *S. henningsii* and *F. sycomorus*. The leaf extract of *S. henningsii* was

TABLE 2: Total phenolic contents of DCM extracts of *S. henningsii* and *F. sycomorus*.

Groups	Mass in mg/g gallic acid equivalent				
	0.1 mg/g	0.2 mg/g	0.3 mg/g	0.4 mg/g	0.5 mg/g
<i>S. henningsii</i>	4.3 ± 0.01 <sup>a</sup>	10.7 ± 0.01 <sup>a</sup>	16.7 ± 0.01 <sup>a</sup>	22.0 ± 0.01 <sup>a</sup>	29.0 ± 0.01 <sup>a</sup>
<i>F. sycomorus</i>	2.3 ± 0.01 <sup>b</sup>	5.0 ± 0.01 <sup>b</sup>	9.7 ± 0.01 <sup>b</sup>	16.0 ± 0.01 <sup>b</sup>	23.0 ± 0.015 <sup>b</sup>

Values are expressed as mean ± SEM of the three replicates. Column means followed by the same superscript letters are not significantly different ( $p > 0.05$ ) by the unpaired Student's *t*-test.

TABLE 3: Total flavonoid content of DCM extracts of *S. henningsii* and *F. sycomorus*.

Groups	Mass in mg/g rutin equivalent				
	0.2 mg/g	0.4 mg/g	0.6 mg/g	0.8 mg/g	1.0 mg/g
<i>S. henningsii</i>	15.7 ± 0.00 <sup>a</sup>	22.7 ± 0.00 <sup>a</sup>	37.7 ± 0.00 <sup>a</sup>	44.3 ± 0.00 <sup>a</sup>	53.0 ± 0.00 <sup>a</sup>
<i>F. sycomorus</i>	10.3 ± 0.01 <sup>a</sup>	16.0 ± 0.01 <sup>b</sup>	23.7 ± 0.01 <sup>b</sup>	34.3 ± 0.00 <sup>b</sup>	38.3 ± 0.00 <sup>b</sup>

Values are expressed as mean ± SEM of the three replicates. Column means followed by the same superscript letters are not significantly different ( $p > 0.05$ ) by the unpaired Student's *t*-test.




FIGURE 1: Percentage hydrogen peroxide inhibition of the DCM leaf extract of *S. henningsii*, the stem bark extract of *F. sycomorus*, and the standard ascorbic acid. Values are expressed as mean ± SEM of the three replicates. Column means followed by the same superscript letters were not significantly different ( $p > 0.05$ ) by one-way ANOVA followed by Tukey's post hoc test.

TABLE 4: % in vitro hydrogen peroxide radical scavenging activities of DCM extract of *S. henningsii* and *F. sycomorus*.

Groups	Concentrations					IC <sub>50</sub> value
	0.1 mg/ml	0.2 mg/ml	0.3 mg/ml	0.4 mg/ml	0.5 mg/ml	
Ascorbic acid %	27.72 ± 2.38 <sup>a</sup>	42.25 ± 2.27 <sup>a</sup>	60.38 ± 1.77 <sup>a</sup>	74.24 ± 2.68 <sup>a</sup>	88.05 ± 2.55 <sup>a</sup>	0.245 mg/ml
<i>F. sycomorus</i> %	15.71 ± 1.47 <sup>b</sup>	31.07 ± 4.15 <sup>b</sup>	47.24 ± 2.38 <sup>b</sup>	61.41 ± 2.81 <sup>b</sup>	75.85 ± 1.66 <sup>b</sup>	0.330 mg/ml
<i>S. henningsii</i> %	14.22 ± 2.69 <sup>b</sup>	30.45 ± 2.64 <sup>b</sup>	48.27 ± 2.50 <sup>b</sup>	61.57 ± 2.92 <sup>b</sup>	75.22 ± 2.12 <sup>b</sup>	0.325 mg/ml

Values are expressed as mean ± SEM for three replicates. Column means with the same superscript were not significantly different by ANOVA followed by Tukey's post hoc test ( $p > 0.05$ ).

comparable to that of standard ascorbic acid at the lowest concentration of 0.2 mg/ml. However, at this concentration, the stem bark extract of *F. sycomorus* had significantly lower ferric reducing power than ascorbic acid. At all the other tested concentrations (0.4–1 mg/ml), the ferric reducing power activities of the two extracts had no significant difference ( $p > 0.05$ ). However, they were significantly different from that of the standard (ascorbic acid) ( $p \leq 0.05$ ; Figure 3 and Table 7).

## 4. Discussion

4.1. *Qualitative and Quantitative Analysis.* The search for natural antioxidants has grown rapidly amongst clinical and medical practitioners due to the interest generated by reactive oxygen species (ROS) and pathogenesis of oxidative stress-related disorders [63, 64]. Plant leaves, stems, flowers, fruits, and roots have been known for centuries to possess therapeutic value and, therefore, have been extensively




FIGURE 2: Percentage DPPH scavenging activities of *S. henningsii*, *F. sycomotorus*, and standard ascorbic acid. Values are expressed as mean  $\pm$  SEM of the three replicates. Column means followed by the same superscript letters were not significantly different ( $p > 0.05$ ) by one-way ANOVA followed by Tukey's post hoc test.

TABLE 5: % in vitro DPPH radical scavenging activities of DCM extracts of *S. henningsii* and *F. sycomotorus*.

Groups	Concentrations					IC <sub>50</sub> value
	0.0125 mg/ml	0.025 mg/ml	0.05 mg/ml	0.1 mg/ml	0.2 mg/ml	
Ascorbic acid %	42.84 $\pm$ 1.05 <sup>a</sup>	50.00 $\pm$ 1.85 <sup>a</sup>	64.06 $\pm$ 1.75 <sup>a</sup>	80.13 $\pm$ 2.06 <sup>a</sup>	91.39 $\pm$ 1.38 <sup>a</sup>	0.025 mg/ml
<i>S. henningsii</i> %	30.38 $\pm$ 1.61 <sup>b</sup>	34.93 $\pm$ 2.42 <sup>b</sup>	46.60 $\pm$ 1.49 <sup>b</sup>	61.71 $\pm$ 1.69 <sup>b</sup>	78.33 $\pm$ 1.65 <sup>b</sup>	0.068 mg/ml
<i>F. sycomotorus</i> %	29.73 $\pm$ 2.21 <sup>b</sup>	37.94 $\pm$ 1.15 <sup>b</sup>	46.45 $\pm$ 2.74 <sup>b</sup>	63.81 $\pm$ 2.27 <sup>b</sup>	76.38 $\pm$ 1.92 <sup>b</sup>	0.062 mg/ml

Values are expressed as mean  $\pm$  SEM of the three replicates. Column means followed by the same superscript letters were not significantly different ( $p > 0.05$ ) by one-way ANOVA followed by Tukey's post hoc test.

TABLE 6: IC<sub>50</sub> values of DCM extracts of *S. henningsii* and *F. sycomotorus*.

	Hydrogen peroxide	DPPH
<i>S. henningsii</i>	0.330 mg/ml	0.068 mg/ml
<i>F. sycomotorus</i>	0.325 mg/ml	0.062 mg/ml
Ascorbic acid	0.245 mg/ml	0.025 mg/ml

studied to provide alternative answers to the diverse stress-related disorders. Arguably, plants tend to offer better alternatives for varied sources of medicinal remedy including antioxidants.

Based on phytochemical studies, the yield and the antioxidant activity of a plant extract depend on the selected extraction solvent [65–67]. Different solvents are employed to isolate different antioxidant compounds based on their disparities in polarities [68]. In this study, dichloromethane solvents were used as a midpolar solvent for extraction of polyphenols and other midpolar phytochemicals. Several assays (DPPH radical scavenging assay, hydroxyl radical assay, hydrogen peroxide radical scavenging assay, ferric reducing power, and total determination of flavonoid and phenolic contents, among others) have been developed for the determination of in vitro nonenzymatic antioxidant activities of medicinal plants [69, 70].




FIGURE 3: Analysis of ferric reducing power of *S. henningsii*, *F. sycomotorus*, and the standard ascorbic acid. Values are expressed as mean  $\pm$  SEM of the three replicates. Column means followed by the same superscript letters were not significantly different ( $p > 0.05$ ) by one-way ANOVA followed by Tukey's post hoc test.

Flavonoids and phenols naturally exhibit strong scavenging abilities for free radicals due to their hydroxyl groups [71, 72], which are attached to their aromatic ring structures and help to quench the radicals either by donating their electrons thus neutralizing them or via the electron

TABLE 7: In vitro ferric reducing power activities of the DCM extracts of *S. henningsii* and *F. sycomorus*.

Groups	Concentrations				
	0.2 mg/ml	0.4 mg/ml	0.6 mg/ml	0.8 mg/ml	1.0 mg/ml
Ascorbic acid (nm)	0.427 ± 0.02 <sup>a</sup>	0.612 ± 0.02 <sup>a</sup>	0.705 ± 0.04 <sup>a</sup>	0.864 ± 0.02 <sup>a</sup>	0.915 ± 0.02 <sup>a</sup>
<i>S. henningsii</i> (nm)	0.317 ± 0.04 <sup>ab</sup>	0.452 ± 0.02 <sup>b</sup>	0.524 ± 0.02 <sup>b</sup>	0.643 ± 0.02 <sup>b</sup>	0.730 ± 0.02 <sup>b</sup>
<i>F. sycomorus</i> (nm)	0.217 ± 0.04 <sup>b</sup>	0.316 ± 0.04 <sup>b</sup>	0.486 ± 0.02 <sup>b</sup>	0.603 ± 0.02 <sup>b</sup>	0.703 ± 0.02 <sup>b</sup>

Values are expressed as mean ± SEM of the three replicates. Column means followed by the same superscript letters were not significantly different ( $p > 0.05$ ) by one-way ANOVA followed by Tukey's post hoc test.

delocalization over all three ring systems achieved by ortho-dihydroxy of the B-ring and 4-oxo group of the ring C of the flavonoid, which actively reduce radicals such as DPPH and  $\text{Fe}^{3+}$  to  $\text{Fe}^{2+}$  ions [73, 74]. Thus, polyphenols directly augment the antioxidant potential through the restoration of redox balance [75].

In most plant extracts, there is a positive correlation between antioxidant activity and the amount of polyphenolic compounds [76, 77]. However, some studies have reported that there is no positive relationship between the polyphenolic compounds and their antioxidant activities [78, 79]. The total phenolic and flavonoid content of the DCM stem bark extract of *F. sycomorus* was lower than that of the leaf extract of *S. henningsii* at all the tested concentrations. This partly explains why *S. henningsii* had a better result against % DPPH radical scavenging activities and ferric reducing power assay (Figures 1 and 2) than *F. sycomorus*. Nevertheless, the difference in polyphenolic content between the two extracts was not statistically significant against hydrogen peroxide scavenging abilities (Figure 3). In this study, the antioxidant activity observed against hydrogen peroxide, DPPH radicals, and FARP were due to the presence of phenolic and flavonoid components. Therefore, a positive correlation was noted between the amount of polyphenolics and the antioxidant activity.

**4.2. Statistical Analysis.** Hydrogen peroxide ( $\text{H}_2\text{O}_2$ ) is a harmless and less reactive molecule, which becomes harmful, toxic, and reactive to the cell when it is converted to hydroxyl radical [80, 81]. Hydroxyl compounds are among the most deleterious ROS produced by mitochondria, causing oxidative damage, and are clinically linked to causes of various stress-related disorders [82]. Thus, the removal of  $\text{H}_2\text{O}_2$  is a critical step for maintaining a functional antioxidant defense system in cells or food systems [83, 84].

DCM leaf extract of *S. henningsii* and stem bark extract of *F. sycomorus* showed significant antioxidant activity against hydrogen peroxide in a dose-associated trend. Sirisha et al. [85] also demonstrated similar in vitro antioxidant activity while working on the methanolic leaf extract of *F. carica*. The dose-dependent activities of both extracts showed that an increase in the concentration of the extract increased the levels of bioactive antioxidant compounds [86]. Similarly, studies conducted on different *Ficus* species by Ahoua et al. reported the species exhibit strong hydrogen peroxide antioxidant activity in a dose-dependent manner [87]. The strong scavenging potential of the extracts is

reflected in their low  $\text{IC}_{50}$  value. A lower  $\text{IC}_{50}$  is normally associated with a higher radical scavenging activity [88].

The findings of this study showed that both *S. henningsii* and *F. sycomorus* extracts had low  $\text{IC}_{50}$  of 0.330 mg/ml and 0.325 mg/ml, respectively, against the  $\text{H}_2\text{O}_2$  radicals. Thus, the low  $\text{IC}_{50}$  values obtained from the study indicate that the two extracts have strong  $\text{H}_2\text{O}_2$  scavenging activities. The  $\text{IC}_{50}$  value obtained for *F. sycomorus* was similar to that obtained by Deo et al. [89] working on some selected herbal extract inhibitory properties against protein glycation and angiotensin enzyme linked to type II diabetes. Additionally, *P. amarus* and *L. pumila* var. *alata* medicinal plants have been shown to possess potent radical inhibiting properties with low  $\text{IC}_{50}$  values of 3.4 and 5.7  $\mu\text{g}/\text{ml}$ , respectively [90].

The potential of plant extracts to inhibit the DPPH radical is strongly linked to their ability to donate electrons to the radical [46, 91]. Normally, DPPH radical is stable in various solvents including methanol, ethanol, and water. Therefore, the radical is usually prepared in a solution of either ethanol or methanol [70, 92]. The results obtained in this study showed dose-dependent DPPH scavenging activities of the two extracts. It was, however, noted that the DCM leaf extract of *S. henningsii* and *F. sycomorus* stem bark extract had lower DPPH scavenging abilities than ascorbic acid [93]. This could be due to the crude nature of extract as compared to the refined standard drug. This result corresponded to the observations of Igbiosa et al. [94] who found that *Jatropha curcas* had lower DPPH activities than ascorbic acid (standard).

The results of the stem bark extract of *F. sycomorus* agree with a study by Kampli et al. [95], who found that the DPPH scavenging activity of *F. racemosa* was considerable but not higher than that of the standard drug. The good antioxidant property of *F. sycomorus* stem bark extract against DPPH corroborates well with the findings of Santiago and Mayor [96], who noted that *F. odorata* had a good antioxidant activity against DPPH radicals. The DCM leaf extract of *S. henningsii* also showed a good antioxidant activity against DPPH. However, studies conducted by Oyedemi et al. [48], while working on the stem bark extract of *S. henningsii*, reported a weaker antioxidant activity against DPPH radicals and attributed it to low levels of flavonoids in the stem bark extract compared to that of the leaf. The potential antioxidant activities of the DCM extract of *S. henningsii* and *F. sycomorus* against DPPH radicals in this study can be positively related to their higher total phenolic ( $R^2 = 0.9782$ ) and flavonoid ( $R^2 = 0.9778$ ) compounds (Tables 2 and 3). This higher antioxidant activity of the two extracts was a reflection of the lower  $\text{IC}_{50}$  values obtained (0.062 mg/ml

and 0.068 mg/ml, Table 1). In DPPH assay, it has been strongly suggested that samples with lower IC<sub>50</sub> values of 50 µg/ml are very strong antioxidants, while with a range of 50–100 µg/ml are strong antioxidants, and with values above 150 µg/ml are weaker antioxidants [97].

The potential of the two extract samples to reduce Fe<sup>3+</sup> to Fe<sup>2+</sup> via electron donation was determined by the amount of Fe<sup>2+</sup> complex generated and measured in terms of the absorbance of Perl's Prussian blue colour at 700 nm wavelength [98]. The change of yellow colour of the test solution to various shades of green and blue indicated the reducing power of the extract [99]. The phytochemicals in an extract either directly bind the metal ions or indirectly suppress their chelating reactivity by occupying their coordination sites [100, 101]. The ultimate outcome of the reduction reaction in the antioxidant defense system is to terminate the radical chain reactions, which may otherwise be very detrimental to tissues.

It was evident that increased extract concentrations increased the ferric reducing power of the two DCM extracts. Basically, as a result of more Fe<sup>3+</sup> being reduced to Fe<sup>2+</sup> as more electrons were being donated by antioxidant components [102, 103]. High absorbance is an indicative of the increased ferric reducing ability of the extracts. Both extracts were, however, found to have a lower reducing capacity at all the tested concentrations compared to ascorbic acid, the reference compound. This observation was starkly different from the one done by Daniel and Dluya [46] in which they demonstrated that the methanolic stem bark extract of *F. sycomorus* had a higher ferric reducing power than ascorbic acid (reference drug) which could be the case of different solvents used. Nevertheless, the findings correlate with one done by Ahoua et al. [87] on eight different *Ficus* species and it was noted that the majority had significantly lower reducing power against the ascorbic acid.

## 5. Conclusions and Recommendations

The findings of this study demonstrate that the DCM extracts of *S. henningsii* and *F. sycomorus* possess alkaloids, phenols, saponins, cardiac glycosides, flavonoids, and steroids while *F. sycomorus* contains saponins, flavonoids, alkaloids, steroids, phenols, cardiac glycosides, and terpenoids. The total phenolic and flavonoid content of the DCM stem bark extract of *F. sycomorus* was lower than that of the leaf extract of *S. henningsii*. The said phytochemicals possess antioxidant activity. The DCM extracts of *S. henningsii* and *F. sycomorus* significantly exhibited strong radical scavenging activities against hydrogen peroxide and DPPH solution at different concentrations used. The IC<sub>50</sub> values of *S. henningsii* and *F. sycomorus* were 0.325 mg/ml and 0.330 mg/ml for hydrogen peroxide and 0.068 mg/ml and 0.062 mg/ml for DPPH, respectively. Both the DCM leaf extract and stem bark extract of *S. henningsii* and *F. sycomorus* were found to have strong ferric reducing power at all the tested concentrations. Therefore, both extracts exhibited significant nonenzyme-based antioxidant activities.

Therefore, the DCM leaf extract of *S. henningsii* and stem bark extract of *F. sycomorus* studied plants can be potential antioxidant compound sources and alternatives for the management of oxidative stress. In addition, studies aimed at investigating the in vivo antioxidant efficacy of the studied plant extracts are encouraged.

## Abbreviations

ANOVA:	Analysis of variance
DCM:	Dichloromethane
DPPH:	2,2-Diphenyl-1-picrylhydrazyl
ROS:	Reactive oxidative species
SEM:	Standard error of the mean
GAE:	Gallic acid equivalent
RE:	Rutin equivalent.

## Data Availability

The data used to support the findings of this study are provided in this article. However, any additional information can be provided by the corresponding author upon request.

## Ethical Approval

All the reagents used in this study were prepared, used, and disposed of according to the set laboratory guidelines and the material safety and data sheets (MSDS).

## Disclosure

This article is part of the Kenedy Wanjala Wafula Thesis (Wanjala) [104], and no preprint version had been deposited in a journal in any form.

## Conflicts of Interest

The authors declare that they have no conflicts of interest.

## Authors' Contributions

Kenedy Wanjala Wafula, Joseph Kiambi Mworira, and Mathew Piero Ngugi designed the project, collected the specimen, performed the experiments, analyzed and interpreted data, and read and approved the manuscript for publication.

## Acknowledgments

The authors sincerely thank the members of Biochemistry, Microbiology, and Biotechnology, Kenyatta University, Nairobi, Kenya, for their unswerving support and contribution in this research project. The authors also thank Mr. Daniel Gitonga of Kenyatta University (Department of Biochemistry, Microbiology, and Biotechnology) for their technical support during the entire period of carrying out this research.



## References

- [1] H. Ushijima and R. Monzaki, "An in vitro evaluation of the antioxidant activities of necroptosis and apoptosis inhibitors: the potential of necrostatin-1 and necrostatin-1i to have radical scavenging activities," *Pharmacological Reports: PROtech*, pp. 1–8, 2023.
- [2] C. Moens, C. J. F. Muller, and L. Bouwens, "In vitro comparison of various antioxidants and flavonoids from Rooibos as beta cell protectants against lipotoxicity and oxidative stress-induced cell death," *PLoS One*, vol. 17, no. 5, Article ID e0268551, 2022.
- [3] L. N. Khanal, K. R. Sharma, Y. R. Pokharel, and S. K. Kalauni, "Phytochemical analysis and in vitro antioxidant and antibacterial activity of different solvent extracts of *Beilschmiedia roxburghiana* nees stem barks," *The Scientific World Journal*, vol. 2022, Article ID 6717012, 7 pages, 2022.
- [4] X. Chen, M. Bi, J. Yang et al., "Cadmium exposure triggers oxidative stress, necroptosis, Th1/Th2 imbalance and promotes inflammation through the TNF- $\alpha$ /NF- $\kappa$ B pathway in swine small intestine," *Journal of Hazardous Materials*, vol. 421, Article ID 126704, 2022.
- [5] N. U. Rehman, M. Shah, S. Ullah et al., "Enzymes inhibition and antioxidant potential of medicinal plants growing in Oman," *BioMed Research International*, vol. 2022, Article ID 7880387, 9 pages, 2022.
- [6] K. Tejchman, K. Kotfis, and J. Sierko, "Biomarkers and mechanisms of oxidative stress-last 20 years of research with an emphasis on kidney damage and renal transplantation," *International Journal of Molecular Sciences*, vol. 22, no. 15, p. 8010, 2021.
- [7] V. Kumar and A. Salam, "A review on reactive oxygen and nitrogen species," *Era's Journal of Medical Research*, vol. 5, no. 1, pp. 59–66, 2018.
- [8] K. Luangpraditkun, M. Tissot, A. Joompang et al., "Prevention by the natural artocarpin of morphological and biochemical alterations on UVB-induced HaCaT cells," *Oxidative Medicine and Cellular Longevity*, vol. 2021, Article ID 5067957, 13 pages, 2021.
- [9] N. Seyed Mohammad, A. A. Dehpour, M. A. Ebrahimzadeh, and N. Seyed Fazel, "Antioxidant activity of the methanol extract of *Ferula assafoetida* and its essential oil composition," *Grasas y Aceites*, vol. 60, no. 4, pp. 405–412, 2009.
- [10] L. A. Pham-Huy, H. He, and C. Pham-Huy, "Free radicals, antioxidants in disease and health," *International journal of biomedical science: IJBS*, vol. 4, no. 2, pp. 89–96, 2008.
- [11] R. Singh, S. Devi, and R. Gollen, "Role of free radical in atherosclerosis, diabetes and dyslipidaemia: larger-than-life," *Diabetes/metabolism research and reviews*, vol. 31, no. 2, pp. 113–126, 2015.
- [12] M. C. Polidori and P. Mecocci, "Modeling the dynamics of energy imbalance: the free radical theory of aging and frailty revisited," *Free Radical Biology and Medicine*, vol. 181, pp. 235–240, 2022.
- [13] A. T. Dharmaraja, "Role of reactive oxygen species (ROS) in therapeutics and drug resistance in cancer and bacteria," *Journal of Medicinal Chemistry*, vol. 60, no. 8, pp. 3221–3240, 2017.
- [14] T. Zhao, W. Wu, L. Sui et al., "Reactive oxygen species-based nanomaterials for the treatment of myocardial ischemia reperfusion injuries," *Bioactive Materials*, vol. 7, pp. 47–72, 2022.
- [15] A. Singh, R. Kukreti, L. Saso, and S. Kukreti, "Oxidative stress: a key modulator in neurodegenerative diseases," *Molecules*, vol. 24, no. 8, p. 1583, 2019.
- [16] P. Thiviya, A. Gamage, D. Piumali, O. Merah, and T. Madhujith, "Apiaceae as an important source of antioxidants and their applications," *Cosmetics*, vol. 8, no. 4, p. 111, 2021 Dec.
- [17] K. Kośmider, M. Kamieniak, S. J. Czuczwar, and B. Miziak, "Second generation of antiepileptic drugs and oxidative stress," *International Journal of Molecular Sciences*, vol. 24, no. 4, p. 3873, 2023.
- [18] M. Kumar, G. Gupta, N. P. Muhammed et al., "Toxicity ameliorative effect of vitamin E against super-paramagnetic iron oxide nanoparticles on haemato-immunological responses, antioxidant capacity, oxidative stress, and metabolic enzymes activity during exposure and recovery in *Labeo rohita* fingerlings," *Aquaculture International*, vol. 30, no. 4, pp. 1711–1739, 2022.
- [19] Y. Longobucco, A. Masini, S. Marini et al., "Exercise and oxidative stress biomarkers among adult with cancer: a systematic review," *Oxidative Medicine and Cellular Longevity*, vol. 2022, Article ID 2097318, 16 pages, 2022.
- [20] A. Roy, S. Das, I. Chatterjee, S. Roy, and R. Chakraborty, "Anti-inflammatory effects of different dietary antioxidants," in *Plant Antioxidants and Health*, pp. 1–25, Springer International Publishing, Cham, 2022.
- [21] M. Grzesik, G. Bartosz, I. Stefaniuk, M. Pichla, J. Namieśnik, and I. Sadowska-Bartosz, "Dietary antioxidants as a source of hydrogen peroxide," *Food Chemistry*, vol. 278, pp. 692–699, 2019.
- [22] F. Shahidi, "Antioxidants in food and food antioxidants," *Nahrung-Food*, vol. 44, no. 3, pp. 158–163, 2000.
- [23] B. L. Tan, M. E. Norhaizan, W. P. P. Liew, and H. Sulaiman Rahman, "Antioxidant and oxidative stress: a mutual interplay in age-related diseases," *Frontiers in Pharmacology*, vol. 9, p. 1162, 2018.
- [24] C. Rice-Evans, B. Halliwell, G. G. Lunt, and C. Rice-Evans, "Plant polyphenols: free radical scavengers or chain-breaking antioxidants?" in *Biochemical Society Symposia*, vol. 61, pp. 103–116, Portland Press, 1995, November.
- [25] E. Parthiban, C. Arokiyaraj, S. Janarthanan, and R. Ramanibai, "Antioxidant and GC-MS analysis of *Annona reticulata* leaves extract against unsecure free radicals," *SN Applied Sciences*, vol. 1, no. 4, pp. 349–9, 2019.
- [26] Y. J. Hsueh, Y. N. Chen, Y. T. Tsao, C. M. Cheng, W. C. Wu, and H. C. Chen, "The pathomechanism, antioxidant biomarkers, and treatment of oxidative stress-related eye diseases," *International Journal of Molecular Sciences*, vol. 23, no. 3, p. 1255, 2022.
- [27] H. S. El-Beltagi, H. I. Mohamed, M. I. Aldaej et al., "Production and antioxidant activity of secondary metabolites in Hassawi rice (*Oryza sativa* L.) cell suspension under salicylic acid, yeast extract, and pectin elicitation," *In Vitro Cellular and Developmental Biology - Plant*, vol. 58, no. 4, pp. 615–629, 2022.
- [28] B. Akbari, N. Baghaei Yazdi, M. Bahmaie, and F. Mahdavi Abhari, "The role of plant derived natural antioxidants in reduction of oxidative stress," *BioFactors*, vol. 48, no. 3, pp. 611–633, 2022.
- [29] H. Peng and F. Shahidi, "The effects of acyl chain length on antioxidant efficacy of mono-andmulti-acylated resveratrol: a comparative assessment," *Molecules*, vol. 27, no. 3, p. 1001, 2022.

- [30] D. I. Egodavitharana, B. V. A. S. Manori Bambaranda, and D. C. Mudannayake, "Phytochemical composition of two green seaweeds (*Ulva lactuca* and *Ulva fasciata*) and their utilization as a functional ingredient in crackers," *Journal of Aquatic Food Product Technology*, vol. 32, no. 2, pp. 158–174, 2023.
- [31] M. A. Anagnostopoulou, P. Kefalas, V. P. Papageorgiou, A. N. Assimopoulou, and D. Boskou, "Radical scavenging activity of various extracts and fractions of sweet orange peel (*Citrus sinensis*)," *Food Chemistry*, vol. 94, no. 1, pp. 19–25, 2006.
- [32] I. Khalil, W. A. Yehye, A. E. Etxeberria et al., "Nano Antioxidants: recent trends in antioxidant delivery applications," *Antioxidants*, vol. 9, no. 1, p. 24, 2019.
- [33] H. Hosseinzadeh-Bandbafha, D. Kumar, B. Singh et al., "Biodiesel antioxidants and their impact on the behavior of diesel engines: a comprehensive review," *Fuel Processing Technology*, vol. 232, Article ID 107264, 2022.
- [34] S. Rathod, S. Arya, S. Kanike, S. A. Shah, P. Bahadur, and S. Tiwari, "Advances on nanoformulation approaches for delivering plant-derived antioxidants: a case of quercetin," *International Journal of Pharmaceutics*, vol. 625, Article ID 122093, 2022.
- [35] E. Salmerón-Manzano, J. A. Garrido-Cardenas, and F. Manzano-Agugliaro, "Worldwide research trends on medicinal plants," *International Journal of Environmental Research and Public Health*, vol. 17, no. 10, p. 3376, 2020.
- [36] K. M. A. A. Naqbi, K. Karthiashwaran, S. S. Kurup, M. Abdul Muhsen Alyafei, and A. Jaleel, "Phytochemicals, proximate composition, mineral analysis and in vitro antioxidant activity of *Calligonum crinitum* (boiss)," *Horticulturae*, vol. 8, no. 2, p. 156, 2022.
- [37] L. K. Keter and P. C. Mutiso, "Ethnobotanical studies of medicinal plants used by traditional health practitioners in the management of diabetes in lower eastern province, Kenya," *Journal of Ethnopharmacology*, vol. 139, no. 1, pp. 74–80, 2012.
- [38] M. Sharifi-Rad, N. V. Anil Kumar, P. Zucca et al., "Lifestyle, oxidative stress, and antioxidants: back and forth in the pathophysiology of chronic diseases," *Frontiers in Physiology*, vol. 11, p. 694, 2020.
- [39] J. Kiambi Mworira, C. Mwit Kibiti, J. Jn Ngeranwa, and M. Piero Ngugi, "Anti-inflammatory potential of dichloromethane leaf extracts of *Eucalyptus globulus* (Labill) and *Senna didymobotrya* (Fresenius) in mice," *African Health Sciences*, vol. 21, no. 1, pp. 397–409, 2021.
- [40] B. M. Guchu, A. K. O. Machocho, S. K. Mwihi, and M. P. Ngugi, "In vitro antioxidant activities of methanolic extracts of *Caesalpinia volkensii* (Harms.), *Vernonia lasiopus* O. Hoffm., and *Acacia hockii* De Wild," *Evidence-based Complementary and Alternative Medicine*, vol. 2020, Article ID 3586268, 2020.
- [41] N. T. T. Nguyen, L. M. Nguyen, T. T. T. Nguyen, T. T. Nguyen, D. T. C. Nguyen, and T. V. Tran, "Formation, antimicrobial activity, and biomedical performance of plant-based nanoparticles: a review," *Environmental Chemistry Letters*, vol. 20, no. 4, pp. 2531–2571, 2022.
- [42] E. S. S. Abdel-Hameed, "Total phenolic contents and free radical scavenging activity of certain Egyptian *Ficus* species leaf samples," *Food Chemistry*, vol. 114, no. 4, pp. 1271–1277, 2009.
- [43] A. Walia, N. Kumar, R. Singh et al., "Bioactive compounds in *Ficus* fruits, their bioactivities, and associated health benefits: a review," *Journal of Food Quality*, vol. 2022, Article ID 6597092, 19 pages, 2022.
- [44] O. Olaoluwa, O. Taiwo, L. Nahar, and S. D. Sarker, "Ethnopharmacology, phytochemistry and biological activities of the African species of the genus *Ficus* L.," *Trends in Phytochemical Research*, vol. 6, no. 1, pp. 46–69, 2022.
- [45] N. A. Igbokwe, I. O. Igbokwe, and U. K. Sandabe, "Effect of prolonged oral administration of aqueous *Ficus sycomorus*-stem-bark extract on testicular size of growing albino rat," *International Journal of Morphology*, vol. 28, no. 4, pp. 1315–1322, 2010.
- [46] D. Daniel and T. Dluya, "In vitro biochemical assessments of methanol stem bark extracts of *Ficus sycomorus* plant," *Jordan Journal of Biological Sciences*, vol. 9, no. 1, pp. 63–68, 2016.
- [47] M. H. S. Al-Shabibi, S. S. J. Al-Touby, and M. A. Hossain, "Isolation, characterization and prediction of biologically active glycoside compounds quercetin-3-rutinoside from the fruits of *Ficus sycomorus*," *Carbohydrate Research*, vol. 511, Article ID 108483, 2022.
- [48] S. O. Oyedemi, G. Bradley, and A. J. Afolayan, "In-vitro and in-vivo antioxidant activities of aqueous extract of *Strychnos henningsii* (Gilg)," *Afr J Pharm Pharmacol*, vol. 4, no. 2, pp. 70–78, 2010.
- [49] N. Ndlovu and S. F. Van Vuuren, "The use of South African botanical species for the control of blood sugar," *Journal of Ethnopharmacology*, vol. 264, Article ID 113234, 2021.
- [50] N. M. Piero, M. N. Joan, K. M. Cromwell, N. J. Joseph, and N. M. Wilson, "Hypoglycemic activity of some Kenyan plants traditionally used to manage diabetes mellitus in eastern province," *Journal of Diabetes & Metabolism*, vol. 2, no. 8, p. 2, 2011.
- [51] A. J. Harborne, *Phytochemical Methods a Guide to Modern Techniques of Plant Analysis*, Springer science & business media, Berlin, Germany, 1998.
- [52] K. Kotake, *Practical Pharmacognosy*, Vallabh prakashan, New Delhi, India, 2000.
- [53] G. A. Spanos, R. E. Wrolstad, and D. A. Heatherbell, "Influence of processing and storage on the phenolic composition of apple juice," *Journal of Agricultural and Food Chemistry*, vol. 38, no. 7, pp. 1572–1579, 1990.
- [54] E. Lister and P. Wilson, *Measurement of Total Phenolics and ABTS Assay for Antioxidant Activity (Personal Communication)*, Crop Research Institute, Lincoln, New Zealand, 2001.
- [55] F. Medini, H. Fellah, R. Ksouri, and C. Abdely, "Total phenolic, flavonoid and tannin contents and antioxidant and antimicrobial activities of organic extracts of shoots of the plant *Limonium delicatulum*," *Journal of Taibah University for Science*, vol. 8, no. 3, pp. 216–224, 2014.
- [56] J. L. Lamaison and A. Carnat, "Levels of principal flavonoids in flowers and leaves of *Crataegus-monogyna* (jacq) and *Crataegus laevigata* (poiret) dc (rosaceae)," *Pharmaceutica Acta Helveticae*, vol. 65, no. 11, pp. 315–320, 1990.
- [57] W. Nurcholis, D. N. Sya'bani Putri, H. Husnawati, S. I. Aisyah, and B. P. Priosoeryanto, "Total flavonoid content and antioxidant activity of ethanol and ethyl acetate extracts from accessions of *Amomum compactum* fruits," *Annals of Agricultural Science*, vol. 66, no. 1, pp. 58–62, 2021.
- [58] R. J. Ruch, S. J. Cheng, and J. E. Klaunig, "Prevention of cytotoxicity and inhibition of intercellular communication by antioxidant catechins isolated from Chinese green tea," *Carcinogenesis*, vol. 10, no. 6, pp. 1003–1008, 1989.

- [59] S. Mehrotra, V. Kirar, P. Vats, S. P. Nandi, P. S. Negi, and K. Misra, "Phytochemical and antimicrobial activities of himalayan cordyceps sinensis (berk) sacc.," *Indian Journal of Experimental Biology*, vol. 53, no. 1, 2015.
- [60] W. Dong, D. Chen, Z. Chen, H. Sun, and Z. Xu, "Antioxidant capacity differences between the major flavonoids in cherry (*Prunus pseudocerasus*) in vitro and in vivo models," *Lebensmittel-Wissenschaft & Technologie*, vol. 141, Article ID 110938, 2021.
- [61] N. G. Hristova-Avakumova, E. P. Valcheva, N. O. Anastassova et al., "In vitro and in silico studies of radical scavenging activity of salicylaldehyde benzoylhydrazones," *Journal of Molecular Structure*, vol. 1245, Article ID 131021, 2021.
- [62] M. Oyaizu, "Studies on products of browning reaction antioxidative activities of products of browning reaction prepared from glucosamine," *The Japanese journal of nutrition and dietetics*, vol. 44, no. 6, pp. 307–315, 1986.
- [63] D. M. Teleanu, A. G. Niculescu, I. I. Lungu et al., "An overview of oxidative stress, neuroinflammation, and neurodegenerative diseases," *International Journal of Molecular Sciences*, vol. 23, no. 11, p. 5938, 2022.
- [64] D. J. Cherubim, C. V. Martins, L. Fariña, and R. A. Lucca, "Polyphenols as natural antioxidants in cosmetics applications," *Journal of Cosmetic Dermatology*, vol. 19, no. 1, pp. 33–37, 2020.
- [65] Q. D. Do, A. E. Angkawijaya, P. L. Tran-Nguyen et al., "Effect of extraction solvent on total phenol content, total flavonoid content, and antioxidant activity of *Limnophila aromatica*," *Journal of Food and Drug Analysis*, vol. 22, no. 3, pp. 296–302, 2014.
- [66] Y. Gong, X. Liu, W. H. He, H. G. Xu, F. Yuan, and Y. X. Gao, "Investigation into the antioxidant activity and chemical composition of alcoholic extracts from defatted marigold (*Tagetes erecta* L.) residue," *Fitoterapia*, vol. 83, no. 3, pp. 481–489, 2012.
- [67] A. C. Akinmoladun, O. E. Falaiye, O. B. Ojo, A. Adeoti, Z. A. Amoo, and M. T. Olaleye, "Effect of extraction technique, solvent polarity, and plant matrix on the antioxidant properties of *Chrysophyllum albidum* G. Don (African Star Apple)," *Bulletin of the National Research Centre*, vol. 46, no. 1, pp. 40–49, 2022.
- [68] W. C. Chen, S. W. Wang, C. W. Li et al., "Comparison of various solvent extracts and major bioactive components from *Portulaca oleracea* for antioxidant, anti-tyrosinase, and anti- $\alpha$ -glucosidase activities," *Antioxidants*, vol. 11, no. 2, p. 398, 2022.
- [69] I. Alfarrayeh, K. Tarawneh, D. Almajali, and W. Al-Awaida, "Evaluation of the Antibacterial and Antioxidant properties of the Methanolic extracts of four Medicinal plants selected from Wadi Al-Karak, Jordan related to their Phenolic contents," *Research Journal of Pharmacy and Technology*, vol. 15, no. 5, pp. 2110–2116, 2022.
- [70] K. Shimada, K. Fujikawa, K. Yahara, and T. Nakamura, "Antioxidative properties of xanthan on the autoxidation of soybean oil in cyclodextrin emulsion," *Journal of Agricultural and Food Chemistry*, vol. 40, no. 6, pp. 945–948, 1992.
- [71] Z. Haida and M. Hakimian, "A comprehensive review on the determination of enzymatic assay and nonenzymatic antioxidant activities," *Food science & nutrition*, vol. 7, no. 5, pp. 1555–1563, 2019.
- [72] A. A. Mohamed, A. A. Khalil, and H. E. S. El-Beltagi, "Antioxidant and antimicrobial properties of kaff maryam (*Anastatica hierochuntica*) and doum palm (*Hyphaene thebaica*)," *Grasas y Aceites*, vol. 61, no. 1, pp. 67–75, 2010.
- [73] M. A. Al-Mamary and Z. Moussa, "Antioxidant activity: the presence and impact of hydroxyl groups in small molecules of natural and synthetic origin," *Antioxidants—Benefits, sources, mechanisms of action*, pp. 318–377, 2021.
- [74] M. M. Namadina, H. Haruna, and U. Sanusi, "Pharmacognostic, antioxidant and acute toxicity study of *Ficus sycomorus* (linn)(moraceae) root and stem bark," *Fudma journal of sciences*, vol. 4, no. 2, pp. 605–614, 2020.
- [75] J. B. Johnson, D. A. Broszczak, J. S. Mani, J. Anesi, and M. Naiker, "A cut above the rest: oxidative stress in chronic wounds and the potential role of polyphenols as therapeutics," *Journal of Pharmacy and Pharmacology*, vol. 74, no. 4, pp. 485–502, 2022.
- [76] N. Huda-Faujan, A. Noriham, A. S. Norrakiah, and A. S. Babji, "Antioxidant activity of plants methanolic extracts containing phenolic compounds," *African Journal of Biotechnology*, vol. 8, no. 3, 2009.
- [77] Y. Li, Z. J. Qian, B. Ryu, S. H. Lee, M. M. Kim, and S. K. Kim, "Chemical components and its antioxidant properties in vitro: an edible marine brown alga, *Ecklonia cava*," *Bioorganic & Medicinal Chemistry*, vol. 17, no. 5, pp. 1963–1973, 2009.
- [78] A. Rafat, K. Philip, and S. Muniandy, "Antioxidant potential and content of phenolic compounds in ethanolic extracts of selected parts of *Andrographis paniculata*," *Journal of Medicinal Plants Research*, vol. 4, no. 3, pp. 197–202, 2010.
- [79] F. Hesam, G. R. Balali, and R. T. Tehrani, "Evaluation of antioxidant activity of three common potato (*Solanum tuberosum*) cultivars in Iran," *Avicenna journal of phyto-medicine*, vol. 2, no. 2, pp. 79–85, 2012.
- [80] T. J. Zhou, Y. Xu, L. Xing, Y. Wang, and H. L. Jiang, "A harmless-harmful switchable and uninterrupted laccase instructed killer for activatable chemodynamic therapy," *Advanced Materials*, vol. 33, no. 27, Article ID 2100114, 2021.
- [81] T. J. Costa, P. R. Barros, C. Arce et al., "The homeostatic role of hydrogen peroxide, superoxide anion and nitric oxide in the vasculature," *Free Radical Biology and Medicine*, vol. 162, pp. 615–635, 2021.
- [82] R. L. Li, Q. Zhang, J. Liu et al., "Hydroxy- $\alpha$ -sanshool possesses protective potentials on H<sub>2</sub>O<sub>2</sub>-stimulated PC1<sub>2</sub> cells by suppression of oxidative stress-induced apoptosis through regulation of PI3K/Akt signal pathway," *Oxidative Medicine and Cellular Longevity*, vol. 12, Article ID 3481758, 2020.
- [83] C. M. C. Andrés, J. M. Pérez de la Lastra, C. A. Juan, F. J. Plou, and E. Pérez-Lebeña, "Chemistry of hydrogen peroxide formation and elimination in mammalian cells, and its role in various pathologies," *Stresses*, vol. 2, no. 3, pp. 256–274, 2022.
- [84] J. C. Lee, H. R. Kim, J. Kim, and Y. S. Jang, "Antioxidant property of an ethanol extract of the stem of *Opuntia ficus-indica* var. *saboten*," *Journal of Agricultural and Food Chemistry*, vol. 50, no. 22, pp. 6490–6496, 2002.
- [85] N. Sirisha, M. Sreenivasulu, K. Sangeeta, and C. M. Chetty, "Antioxidant properties of *Ficus* species-a review," *International journal of pharmtech research*, vol. 2, no. 4, pp. 2174–2182, 2010.
- [86] P. Sharayei, E. Azarpazhooh, S. Zomorodi, and H. S. Ramaswamy, "Ultrasound assisted extraction of bioactive compounds from pomegranate (*Punica granatum* L.) peel," *Lebensmittel-Wissenschaft & Technologie*, vol. 101, pp. 342–350, 2019.

- [87] A. R. C. Ahoua, M. W. Kone, A. G. Konan, F. H. T. Bi, and B. Bonfoh, "Antioxidant activity of eight plants consumed by great apes in Côte d'Ivoire," *African Journal of Biotechnology*, vol. 11, no. 54, pp. 11732–11740, 2012.
- [88] I. Varzaru, A. E. Untea, and M. Saracila, "In vitro antioxidant properties of berry leaves and their inhibitory effect on lipid peroxidation of thigh meat from broiler chickens," *European Journal of Lipid Science and Technology*, vol. 122, no. 4, Article ID 1900384, 2020.
- [89] P. Deo, E. Hewawasam, A. Karakoulakis et al., "In vitro inhibitory activities of selected Australian medicinal plant extracts against protein glycation, angiotensin converting enzyme (ACE) and digestive enzymes linked to type II diabetes," *BMC Complementary and Alternative Medicine*, vol. 16, no. 1, pp. 435–511, 2016.
- [90] F. C. Saputri and I. Jantan, "Effects of selected medicinal plants on human low-density lipoprotein oxidation, 2, 2-diphenyl-1-picrylhydrazyl (DPPH) radicals and human platelet aggregation," *Journal of Medicinal Plants Research*, vol. 5, no. 26, pp. 6182–6191, 2011.
- [91] M. Mahmudzadeh, H. Yari, B. Ramezanzadeh, and M. Mahdavian, "Highly potent radical scavenging-antioxidant activity of biologically reduced graphene oxide using Nettle extract as a green bio-genic amines-based reductants source instead of hazardous hydrazine hydrate," *Journal of Hazardous Materials*, vol. 371, pp. 609–624, 2019.
- [92] L. R. Fukumoto and G. Mazza, "Assessing antioxidant and pro-oxidant activities of phenolic compounds," *Journal of Agricultural and Food Chemistry*, vol. 48, no. 8, pp. 3597–3604, 2000.
- [93] W. Arika, C. M. Kibiti, J. M. Njagi, and M. P. Ngugi, "In vitro antioxidant properties of dichloromethanolic leaf extract of *Gnidia glauca* (Fresen) as a promising anti-obesity drug," *Journal of Evidence-Based Integrative Medicine*, vol. 24, Article ID 2515690X1988325, 2019.
- [94] O. O. Igbinosa, I. H. Igbinosa, V. N. Chigor et al., "Polyphenolic contents and antioxidant potential of stem bark extracts from *Jatropha curcas* (Linn)," *International Journal of Molecular Sciences*, vol. 12, no. 5, pp. 2958–2971, 2011.
- [95] J. A. Y. Kambli, A. Patil, and R. Keshava, "Phytochemical screening and evaluation of antibacterial, antioxidant and cytotoxic activity of *Ficus racemosa* Linn," *International Journal of Pharmacy and Pharmaceutical Sciences*, vol. 6, no. 4, pp. 1–8, 2014.
- [96] L. A. Santiago and A. B. R. Mayor, "Prooxidant effect of the crude ethanolic leaf extract of *Ficus odorata* Blanco Merr. in vitro: its medical significance," *International Journal of Biotechnology and Bioengineering*, vol. 8, pp. 53–60, 2014.
- [97] M. Saleem, "Natural products as antimicrobial agents—an update," *Novel Antimicrobial Agents and Strategies*, vol. 46, pp. 219–294, 2014.
- [98] B. Jasiewicz, W. Kozanecka-Okupnik, M. Przygodzki et al., "Synthesis, antioxidant and cytoprotective activity evaluation of C-3 substituted indole derivatives," *Scientific Reports*, vol. 11, no. 1, 14 pages, Article ID 15425, 2021.
- [99] A. N. M. Ramli, N. W. A. Manap, P. Bhuyar, and N. I. W. Azelee, "Passion fruit (*Passiflora edulis*) peel powder extract and its application towards antibacterial and antioxidant activity on the preserved meat products," *SN Applied Sciences*, vol. 2, no. 10, pp. 1748–1811, 2020.
- [100] H. O. Edeoga, D. E. Okwu, and B. O. Mbaebie, "Phytochemical constituents of some Nigerian medicinal plants," *African Journal of Biotechnology*, vol. 4, no. 7, pp. 685–688, 2005.
- [101] O. V. Njoku and C. Obi, "Phytochemical constituents of some selected medicinal plants," *African Journal of Pure and Applied Chemistry*, vol. 3, no. 11, pp. 228–233, 2009.
- [102] K. Mitra and N. Uddin, "Total phenolics, flavonoids, proanthocyanidins, ascorbic acid contents and in-vitro antioxidant activities of newly developed isolated soya protein," *Discourse Journal of Agriculture and Food Sciences*, vol. 2, no. 5, pp. 160–168, 2014.
- [103] F. A. Alsulaymani, M. F. Elmhdwi, S. Gaber, N. M. El Aali, M. I. Mohammed, and A. A. Abdualsalam, "In vitro antioxidant and antibacterial activity of olive leave extract," *Journal of Pharmacy and Applied Chememistry*, vol. 7, no. 2, pp. 75–80, 2021.
- [104] W. K. Wanjala, *In-Vitro Antioxidant Activities of Dichloromethane Doctoral Dissertation*, Kenyatta University, Nairobi, Kenya, 2019.