

Research article

Open Access

The risk of pathogenic intestinal parasite infections in Kisii Municipality, Kenya

Robert M Nyarango*^{1,2}, Peninah A Aloo², Ephantus W Kabiru³ and Benson O Nyanchongi^{1,4}

Address: ¹Department of Health Sciences, Gusii Institute of Science and Technology, P.O. Box 222-40200 Kisii, Kenya, ²Department of Zoological Sciences, Kenyatta University P.O. Box 43844-01000 Nairobi, Kenya, ³Department of Clinical Pathology, Kenyatta University P.O. Box 43844-0100 Nairobi, Kenya and ⁴Department of Experimental Medicine and Public Health, Via Camillo Lili n. 55-62032 Camerino (MC), Italy

Email: Robert M Nyarango* - rnyarango6@yahoo.com; Peninah A Aloo - aloopenina@yahoo.com; Ephantus W Kabiru - ewkabiru@yahoo.com; Benson O Nyanchongi - benson.nyanchongi@unicam.it

* Corresponding author

Published: 14 July 2008

Received: 17 November 2007

BMC Public Health 2008, 8:237 doi:10.1186/1471-2458-8-237

Accepted: 14 July 2008

This article is available from: <http://www.biomedcentral.com/1471-2458/8/237>

© 2008 Nyarango et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Intestinal parasitic infections are among the most common infections worldwide. Various epidemiological studies indicate that the prevalence of intestinal parasites is high especially in developing countries, although in many of these, the environmental risk factors have not been clearly elucidated. The objective of this study was to determine the risk of pathogenic intestinal parasites infections in Kisii Municipality.

Methods: Random sampling was used in the selection of the study samples. Stool parasitological profiles of food handlers were done by direct smear and formalin-ethyl acetate sedimentation method. Both vegetable and meat samples were examined for the presence of intestinal parasites. The storage and meat handling practices of the various butchereries were observed.

Results: Types of samples examined for occurrence of intestinal parasites includes, a total of 84 vegetable, 440 meat and 168 stool samples. Fifty five (65.5%) vegetable, 334 (75.9%) meat and 69 (41.1%) of the stool samples were found positive for intestinal parasites indicating a high overall risk (66.18%) for intestinal parasite infections. Of the parasites detected, the most common parasites infesting the foodstuffs and infecting the food handlers were *Ascaris lumbricoides* and *Entamoeba histolytica*. Parasites were significantly less likely to be present on meat that was refrigerated during display than meat that was displayed at ambient temperature.

Conclusion: There is a high risk of infection with intestinal parasites in the sampled Municipal markets. About half of the food handlers surveyed (41.1 %) at the Municipal Hospital had one or more parasitic infections. Furthermore, meat (65.5%) and vegetables (75.9%) sold at the Municipal market were found to be contaminated with parasites hence the inhabitants requires a need for education on food safety, good distribution practices and improvement on sanitary conditions.

Background

Intestinal parasites cause significant morbidity and mor-

tality throughout the world, particularly in developing countries and in persons with comorbidities. The risk of

intestinal parasites infections is measured using the DALY (disability- adjusted life year) and one DALY represents the loss of one year of equivalent full health [1]. The resulting diseases have socio-economic impact in terms of high treatment costs per DALY and hospitalization costs [2]. Globally, millions of people suffer from parasitic infections such as *Ascaris lumbricoides* (1.2 billion), *Trichuris trichiura* (795 million), hookworms (*Ancylostoma duodenale* and *Necator americanus*) (740 million) [3], *Entamoeba histolytica* (50 million) [4] and *Giardia lamblia* (2.8 million) [5]. In man, intestinal parasites are significantly associated with diarrhoea [6]. The fecal oral route is significant in the transmission of parasitic infections to humans via poor personal hygiene [7], environmental conditions like contamination of soil and water sources with human faeces [8] and poor sewage disposal such as use of night soil as fertilizer [9]. When the soil becomes contaminated, the eggs in soil can be transferred onto vegetables then onto the hands and transferred directly into the mouth [10] or ingested by eating raw vegetables [9]. The risk of intestinal parasites is even higher among the inhabitants of towns of the developing countries especially in the shanties and slums where there is poor disposal of garbage, poor health systems and overcrowding [11]. Intestinal parasites have been found to adhere to vegetables, fruits, fingers, utensils, door handles [12] and money [11]. Additionally, they can be transmitted through flies [13] and contaminated fingernails [14].

Epidemiological surveys done in Kenya's poor periurban and urban school children revealed a high prevalence of intestinal parasitic infections with *Ascaris lumbricoides* (82%), *Trichuris trichiura* (60%), *Entamoeba histolytica* (41%) and *Giardia lamblia* (30%) [15]. Recent studies done at various locations in Kisii Highlands, Western Kenya, showed that there was high prevalence of many intestinal parasites such as *A. lumbricoides* (10%), hookworm (4%) and *T. trichiura* (0.1%) [16]. The Kisii Municipal markets are characterized by a lack of bathrooms and washing facilities, poor sanitary conditions of latrines which lack water supplies and a frequent presence of piles of garbage that provide a fertile environment for transmission of intestinal parasites. The objective of this study was to determine the prevalence of intestinal parasites among food handlers and foodstuffs sold at Kisii Municipality.

Methods

Study area

The study was carried out between December 2004 and June 2005 in Kisii Municipality, which is located at the southern end of the western Kenya highlands at an altitude of 1660 m above sea level. The area receives average rainfall of over 1500 mm per annum distributed almost throughout the year although there are two rainy seasons (March to May and October to November). Temperatures

range from 10°C to 30°C with relative humidity of 88%. The area is densely populated with a population of 37,531 people and a density of 1295 people/km² [17]. Majority of the population depends on open air markets as a source of both meat and vegetable foodstuffs although there is poor drainage and sanitation characterized by presence of refuse dump sites nearby. The parasitological survey of the food handlers and foodstuffs was carried out at the Kisii District Hospital.

Study population

A correlative descriptive study design was used to determine the association between various risk factors and the occurrence of intestinal parasites. Two open air markets were randomly selected by lottery method from the list of the five markets of Kisii Municipality. The sample size for foodstuffs and food handlers was calculated on a prevalence of 10%, $d = 0.05$ at a confidence level of 95%. Each of the four varieties of vegetables commonly sold in the study area was selected by random sampling from the vendors in the open air markets. Samples of vegetables leaves weighing 250 g for each variety (kales (*Brassica oleracea* var. *acephala*), cabbage (*Brassica oleracea capitata*), spider flower (*Gynandropsis gynandra*) and black nightshade (*Solanum nigrum*)) were collected each day from the vendors by random sampling. The leaves of the four vegetable varieties were chosen because they are the major ones eaten by residents of Kisii Municipality. Twenty two butcheries were randomly selected using lottery method from using a list of the total of 125 butcheries in the Municipality. The asymptomatic adult male and female food handlers from the sampled markets and butcheries were randomly selected by lottery method from a list of attendance at Kisii District Hospital for routine examination of stool parasitological profile according to the Kenyan Ministry of Health Regulations (Public Health act Chapter 242) [18]. The meat (beef, mutton and goat) samples each weighing 250 g was collected from the 22 butcheries selected. The samples were collected weekly, between 9.00 a.m. and 12.00 mid day. The meat handling practices at the butcheries were observed on every sampling occasion using a checklist [see Additional file 1] and the observations recorded. However, this study did not examine parasitological profile of slaughter houses and hand washes of foodstuff vendors. Each of the food handler selected was given a plastic stool container and asked to bring bean-sized stool sample within 24 hours. This study was approved by the research and ethics committee of Kenyatta University as well as the Ministry of Science and Technology, Kenya. Consent of the study participants was obtained before taking the samples for stool examination. The food handlers alternate between the five markets in Kisii Municipality, although the two markets chosen in this study are the major ones. Double counting was

avoided by asking the food handlers if they had previously been examined for intestinal parasite profile.

Sample analysis

A 250 g sample of each vegetable or meat sample was examined for intestinal parasite profile (described by [12]). The sample was washed in distilled water and the suspension was strained through a sterile sieve to remove undesirable materials. The filtrate was centrifuged and supernatant discarded while the deposit was suspended in magnesium sulphate floatation fluid of specific gravity 1.3 and recentrifuged. The floatation fluid was filled to the brim and a cover slip was superimposed. The cover slip was lifted and examined under a light microscope. The cysts and eggs of various parasite species present were identified [19]. Each parasite eggs, larvae or cysts present in the samples were counted and densities of each species were expressed as "many" (>three cysts per high-power field; >20 eggs or larvae per mount); "moderate"(two cysts per high-power field; 10 to 19 eggs or larvae per mount); "few" (one cyst per high-power field; three to nine eggs or larvae per mount); and "rare" (two to five cysts and <two eggs or larvae per mount). For simplification, numerical values were assigned to each density: many, 4; moderate, 3; few, 2; rare, 1; and none, 0 [20]. The food handling practices of the butchereries including the meat storage method(s), the handlers' hygiene standards and the presence of houseflies on meat samples were observed using a check list and recorded during each sampling day. A direct saline smear preparation of the stool sample specimens obtained from the food handlers were made for examination of trophozoites, ova and cysts of intestinal parasites using Lugol's Iodine solution and formalin-ethyl acetate sedimentation method [21].

Statistical analysis

A computer program (SPSS 11.5 for Windows) was used for data analysis. Multiple logistic regression analysis was used. The differences were considered to be statistically significant when the p-value obtained was less than 0.05

Results

Parasites in the foodstuffs

A total of eighty four vegetable samples comprising of 21 samples for each species were collected, kales cabbage,

spider flower and black night shade. In all, 55 (65.5%) vegetable samples were infested with intestinal helminthes. Eleven (52.4%) of kales, 12 (57.1%) cabbages, 17 (81.0%) spider flower and 15 (71.4%) of black night shade tested positive. An average parasite score density of 2.37 was observed in the vegetables (Table 1). A statistically significant difference was observed between infestation rates of intestinal parasites and types of vegetables (p = 0.000)).

The risk factors (independent variables) influencing infection with intestinal parasites (dependent variable) were investigated in 22 butchereries (Table 2). These were the meat storage methods, presence or absence of cashier, the presence or absence of protective clothing and the presence or absence of houseflies on meat samples. The storage methods used included placing the meat in refrigerator 4 butchereries (18.2%), placing the meat on the surface of the table 10 (45.5%), and hanging the meat in a wire mesh 8 (36.4%). In the meat stored in a refrigerator, there were 15 (18.8%) samples with intestinal parasites, while those placed in the open surface had 73 (45.6%) samples and wire mesh placed had 73 (36.5%) samples with intestinal parasites. Parasites were significantly less likely to be present on meat that was refrigerated during display than meat that was displayed at ambient temperature (p = 0.002).

Fifteen (68.2%) butchereries had the same person handling the meat and the cash (no cashier) while seven (31.8%) had a food handler and a cashier. There was a significantly high prevalence of intestinal parasites in meat samples where there was no cashier 154 (59.2%) than where the cashier was present 74 (40.9%) (p = 0.000).

The prevalence of intestinal parasites in meat samples collected from (47.1%) whereas those present from butchereries where personnel were not wearing protective clothing were 233 (52.9%).

The numbers of butchereries where there were houseflies on the meat samples collected were 13 (59.1%) while those that did not have houseflies were 9 (40.9%). The average parasite score density in the meat samples was ranging from 1.40 to 1.87.

Table 1: The prevalence and density of intestinal parasites among vegetable foods samples.

Samples	n	Frequency of intestinal parasites (%)	Score of parasite density
Kales	21	11(52.4)	2.43
Cabbage	21	12(57.1)	1.86
Spider flower	21	17(81)	2.57
Black night shade	21	15(71.4)	2.62
Overall	84	55(65.5)	2.37

Table 2: The relationship between risk factors for meat handling practices and prevalence of intestinal parasites.

Risk factor	No of samples	Frequency of intestinal parasites (%)	Score of parasite density
Storage Method			
Refrigerator	80	31 (7.0)	1.40
Open surface	200	165 (37.5)	1.87
Wire mesh	160	138 (31.4)	1.63
Cashier			
Absent	300	263 (59.8)	1.85
Present	140	71(16.1)	1.64
Protective clothing			
Present	200	134(30.5)	1.50
Absent	240	200(45.5)	1.61
Houseflies			
Present	300	263(59.8)	1.83
Absent	140	71(16.1)	1.64

Parasitological profiles of food handlers

Stool specimens were collected from the food handlers attending the Kisii District Hospital. Out of 168 food handlers, 69 (41.1%) were infected with one or more intestinal parasites of whom 27 (16.1%) were infected with one species of protozoan, 37 (22.0%) were infected with one species of helminth, 5(3.0%) had mixed infections, with 3 (1.8%) having *Entamoeba histolytica* and *Ascaris lumbricoides*, and 2 (1.2%) had *E. histolytica* and *Giardia lamblia* (Table 3). Of the most common intestinal parasite species were *A. lumbricoides* 22 (13.1%) and *E. histolytica* 20 (11.9%). The others were hookworms 13 (7.7%), *G. lamblia* 6 (3.6%), *Trichuris trichiura* 2 (1.2%) and *Balantidium coli* 1(0.6%), in their decreasing order of prevalence. There was a statistically significant difference between the numbers of the various intestinal parasite species among the food handlers ($p < 0.010$). The parasite score density in the faecal specimens observed ranged between 1.50 for

Trichuris trichiura whilst highest parasite score density was found among multiple infections with *Entamoeba histolytica* and *Ascaris lumbricoides* with 2.63.

Discussion

The study showed high intestinal parasitic infestation of both meat and vegetable foodstuffs as 65.5 % and 75.9 %, respectively. Additionally, the burden of infection with intestinal parasites among the food handlers was almost half (41%). The presence of moderate parasite score densities in the foodstuffs and food handlers indicate high transmission risks of such parasites to inhabitants of Kisii Municipality. These findings indicate a public health priority and strongly support the need for education on food safety, Good Manufacturing Practice (GMP), Hazard Analysis Critical Control Point (HACCP), improvement of sanitation conditions in and around the market by waste collection, management and handling of foodstuffs so as

Table 3: Distribution of intestinal parasites among examined food handlers.

Parasite class/species	Number of infected cases (n)	Infection rates (%)	% Of those examined. N = 168	Score of parasite density
Single Infections				
Protozoa				
<i>Entamoeba histolytica</i>	20	74.1	11.9	2.50
<i>Giardia lamblia</i>	6	22.2	3.6	1.67
<i>Balantidium coli</i>	1	3.70	0.6	2.00
Sub Total protozoa	27	100	16.1	
Helminthes				
<i>Ascaris lumbricoides</i>	22	59.5	13.1	2.55
Hookworms	13	35.1	7.7	2.38
<i>Trichuris trichura</i>	2	5.4	1.2	1.50
Sub Total Helminthes	37	100	22.0	
Multiple infections				
<i>E. histolytica</i> + <i>A. lumbricoides</i>	3	60.0	1.8	2.63
<i>E. histolytica</i> + <i>G. lamblia</i>	2	40.0	1.2	2.56
Sub -total mixed infections	5	100	3.0	
Overall total of individuals	69		41.1	

to reduce the prevalence of infections in both food handlers and food stuffs sold at the market.

Examination of vegetable samples revealed that all were equally highly contaminated with infestation rates of 52.4%, 57.1%, 81.0% and 71.4% respectively for kales, cabbage, spider flower, and black nightshade. This may be attributed to handling techniques of the vegetables [9]. The intestinal parasites found on the vegetable samples include protozoa: *Entamoeba histolytica*, *Giardia lamblia* and *Balantidium coli* and helminthes: *Ascaris lumbricoides*, *Trichuris trichiura* and hookworms. The risk of infection with intestinal parasites to the population is increased because these contaminated vegetables are sometimes eaten raw, undercooked to retain the natural taste and preserve heat-labile nutrients, or unclean [22]. Additionally, vegetables purchased at urban markets have been found to have higher rates of infestation with intestinal parasites [23].

The type of meat storage practice influenced the prevalence of intestinal parasites, with the refrigerator having least number (7%) while both meat placed in wire mesh and on open surface had higher number of intestinal parasites, 31.4% and 37.5% respectively. This indicates that proper storage methods like refrigeration should be used to minimize risk of contamination as only 18.1% of the butcheries sampled used refrigeration. Poor storage methods expose meat to mechanical vectors like houseflies, cockroaches and rats that transfer eggs, cysts of intestinal parasites to improperly stored meat [13]. Additionally, strong winds may blow eggs of intestinal parasites in dust to the exposed meat [24]. The butcheries where the handlers handled money while serving meat had 59.8% parasites prevalence while those where different personnel handled money and the meat serving had low prevalence (16.1%), this is major factor enhancing the risk of acquiring intestinal parasites as shown in other studies [11,25]. Besides, the cysts of intestinal parasites have been found to adhere on meat carcasses [26], this enhances the risk of transmission of intestinal parasites especially where meat is undercooked or eaten raw.

The most common intestinal parasites affecting the food handlers were *A. lumbricoides* (13.1%) and *E. histolytica* (11.9%), as was observed in Kisii District [16] although the study did not show intestinal protozoans. This contrasts sharply with earlier studies done in poor periurban and urban communities in Nairobi that had *A. lumbricoides* prevalence of 82% [27]. The prevalence of infection with intestinal parasites among the food handlers was almost half (41.1%) of those examined, with 3% having multiple protozoan and helminth infections. This high prevalence is a risk to patrons especially if the handlers fail to adequately sanitize hands, use food handling tools

(tongs, spoons, utensils or bakery or serving papers) and handle money while serving food [28]. Foods sold in markets may be contaminated from hands that have not been washed after defecation or from flies that land on both food and faeces hence increasing risks of transmission of intestinal parasites to consumers [29].

Conclusion

About half of the people surveyed (41.1 %) at the Municipal Hospital had one or more parasitic infections. Furthermore, meat (65.5%) and vegetables (75.9%) sold at the Municipal market were found to be contaminated with parasites. The handlers of meat should employ proper storage methods like refrigeration, employ different personnel to handle money and serve meat to consumers. Additionally, they should use insecticides to kill the houseflies. Media programmes should be launched to enlighten the public on the necessity of good sanitation hygiene and risks of acquiring intestinal parasites.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

RMN designed, performed sampling, parasitological examination and manuscript development. PAA Contributed in planning the research, discussing the results and revising the manuscript. EWK participated in the study design, coordinated the study and revised the manuscript. BON conducted statistical analysis, discussing the results and developing the manuscript. All authors read and approved the final manuscript.

Acknowledgements

The authors wish to thank the people of Kisii Municipality who participated in the study and the staff of the Ministry of Health at Kisii District Hospital, who assisted in this study. We particularly appreciate the work by Henry Sendora during specimen examination and Dr Wycliffe Mogo for his technical and logistical support. Special thanks to the Teachers Service Commission (Kenya) for granting a study leave to RMN. This study was supported by Kisii District Hospital and Gusii Institute of Science and Technology and we thank the two institutions.

References

1. Murray CJL, Lopenz AD: **The global burden of disease: a comprehensive assessment of mortality and disability from diseases, injuries, and risk factors in 1990 and projected to 2020.** Cambridge (MA): Harvard University Press; 1996.
2. Utzinger J, Keiser J: **Schistosomiasis and soil-transmitted helminthiasis: common drugs for treatment and control.** *Expert Opin Pharmacother* 2004, **5**:263-285.
3. de Silva NR, Brooker S, Hotez PJ, Montresor A, Engels D, Savioli L: **Soil transmitted helminth infections: updating the global picture.** *Trends Parasitol* 2003, **19**:547-551.
4. Samuel L, Stanley Jr, Sharon L: **Reed: Microbes and Microbial Toxins: Paradigms for Microbial-Mucosal Interactions. VI Entamoeba histolytica: parasite-host interactions.** *Am J Physiol Gastrointest Liver Physiol* 2001, **280**:1049-1054.
5. Ali SA, Hill DR: **Giardia intestinalis.** *Curr Opin Infect Dis* 2003, **16**:453-460.

6. Utzinger J, N'Goran EK, Marti HP, Tanner M, Lengeler C: **Intestinal amoebiasis, giardiasis and geohelminthiasis: their association with other intestinal parasites and reported intestinal symptoms.** *Trans R Soc Trop Med Hyg* 1999, **93**:137-141.
7. Okyay P, Ertug S, Gultekin B, Onen O, Beser E: **Intestinal parasites prevalence and related factors in school children, a western city sample-Turkey.** *BMC Public Health* 2004, **4**:64. doi: 10.1186/1471-2458-4-64
8. Muttalib MA, Huq M, Huq JA, Suzuki N: **Soil pollution with Ascaris ova in three villages of Bangladesh.** *Yokogawa, collected paper on the control of soil transmitted helminthiasis, APCO, Tokyo* 1983, **11**:66-71.
9. Mustafa U, Adnan S, Gonul A, Hatice O, Suleyman A: **Environmental Pollution with Soil-transmitted Helminths in Sanliurfa, Turkey.** *Mem Inst Oswaldo Cruz, Rio de Janeiro* 2001, **96**(7):903-909.
10. Koyabashi A: **Ascaris.** *JICA, Textbook for Seminar on Parasite Control Administration for Senior Officers – A Step Towards Primary Health Care, Tokyo* 1999, **5**:233-242.
11. Crompton DWT, Savioli L: **Intestinal parasitic infections and urbanization.** *Buletin of the World Health Organization* 1993, **71**(1):1-7.
12. Kagei N: **Techniques for the measurement of environmental pollution by infective stage of soil transmitted helminths.** *M Yokogawa, Collected Paper on the Control of Soil Transmitted Helminthiasis, APCO, Tokyo* 1983, **11**:27-46.
13. Monzon RB, Sanchez AR, Tadiaman BM, Najos OA, Valencia EG, Rueda RR, Ventura JV: **A comparison of the role of Musca domestica (Linnaeus) and Chrysomya megacephala (Fabricius) as mechanical vectors of helminthic parasites in a typical slum area of metropolitan manila.** *Southeast Asian J Trop Med Public Health* 1991, **22**:222-8.
14. Ismid S, Rukmono B: **Nail and dust examination for helminth eggs in orphanages.** In *Collected papers on the control of soil transmitted helminthiasis II Volume 5.* Tokyo. Asian Parasite Control Organisation; 1983:1-53.
15. Rijsptra AC: **Results of duplicated series of stool examinations for all intestinal parasites by different methods in school children in East Africa with remarks on serological aspects of amoebiasis and schistosomiasis.** *Annal Soc Belge Med Trop* 1975, **55**:415-425.
16. Akhwale SW, Koji JL, Kaneko A, Eto H, Obonyo C, Bjorkman A, Kobayakawa T: **Anaemia and malaria at different altitudes in the western highlands of Kenya.** *Acta Tropica* 2004, **91**:167-175.
17. CBS: **1999 Population and housing census: Counting people for our development, Population distribution by Administration and Urban Centres.** *Central Bureau of Statistics, Ministry of Finance, Government of Kenya, Nairobi* 2001, **1**.
18. Republic of Kenya: **Laws of Kenya, The Public Health Act Chapter 242, Revised edition.** *Government of Kenya, Nairobi* 1972.
19. Suzuki N: **Colour Atlas for Human Helminth Eggs.** APS & JOICFP, Hoken Kaikan Group, Tokyo; 1981:44-56.
20. Young KH, Bullock SL, Melvin DM, Spruill CL: **Ethyl acetate as a substitute for diethyl ether in the Formalin-ether sedimentation technique.** *J Clinical Microbiol* **10**:852-853.
21. Truant AL, Elliott SH, Kelly MT, Smith JH: **Comparison of formalin-ethyl ether sedimentation, formalin-ethyl acetate sedimentation, and zinc sulfate flotation techniques for detection of intestinal parasites.** *Journal of Clinical Microbiology* 1981, **13**(5):882-884.
22. Slifko TR, Smith HV, Rose JB: **Emerging parasite zoonoses associated with water and food.** *International Journal for Parasitology* 2000, **30**:1379-1393.
23. Cifuentes E, Gomez M, Blumenthal U, Tellez-Rojo MM, Romieu I, Ruiz-Palacios G, Ruiz-Velazco S: **Risk factors for Giardia intestinalis infection in agricultural villages practicing wastewater irrigation in Mexico [abstract].** *Am J Trop Med Hyg* 2000, **62**(3):388-392.
24. Kunii C: **It all started from worms. The 45-year record of Japan's post-World War II national health and family planning movement.** Tokyo: The Hoken Kaikan Foundation; 1992.
25. Singh LA, Das CS, Baruah I: **Observations on the soil contamination with zoonotic canine gastrointestinal parasites in selected areas of Tezpur, Assam, India.** *Journal of parasitic diseases* 2004, **28**(2):121-123.
26. Rajinish S, Sharma DK, Juyal PD, Sharma JK: **Epidemiology of Taenia solium cysticercosis in pigs of Northern Punjab, India.** *Journal of Parasitic Diseases* 2004, **28**(2):124-126.
27. World Health Organization: **Control of Tropical Diseases.** World Health Organization Geneva; 1998:201.
28. Michaels B: **Handling money and serving ready-to-eat food Food Service Technology.** 2002, **2**(1):1.
29. Nichols GL: **Food borne protozoa.** *Br Med Bull* 1999, **55**(4):209-235.

Pre-publication history

The pre-publication history for this paper can be accessed here:

<http://www.biomedcentral.com/1471-2458/8/237/prepub>

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

