

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/372915002>

Numerical Investigation of HIV/AIDS Dynamics Among the Truckers and the Local Community at Malaba and Busia Border Stops

Article · June 2023

DOI: 10.5923/j.ajcam.20231301.02

CITATIONS

0

READS

10

1 author:



Ancent Makau Kimulu
Kenyatta University

5 PUBLICATIONS 13 CITATIONS

[SEE PROFILE](#)

Numerical Investigation of HIV/AIDS Dynamics Among the Truckers and the Local Community at Malaba and Busia Border Stops

Ancent Makau Kimulu

Department of Mathematics and Actuarial Science, Kenyatta University, Nairobi, Kenya

Abstract Busia and Malaba are the busiest border crossing points for truckers using the Northern Corridor connecting landlocked countries of Uganda, DR Congo, South Sudan and parts of Rwanda to Mombasa port. On average, Malaba and Busia clears 1000 and 600 trucks respectively per day. Delays in clearance of the trucks causes long queues at crossing points and influx of truckers in these respective towns. The truckers spent this time in the company of commercial sex workers and interact with the local communities in these towns, hence this is a conduit of spread of HIV/AIDS in these regions. This paper designed a model to study the HIV/AIDS dynamics in this border towns. The reproduction number, the disease-free equilibrium and endemic equilibrium points determined using the Next Generation Matrix (NGM) method. From the analysis of the model, it is found that delays in clearance time increases the force of infections from females to males. It is found out that also found out that increase in force if infections to males from females increases the male HIV infections. This is due to prolonged time of sexual interactions as the trucks await clearance and also due to higher number of male truckers than females. Furthermore, the analysis shows that delays in clearance time leads to increase in both male and female HIV/AIDS infections which causes an increase in the number of AIDS cases in this border crossing points.

Keywords Truckers, HIV, AIDS, Next Generation Matrix

1. Introduction

Kenya has many border stations, however Busia and Malaba are most commonly used by heavy commercial vehicles. The Malaba border is the busiest crossing point for truckers using the Northern Corridor and connects landlocked countries of Uganda, DR Congo, South Sudan and parts of Rwanda to Mombasa port. Approximately 1,000 cargo trucks pass through the Malaba one stop border post daily while Busia handles an average of 600 trucks according to Bulzomi *et al.* [1]. According to the first volume of the final report of the Africa Trade Mark East Africa in 2014, the high volumes of traffic on the Northern Corridor have stretched these stations; causing long queues of trucks and delays in clearance of freight [2]. These delays make the truckers wait for long time for clearance while interacting with the local communities at these border towns. Malaba border post clears over 1,000 trucks daily which is a tall order compared to the high number of trucks using the route since 80% of cargo to Uganda, South Sudan, DR Congo, Rwanda and Burundi passes here [3]. Clearance backlog at

the Malaba border is threatening to increase costs and expose importers to higher storage charges. These delays are caused by frequent downtimes on Kenya Revenue Authority's (KRA) scanner at the border [4]. According to Maredi [5], truckers spend a few hours to several days at the border while the trucks are loaded with new goods or while customs processes are being cleared. Truckers spent this time in the company of commercial sex workers since they spent approximately less than 40 days a year at home with their families. East African countries have the highest HIV prevalence and high number of people living with HIV and AIDS [6]. Also the mobile populations which includes the truckers were key in the spread of COVID-19 and HIV pandemics. According to the NACC in charge of HIV policy in Kenya, border towns and counties in Kenya had the highest rates of new HIV infections [7].

Truckers are in the category of vulnerable populations to HIV and AIDS and hence can be a conduit of spread of the disease [8-10]. Results found that the HIV prevalence among the truckers in Kenya was 26-27%. Hence unprotected sexual interaction between the truckers and local community at Malaba and Busia would lead to spread of the pandemic within the locality. Also according to study by Nzyuko *et al.* [11] carried out at Malaba and Sachangwan indicated that

* Corresponding author:

ancentk@gmail.com (Ancent Makau Kimulu)

Received: Jan. 29, 2023; Accepted: Feb. 13, 2023; Published: Jun. 26, 2023

Published online at <http://journal.sapub.org/ajcam>

46% of girls reported usually having sex with truck drivers, 78% of girls reported usually exchanging sex for gifts or money and 59% of boys reported usually giving gifts or money for sex. This can lead to the spread of HIV from truck drivers and sex workers to adolescents in this stop overs along the Trans African highway in Kenya. According to [12], the tuckers along the Trans African highway are vulnerable to HIV and AIDS as a result of more time they spent away from their families. Other challenges on the highway includes poor road conditions, delays at the border crossing points which increases the time of travel between the designated destinations and lack of security in some sections of the highway especially as they drive at night. In Kenya, small towns offering truckers stops and rest bays like Mariakani, Voi, Mlolongo, Naivasha, Malaba and Busia, expose the truckers to environmental hazards such as HIV/ AIDS, drug and alcohol abuse. This could also lead to a similar exposure of the risk of HIV/ AIDS infection to the locals. According to Mulholland *et al.* [13], sexual networks include partnerships across resident, migrant, and mobile populations in the cross-border areas like Malaba and Busia. This implies that there is a lot of sexual activities at cross-border towns between truckers and the local community. Morris & Ferguson [14] found that there was an estimated population of 8000 female sex workers along the Northern corridor highway in Kenya. Furthermore, the female sex workers had 129 different sex partners and 634 sexual acts per year. The results also indicated that there were approximately 3200 – 4148 new HIV infections per year.

The above literature indicates that the truckers and their cohort can serve as conduits for the spread of HIV and AIDS along the corridor and to the local communities in the various stop-overs along the highway. This study uses the knowledge of mathematical modeling to study the effects of clearance time on HIV and AIDS dynamics of spread between the truckers and the local communities at Malaba and Busia border stops. The force of infection is considered

as a decaying exponential function while the population is largely classified into the binary compartment of male and female.

2. Methodology

The model flowchart is shown in figure 1. To formulate the model, only the adult populations are considered and some reasonable assumptions are placed. These assumptions include the following:

- i. Truckers involves both male and female drivers and their assistants.
- ii. Heterosexual intercourse is the main means of HIV transmission between truckers and local community.
- iii. There was heterogeneous mixing between the truckers and the local communities at Malaba and Busia.

By considering the flowchart (figure 1), the deterministic mathematical model governing the dynamics of the spread of HIV and AIDS among the truckers and the local communities at Malaba and Busia Border Stops is formulated below as;

$$\left. \begin{aligned} \frac{dS_m}{dt} &= r\Lambda - \tau\lambda_{fm}S_mI_f - \mu S_m \\ \frac{dS_f}{dt} &= (1-r)\Lambda - \tau\lambda_{mf}S_fI_m - \mu S_f \\ \frac{dI_m}{dt} &= \tau\lambda_{fm}S_mI_f - (\mu + \delta + \gamma_m)I_m \\ \frac{dI_f}{dt} &= \tau\lambda_{mf}S_fI_m - (\mu + \delta + \gamma_f)I_f \\ \frac{dA}{dt} &= \gamma_m I_m + \gamma_f I_f - (\mu + \delta)A \end{aligned} \right\} \quad (1)$$

where

$$\lambda_{fm} = \beta_{fm} e^{(-\rho_m)\rho_f}, \lambda_{mf} = \beta_{mf} e^{(-\rho_f)\rho_m},$$

and all parameters are between 0 and 1, i.e.,

$$0 < r, \Lambda, \lambda_{mf}, \lambda_{fm}, \beta_{mf}, \beta_{fm}, \rho_m, \rho_f, \tau, \gamma_m, \gamma_f, \mu, \delta < 1.$$




Figure 1. Model flowchart

3. Qualitative Analysis of the Model

Theorem 1: (Positivity and boundedness)

For time $t \geq 0$, the solutions of system (1) are confined in the set

$$\{S_m, S_f, I_m, I_f, A \in \mathbb{R}_+^5 : N = S_m + S_f + I_m + I_f + A\}.$$

Proof:

Let $\{S_m(t), S_f(t), I_m(t), I_f(t), A(t)\}$ be any solution set with positive initial conditions. The total population satisfies the equation

$$N(t) = S_m(t) + S_f(t) + I_m(t) + I_f(t) + A(t)$$

whose time derivative is given by,

$$\begin{aligned} \frac{dN}{dt} &= \frac{dS_m}{dt} + \frac{dS_f}{dt} + \frac{dI_m}{dt} + \frac{dI_f}{dt} + \frac{dA}{dt} \Rightarrow \frac{dN}{dt} \\ &= \Lambda - \mu N - \delta(I_m + I_f + A) \leq \Lambda - \mu N \end{aligned}$$

Using theory of differential equations (see [12]), then

$$N(t) \rightarrow \frac{\Lambda}{\mu} \text{ as } t \rightarrow \infty.$$

Hence, our solution to system (1) is biologically feasible in the region

$$\Psi = \left\{ (S_m, S_f, I_m, I_f, A) : S_m, S_f, I_m, I_f, A \in \mathbb{R}_+^5 \text{ and } N \leq \frac{\Lambda}{\mu} \right\}$$

$$\begin{aligned} S_m^* &= \frac{r\Lambda - (\mu + \delta + \gamma_m)I_m^*}{\mu}, S_f^* = \frac{(1-r)\Lambda}{\lambda_{mf}I_m^* - \mu}, I_f^* = \frac{\Lambda\tau\lambda_{mf}(1-r)I_m^*}{(\mu + \delta + \gamma_m)(\tau\lambda_{fm}I_m^* + \mu)}, \\ A^* &= \frac{\gamma_m I_m^* (\mu + \delta + \gamma_m) (\lambda_{mf} I_m^* - \mu) + \Lambda \tau \lambda_{mf} \gamma_f (1-r) I_m^*}{(\mu + \delta + \gamma_m) (\tau \lambda_{fm} I_m^* + \mu)}. \end{aligned}$$

3.2. Reproduction Number

Kaur *et al.* [15] and Kimulu *et al.* [16] defined the reproduction number R_0 as the average number of new HIV infections caused by a single infectious individual in a susceptible population in an entire infectious period. It is determined by the next generation matrix (NMG) method where \mathcal{F} and \mathcal{V} represents matrices of new infections and transitions out of the infected classes respectively, given by;

$$\mathcal{F} = \begin{pmatrix} \tau\lambda_{fm}S_mI_f \\ \tau\lambda_{mf}S_fI_m \\ 0 \end{pmatrix} \text{ and } \mathcal{V} = \begin{pmatrix} (\mu + \delta + \gamma_m)I_m \\ (\mu + \delta + \gamma_f)I_f \\ -\gamma_m I_m - \gamma_f I_f + (\mu + \delta)A \end{pmatrix}.$$

The Jacobian matrices of \mathcal{F} and \mathcal{V} at the DFE are denoted as \mathcal{F}_0 and \mathcal{V}_0 respectively and thus, we have;

$$\mathcal{F}_0 = \begin{pmatrix} 0 & \tau\lambda_{fm}S_m^0 & 0 \\ \tau\lambda_{mf}S_f^0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \quad (3)$$

$$\mathcal{V}_0 = \begin{pmatrix} \mu + \delta + \gamma_m & 0 & 0 \\ 0 & \mu + \delta + \gamma_f & 0 \\ -\gamma_m & -\gamma_f & \mu + \delta \end{pmatrix} \quad (4)$$

$$\mathcal{V}_0^{-1} = \begin{pmatrix} \frac{1}{\mu + \delta + \gamma_m} & 0 & 0 \\ 0 & \frac{1}{\mu + \delta + \gamma_f} & 0 \\ \frac{\gamma_m}{(\mu + \delta + \gamma_m)(\mu + \delta)} & \frac{\gamma_f}{(\mu + \delta + \gamma_f)(\mu + \delta)} & \frac{1}{\mu + \delta} \end{pmatrix} \quad (5)$$

and is bounded and positively invariant. ■

3.1. Equilibrium Points

The *disease-free equilibrium point* (DFE) is the point at which there is no HIV at Malaba and Busia border stop and its environs. It is obtained by setting $I_m = I_f = A = 0$ in system (1) and we obtain the DFE (E^0) as;

$$E^0 = (S_m^0, S_f^0, I_m^0, I_f^0, A^0) = \left(\frac{r\Lambda}{\mu}, \frac{(1-r)\Lambda}{\mu}, 0, 0, 0 \right).$$

The *Endemic Equilibrium Point* (EEP) is the point at which the HIV persists at Malaba border stop and it is spreading in the population. According to Kaur *et al.* [15], the EEP is obtained by solving the system (2) below

$$\left. \begin{aligned} r\Lambda - \tau\lambda_{fm}S_m^*I_f^* - \mu S_m^* &= 0 \\ (1-r)\Lambda - \tau\lambda_{mf}S_f^*I_m^* - \mu S_f^* &= 0 \\ \tau\lambda_{fm}S_m^*I_f^* - (\mu + \delta + \gamma_m)I_m^* &= 0 \\ \tau\lambda_{mf}S_f^*I_m^* - (\mu + \delta + \gamma_f)I_f^* &= 0 \\ \gamma_m I_m^* + \gamma_f I_f^* - (\mu + \delta)A^* &= 0 \end{aligned} \right\} \quad (2)$$

The EEP is obtained as

$$EEP = (S_m^*, S_f^*, I_m^*, I_f^*, A^*)$$

where

and thus

$$\mathcal{F}_0 \mathcal{V}_0^{-1} = \begin{pmatrix} 0 & \frac{\tau \lambda_{fm} S_m^0}{\mu + \delta + \gamma_f} & 0 \\ \frac{\tau \lambda_{mf} S_f^0}{\mu + \delta + \gamma_m} & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}.$$

The R_0 is obtained as the spectral radius of the next generation matrix FV^{-1} as

$$R_0 = \frac{\tau \Lambda}{\mu} \sqrt{\frac{\lambda_{fm} \lambda_{mf} r(1-r)}{(\mu + \delta + \gamma_m)(\mu + \delta + \gamma_f)}}.$$

3.3. Stability Analysis [17-20]

Theorem 2: (Local stability of Disease-Free Equilibrium). The DFE $E^0 = (S_m^0, S_f^0, I_m^0, I_f^0, A^0)$ is locally asymptotically stable when $R_0 < 1$ and is asymptotically unstable when $R_0 > 1$.

Proof:

The local stability of DFE was studied by determining the Jacobian matrix of system (1) at DFE. The eigenvalues were calculated and their nature was used to determine the stability of the DFE. The Jacobian matrix for system (1) at DFE is given by;

$$J^{DFE} = \begin{pmatrix} -\mu & 0 & 0 & -\tau \lambda_{fm} S_m^0 & 0 \\ 0 & -\mu & -\tau \lambda_{mf} S_f^0 & 0 & 0 \\ 0 & 0 & -(\mu + \delta + \gamma_m) & \tau \lambda_{fm} S_m^0 & 0 \\ 0 & 0 & \tau \lambda_{mf} S_f^0 & -(\mu + \delta + \gamma_f) & 0 \\ 0 & 0 & \gamma_m & \gamma_f & -(\mu + \delta) \end{pmatrix} \tag{6}$$

The eigenvalues are determined from;

$$|J^{DFE} - \lambda| = \begin{vmatrix} -\mu - \lambda & 0 & 0 & -\tau \lambda_{fm} S_m^0 & 0 \\ 0 & -\mu - \lambda & -\tau \lambda_{mf} S_f^0 & 0 & 0 \\ 0 & 0 & -(\mu + \delta + \gamma_m) - \lambda & \tau \lambda_{fm} S_m^0 & 0 \\ 0 & 0 & \tau \lambda_{mf} S_f^0 & -(\mu + \delta + \gamma_f) - \lambda & 0 \\ 0 & 0 & \gamma_m & \gamma_f & -(\mu + \delta) - \lambda \end{vmatrix} = 0 \tag{7}$$

Three of the eigenvalues are

$$\lambda_1 = \lambda_2 = -\mu, \lambda_3 = -\mu - \delta$$

and the remaining two eigenvalues are obtained by solving the quadratic equation;

$$\lambda^2 + ((\mu + \delta + \gamma_m) + (\mu + \delta + \gamma_f))\lambda + (\mu + \delta + \gamma_m)(\mu + \delta + \gamma_f) - \frac{\tau^2 \Lambda^2 \lambda_{fm} \lambda_{mf} r(1-r)}{\mu^2} = 0 \tag{8}$$

Using Routh-Hurwitz criterion for a polynomial of degree 2, then (8) has negative eigenvalues if;

(i) $a_1 = (\mu + \delta + \gamma_m) + (\mu + \delta + \gamma_f) > 0$

(ii) $a_2 = (\mu + \delta + \gamma_m)(\mu + \delta + \gamma_f) - \frac{\tau^2 \Lambda^2 \lambda_{fm} \lambda_{mf} r(1-r)}{\mu^2} > 0$

(i) is always true since all parameters are positive and (ii) implies $R_0 < 1$. Thus, the DFE is locally asymptotically stable if $R_0 < 1$. ■

To prove the global stability, the theorem by Castillo-Chavez *et al.* [21] is used.

Theorem 3: [21] The system (1) can be written in the form:

$$\left. \begin{aligned} X'(t) &= F(X, Y) \\ Y'(t) &= G(X, Y), G(X, 0) = 0 \end{aligned} \right\} \tag{9}$$

where $X = S = (S_m, S_f)$ is the number of uninfected individuals, $Y = (I_m, I_f, A)$ gives the HIV infectives and the AIDS patients. The DFE is denoted by $E^0 = (X_0, 0) = (\frac{r\Lambda}{\mu}, \frac{(1-r)\Lambda}{\mu}, 0, 0, 0)$. The conditions (H_1) and (H_2) stated below must be satisfied for global asymptotic stability to be guaranteed,

H₁: For $X'(t) = F(X_0, 0)$, X_0 is globally asymptotically stable,

H₂: $G(X, Y) = AY - \hat{G}(X, Y)$, $\hat{G}(X, Y) \geq 0$ and $A = D_Y G(X_0, 0)$ is an M-Matrix and in the region where model makes biological sense.

Theorem 4: (Global stability of the DFE). The DFE

$$E^0 = (X_0, 0) = \left(\frac{r\Lambda}{\mu}, \frac{(1-r)\Lambda}{\mu}, 0, 0, 0 \right)$$

is globally asymptotically stable if $R_0 < 1$ and assumptions in equation (9) are satisfied.

Proof:

Consider

$$F(X_0, 0) = \left(\frac{r\Lambda}{\mu}, \frac{(1-r)\Lambda}{\mu} \right), G(X, Y) = AY - \hat{G}(X, Y),$$

and

$$A = \begin{pmatrix} -(\mu + \delta + \gamma_m) & \tau\lambda_{fm}S_m^0 & 0 \\ \tau\lambda_{mf}S_f^0 & -(\mu + \delta + \gamma_f) & 0 \\ \gamma_m & \gamma_f & -(\mu + \delta) \end{pmatrix}, Y = \begin{pmatrix} I_m \\ I_f \\ A \end{pmatrix} \quad (10)$$

$$\hat{G}(X, Y) = \begin{pmatrix} \tau\lambda_{fm}S_mI_f - (\mu + \delta + \gamma_m)I_m \\ \tau\lambda_{mf}S_fI_m - (\mu + \delta + \gamma_f)I_f \\ \gamma_mI_m + \gamma_fI_f - (\mu + \delta)A \end{pmatrix}, AY = \begin{pmatrix} -(\mu + \delta + \gamma_m)I_m + \tau\lambda_{fm}S_m^0I_f \\ \tau\lambda_{mf}S_f^0I_m - (\mu + \delta + \gamma_f)I_f \\ \gamma_mI_m + \gamma_fI_f - (\mu + \delta)A \end{pmatrix} \quad (11)$$

Thus,

$$AY - \hat{G}(X, Y) = \begin{pmatrix} \tau\lambda_{fm}(S_m^0 - S_m)I_f \\ \tau\lambda_{mf}(S_f^0 - S_f)I_m \\ 0 \end{pmatrix} \geq 0 \text{ if}$$

hence $S_m^0 > S_m, S_f^0 > S_f \Rightarrow \hat{G}(X, Y) \geq 0$. A is an M-matrix therefore DFE is globally asymptotically stable when $R_0 < 1$. ■

Theorem 5: (Global stability of EEP). Let $E^* = (S_m^*, S_f^*, I_m^*, I_f^*, A^*)$ be the endemic equilibrium of the system (1) and V be a positive definite function on some neighborhood of E^* , then E^* is globally asymptotically stable if $\frac{dV}{dt} \leq 0$.

Proof:

Let the Lyapunov function be;

$$V = b_1 \left(S_m - S_m^* - S_m^* \ln \frac{S_m}{S_m^*} \right) + b_2 \left(S_f - S_f^* - S_f^* \ln \frac{S_f}{S_f^*} \right) + b_3 \left(I_m - I_m^* - I_m^* \ln \frac{I_m}{I_m^*} \right) + b_4 \left(I_f - I_f^* - I_f^* \ln \frac{I_f}{I_f^*} \right) \quad (12)$$

The time derivative of V is given by,

$$\frac{dV}{dt} = b_1 \left(1 - \frac{S_m^*}{S_m} \right) \frac{dS_m}{dt} + b_2 \left(1 - \frac{S_f^*}{S_f} \right) \frac{dS_f}{dt} + b_3 \left(1 - \frac{I_m^*}{I_m} \right) \frac{dI_m}{dt} + b_4 \left(1 - \frac{I_f^*}{I_f} \right) \frac{dI_f}{dt}. \quad (13)$$

Substituting $\frac{dS_m}{dt}, \frac{dS_f}{dt}, \frac{dI_m}{dt}$, and $\frac{dI_f}{dt}$ we obtain;

$$\begin{aligned} \frac{dV}{dt} &= b_1 \left(1 - \frac{S_m^*}{S_m} \right) (r\Lambda - \tau\lambda_{fm}S_mI_f - \mu S_m) + b_2 \left(1 - \frac{S_f^*}{S_f} \right) ((1-r)\Lambda - \tau\lambda_{mf}S_fI_m - \mu S_f) \\ &+ b_3 \left(1 - \frac{I_m^*}{I_m} \right) (\tau\lambda_{fm}S_mI_f - (\mu + \delta + \gamma_m)I_m) + b_4 \left(1 - \frac{I_f^*}{I_f} \right) (\tau\lambda_{mf}S_fI_m - (\mu + \delta + \gamma_f)I_f) \end{aligned} \quad (14)$$

From system (1), then at endemic equilibrium point the following relations are true;

$$\left. \begin{aligned} r\Lambda &= \tau\lambda_{fm}S_m^*I_f^* + \mu S_m^* \\ (1-r)\Lambda &= \tau\lambda_{mf}S_f^*I_m^* + \mu S_f^* \\ \mu + \delta + \gamma_m &= \frac{\tau\lambda_{fm}S_m^*I_f^*}{I_m^*} \\ \mu + \delta + \gamma_f &= \frac{\tau\lambda_{mf}S_f^*I_m^*}{I_f^*} \end{aligned} \right\} \quad (15)$$

Substituting (15) into (14) and simplifying gives;

$$\begin{aligned} \frac{dV}{dt} = & - \left(b_1 \left(\frac{S_m - S_m^*}{S_m} \right) \left(\tau \lambda_{fm} (S_m I_f - S_m^* I_f^*) + \mu (S_m - S_m^*) \right) + b_2 \left(\frac{S_f - S_f^*}{S_f} \right) \left(\tau \lambda_{mf} (S_f I_m - S_f^* I_m^*) + \mu (S_f - S_f^*) \right) \right) \\ & + \left(b_3 \left(\frac{I_m - I_m^*}{I_m} \right) \tau \lambda_{fm} \left(S_m I_f - \frac{S_m^* I_f^*}{I_m^*} \right) + b_4 \left(\frac{I_f - I_f^*}{I_f} \right) \tau \lambda_{mf} \left(S_f I_m - \frac{S_f^* I_m^*}{I_f^*} \right) \right) \end{aligned} \quad (16)$$

$E^* = (S_m^*, S_f^*, I_m^*, I_f^*, A^*)$ is globally asymptotically stable if $\frac{dV}{dt} < 0$. Using equation (16), then

(i) $\frac{dV}{dt} = 0$ if and only if $S_m = S_m^*, S_f = S_f^*, I_m = I_m^*$, and $I_f = I_f^*$,

(ii) $\frac{dV}{dt} < 0$ if and only if

$$\begin{aligned} & \left(b_3 \left(\frac{I_m - I_m^*}{I_m} \right) \tau \lambda_{fm} \left(S_m I_f - \frac{S_m^* I_f^*}{I_m^*} \right) + b_4 \left(\frac{I_f - I_f^*}{I_f} \right) \tau \lambda_{mf} \left(S_f I_m - \frac{S_f^* I_m^*}{I_f^*} \right) \right) < \\ & - \left(b_1 \left(\frac{S_m - S_m^*}{S_m} \right) \left(\tau \lambda_{fm} (S_m I_f - S_m^* I_f^*) + \mu (S_m - S_m^*) \right) + b_2 \left(\frac{S_f - S_f^*}{S_f} \right) \left(\tau \lambda_{mf} (S_f I_m - S_f^* I_m^*) + \mu (S_f - S_f^*) \right) \right). \blacksquare \end{aligned}$$

4. Numerical Simulation and Analysis

The system (1) which models the effect of clearance time on HIV and AIDS dynamics between truckers and the local community was simulated using MATLAB ODE45 solver [24-26]. The following default parameters were used for simulation.

Parameter	Value	Source	Parameter	Value	Source
r	0.069	[22]	ρ_f	0.056	Kenya AIDS Indicator Survey 2012
Λ	0.3	[16]	τ	4 -14 days	Assumed
λ_{mf}	0.246	Calculated	γ_m	0.89	Assumed
λ_{fm}	0.117	Calculated	γ_f	0.66	Assumed
β_{mf}	0.25	[14]	δ	0.016	[22]
β_{fm}	0.119	Assumed	μ	0.0539	[23]
ρ_m	0.32	[14]			

4.1. Analysis of Results

From the analysis of the results, figures (2-4) shows the effects of force of infection of females to males (λ_{fm}) with time on the HIV/AIDS dynamics between truckers and local community at Malaba and Busia border stops. Figure 2 shows that the susceptible male population increases with increase in λ_{fm} . This is attributed to the influx of truckers at the Malaba and Busia border stops due to delay in the clearance time. Also as λ_{fm} increases the number of HIV infections among the males increases as indicated in figure 3 This is as a result of higher number of susceptible population as indicated in figure 2 increase in the force of infection from females to males which implies more infections in the males. In figure 4, increases in λ_{fm} produces an increase in the number of AIDS cases initially and reduces exponentially with time due to increase in HIV infections as indicated in figure 3.

The effects of the clearance time on the HIV/AIDS dynamics at the Malaba and Busia Border stops are displayed in figures (5-9). Figure 5 shows that susceptible male population increase with the clearance time at the border stops. This can be attributed to the fact that the truckers population has more males than females and hence the males increases rapidly at the border stop overs with increase delays in clearance time. This consequently reduces the number females at the border stops interacting with the male truckers as displayed in Figure 6. In figure 7, increase in duration of clearance time increases the number of male HIV infections due higher number of susceptible male population. Figure 8 shows that the female HIV infections increases slightly as the clearance time increases. The increase in the number of infections consequently increases AIDS cases recorded approximately between $t = 0$ and $t = 5$ then decreases exponentially with time as shown in figure 9.




Figure 2. Variation of S_m with λ_{fm}




Figure 5. Variation of S_m with τ




Figure 3. Variation of I_m with λ_{fm}




Figure 6. Variation of S_f with τ




Figure 4. Variation of *AIDS* with λ_{fm}




Figure 7. Variation of I_m with τ




Figure 8. Variation of I_f with τ




Figure 9. Variation of AIDS with τ

5. Discussion of Results

The numerical simulations in figures (2-4) shows the effects of force of infection from females to males on the HIV/AIDS dynamics at Malaba and Busia border towns. In figure 2, as λ_{fm} increases, the susceptible male population increases. This is attributed to the increase in the susceptible population of males and females hence more sexual interactions at the border stop towns due to the delays in the clearance freight across the border. Increase in λ_{fm} increases the number of HIV infected males as displayed in figure 3. This is as a result of influx of truckers at the border stop towns as a result of delays in clearance across the border. These delays increases the number of interactions between truckers and the local communities increases the number males infected by the females. The increase in HIV male infections results to increase in the number of AIDS cases from figure 4. This is because more males will be progressing to AIDS class because AIDS is not curable.

Figures 5-9 displays the effects of clearance time on HIV/AIDS dynamics at Malaba and Busia border towns. From figure 5, the susceptible males increases with increase

in the length of the clearance time due to influx of truckers at the towns as a result of delays in clearance across the border. The susceptible females slightly decreases as τ increases as shown in figure 6 which can be attributed to the fact that more truckers are males leading to more female infections reducing the susceptible female population. Figures 7 and 8 shows that as τ increases HIV/AIDS infections for both male and female populations. This due longer period of sexual interactions between incoming truckers and the local communities producing more HIV/AIDS infections. The increase in HIV/AIDS infections due to delay in clearance time and hence more sexual interactions consequently leading to increase in AIDS cases at border crossing points.

6. Conclusions

This paper formulated a mathematical model to study the effects of clearance time on HIV/AIDS dynamics between truckers and local community at Malaba and Busia border stops. The model was numerically solved and analyzed using MATLAB ode45 solver. The analysis indicates that delays in clearance of trucks at Malaba and Busia increases the susceptible population due the presence of incoming truckers. This further increases sexual interactions between the truckers and the local community which increases HIV/AIDS infections and hence more AIDS cases at the border crossing points. The Kenyan government through the Kenya Revenue Authority department should collaborate with the Uganda and other countries whose use these points to use most recent technology to clear the cargo at the border points. This may involve clearing the cargoes even before arriving to reduce long queues at the border points. They should also collaboratively set up backup systems to use during KRA downtimes.

Definition of Parameters

τ	Proportion of male population recruited to the susceptible population
Λ	Rate of recruitment into susceptible population
λ_{mf}	Force of infection of males to females
λ_{fm}	Force of infection of females to males
β_{mf}	HIV transmission rate of males to females
β_{fm}	HIV transmission rate of females to males
ρ_m	HIV prevalence of males
ρ_f	HIV prevalence of females
τ	Clearance time across the border in days
γ_m	Rate at which infected males progress to AIDS
γ_f	Rate at which infected females progress to AIDS
δ	Disease induced death
μ	Natural death rate

REFERENCES

- [1] Bulzomi, A., Danssaert, P., Finardi, S., Matthysen, K., & International Peace Information Service (Antwerp). (2014). *Supply chains and transport corridors in East Africa*. International Peace Information Service.
- [2] Africa Trade Mark East Africa (2014). Final Report–1st Volume.
- [3] Ranganathan, R., & Foster, V. (2011). East Africa's infrastructure: a regional perspective. *World Bank Policy Research Working Paper*, (5844).
- [4] Kuria, S. W. (2019). Influence of automated systems on customs performance in Kenya.
- [5] Maredi, L. (2015). *Investigating the causes of delays at border posts: A focus on institutional and infrastructural factors at Beitbridge Border Post* (Master's thesis, University of Cape Town).
- [6] Malinga, T., Wiysonge, C. S., Ndwandwe, D., Okeibunor, J. C., & Talisuna, A. O. (2021). A scoping review of the impact of long-distance truck drivers on the spread of COVID-19 infection. *The Pan African Medical Journal*, 38.
- [7] Shannon, K., Strathdee, S. A., Goldenberg, S. M., Duff, P., Mwangi, P., Rusakova, M., ... & Boily, M. C. (2015). Global epidemiology of HIV among female sex workers: influence of structural determinants. *The Lancet*, 385(9962), 55-71.
- [8] Plummer, F. A., Nagelkerke, N. J., Moses, S., Ndinya-Achola, J. O., Bwayo, J., & Ngugi, E. (1991). The importance of core groups in the epidemiology and control of HIV-1 infection. *AIDS (London, England)*, 5 Suppl 1, S169-76.
- [9] Bwayo, J. (1994). Human Immunodeficiency Virus Infection in Long-Distance Truck Drivers in East Africa. *Archives of Internal Medicine*, 154(12), 1391. <https://doi.org/10.1001/archinte.1994.00420120123013>.
- [10] Mbugua, G. G., Muthami, L. N., Mutura, C. W., Oogo, S. A., Waiyaki, P. G., Lindan, C. P., & Hearst, N. (1995). Epidemiology of HIV infection among long distance truck drivers in Kenya. *East African Medical Journal*, 72(8), 515–518.
- [11] Nzyuko, S., Lurie, P., McFarland, W., Leyden, W., Nyamwaya, D., & Mandel, J. S. (1997). Adolescent sexual behavior along the Trans-Africa Highway in Kenya. *Aids*, 11, S21-S26.
- [12] Kimulu, A. M., Mutuku, W. N., Mwalili, S. M., Malonza, D., & Oke, A. S. (2022). Male Circumcision: A Means to Reduce HIV Transmission between Truckers and Female Sex Workers in Kenya. *Journal of Mathematical Analysis and Modeling*, 3(1), 50-59.
- [13] Mulholland, G. E., Markiewicz, M., Arimi, P., Ssengooba, F., Weir, S., & Edwards, J. K. (2022). HIV prevalence and the HIV treatment Cascade among female sex Workers in Cross-Border Areas in East Africa. *AIDS and Behavior*, 26(2), 556-568.
- [14] Morris, C. N., & Ferguson, A. G. (2006). Estimation of the sexual transmission of HIV in Kenya and Uganda on the trans-Africa highway: the continuing role for prevention in high risk groups. *Sexually transmitted infections*, 82(5), 368-371.
- [15] Kaur, N., Ghosh, M., & Bhatia, S. S. (2014). Mathematical analysis of the transmission dynamics of HIV/AIDS: Role of female sex workers. *Applied Mathematics & Information Sciences*, 8(5), 2491.
- [16] Kimulu, A. M., Mutuku, W. N., Mwalili, S. M., Malonza, D., and Oke, A. S. (2022). Mathematical Modelling of the Effects Funding on HIV Dynamics Among Truckers and Female Sex Workers Along the Kenyan Northern Corridor Highway. *Advances in Applied Sciences*, 7(3): 52–64.
- [17] Bada, O. I., Oke, A. S., Mutuku, W. N., and Aye, P. O. (2020). Analysis of the Dynamics of SI-SI-SEIR Avian Influenza A(H7N9) Epidemic Model with Re-infection. *Earthline Journal of Mathematical Sciences*, pages 43–73.
- [18] Juma, B. A., Oke, A. S., Mutuku, W. N., Ariwayo, A. G., and Ouru, O. J. (2022). Dynamics of Williamson fluid over an inclined surface subject to Coriolis and Lorentz forces. *Engineering and Applied Science Letter*, 5(1): 37–46.
- [19] Oke, A. S. and Bada, O. I. (2019). Analysis of the dynamics of avian influenza A(H7N9) epidemic model with re-infection. *Open Journal of Mathematical Sciences*, 3(1): 417–432.
- [20] Oke, A. S., Bada, O. I., Rasaq, G., and Adodo, V. (2021). Mathematical analysis of the dynamics of COVID-19 in Africa under the influence of asymptomatic cases and re-infection. *Mathematical Methods in the Applied Sciences*, 45(1): 137–149.
- [21] Castillo-Chavez, C., Feng, Z., & Huang, W. (2002). On the computation of ρ and its role on. *Mathematical approaches for emerging and reemerging infectious diseases: an introduction*, 1, 229.
- [22] UNAIDS (2021). *Global Factsheets 2020* <https://aidsinfo.unaids.org/> [Accessed on December 22, 2021]
- [23] AVERT HIV, and AIDS In Kenya (2021). *Kenya - Death Rate, Crude - 2021 Data 2022 Forecast 1960*. www.avert.org/professionals/hiv-around-world/sub-saharan-africa/kenya [Accessed on December 22, 2021]
- [24] Oke, A. (2017). Convergence of Differential Transform Method for Ordinary Differential Equations. *Journal of Advances in Mathematics and Computer Science*, 24(6): 1–17.
- [25] Oke, A. S., Fatunmbi, E. O., Animasaun, I. L., and Juma, B. A. (2022). Exploration of ternary hybrid nanofluid experiencing Coriolis and Lorentz forces: case of three-dimensional flow of water conveying carbon nanotubes, graphene, and alumina nanoparticles. *Waves in Random and Complex Media*, pages 1–20.
- [26] Okundalaye, O., Othman, W., and Oke, A. (2022). Toward an efficient approximate analytical solution for 4-compartment COVID-19 fractional mathematical model. *Journal of Computational and Applied Mathematics*, 416: 114506.