

**ASSESSMENT OF GERMINATION POTENTIAL AND POPULATION STATUS
OF *Melia volkensii* Gürke IN KITUI COUNTY, KENYA**

JANE MUKAMI NJEHU (BSc)

I56/31194/2015

**A THESIS SUBMITTED IN PARTIAL FULFILMENT OF THE
REQUIREMENTS FOR THE AWARD OF THE DEGREE OF MASTER
OF SCIENCE (PLANT ECOLOGY) IN THE SCHOOL OF PURE AND
APPLIED SCIENCES OF KENYATTA UNIVERSITY.**

MAY, 2022

DECLARATION

This thesis is my original work and has not been presented for a degree or other awards in any other University.

Signature..... Jane Mukami Njehu Date..... 31/5/2022

Jane Mukami Njehu – Reg No: I56/31194/2015

Department of Plant Sciences

SUPERVISORS

We confirm that the work reported in this thesis was carried out by the student under our supervision.

Signature..... Emily Wabuye Date..... 3/6/2022

Dr. Emily Wabuye

Department of Plant sciences

Kenyatta University

Signature..... Adelaide Mutune Date..... 3/6/2022

Dr. Adelaide Mutune

Department of Plant sciences

Kenyatta University

DEDICATION

I dedicate my work to the Almighty God for granting me good health, strength, wisdom and sufficient grace to pursue my studies. I am more grateful to my husband Stephen, mum Josephine and my three children Angel, Wisdom and Prudence for giving me moral support throughout my study period.

ACKNOWLEDGEMENTS

I wish to sincerely appreciate my two supervisors from Kenyatta University Department of Plant Sciences - Dr. Emily Wabuye and Dr. Adelaide Mutune. They have been very supportive in offering guidance and encouragement in my studies. Their patience and meticulous input gave me the motivation to soldier on and complete my studies. Much appreciation goes to Kenya Forestry Research Institute (KEFRI) under the management of Dr. Joshua Cheboiwo for assisting me financially in carrying out my research work. My sincere appreciation goes to Mr. Bernard Kamondo and Stephen Kiama, Research Scientists in KEFRI for their guidance in my data collection. Special thanks go to Mr. Daniel Buyela and Mr. Abuid Sayah who have been very instrumental in my data analysis.

I express my gratitude to the technical staff of KEFRI at Muguga and Kitui stations for assisting me in data collection. Mr. Mwandime of the National Museums of Kenya (NMK) is thanked for assisting in specimen identification and not to forget my colleagues at KEFRI especially the scientists for their encouragement throughout my studies. Finally, I thank Mr. Muema, a village elder from Ikutha Sub County in Kitui County providing local names of tree species and directing me in locating the study sites.

TABLE OF CONTENTS

DECLARATION	ii
DEDICATION	iii
ACKNOWLEDGEMENTS	iv
TABLE OF CONTENTS	v
LIST OF TABLES	viii
LIST OF FIGURES	ix
LIST OF PLATES	x
ABBREVIATIONS AND ACRONYMS	xi
ABSTRACT	xii
CHAPTER ONE	1
INTRODUCTION	1
1.1 Background information	1
1.2 Statement of the problem	5
1.3 Justification	6
1.4 Research Hypotheses	6
1.5 Objectives	7
1.5.1 General objective	7
1.5.2 Specific objectives	7
1.6 Significance of the Study	7
CHAPTER TWO	9
LITERATURE REVIEW	9
2.1 Description of <i>Melia volkensii</i>	9
2.2 Economic importance of <i>Melia volkensii</i>	10
2.4 Population status and natural regeneration of trees	12
2.4.1 Biotic and abiotic factors affecting natural regeneration	13
2.5 Plant species associations in nature and its relationship to the environment	14

2.6 Seed storage conditions as determinants of longevity	15
2.6.1 Seed moisture content	16
2.6.2 Seed storage temperature and relative humidity	16
2.6.3 Factors responsible for germination of <i>Melia volkensii</i>	17
CHAPTER THREE	18
MATERIALS AND METHODS	18
3.1 Study area.....	18
3.1.1 Population and livelihood activities in Kitui County	19
3.2 Research design.....	22
3.2.1 Establishment of sampling plots for study of population status of <i>M. volkensii</i> and associated species in Kasaala location	22
3.2.2 Sampling of <i>M. volkensii</i> and associated tree species	22
3.3 Germination and viability experiments	24
3.4 Data Analysis	29
3.4.1 Population status of <i>M. volkensii</i> and diversity of associated tree species	29
3.4.2 Formulae used for determining tree parameters were as follows:.....	30
3.4.3 Germination tests.....	31
3.5 NACOSTI research license.....	31
CHAPTER FOUR.....	32
RESULTS	32
4.1 Population status of <i>M. volkensii</i> in study area (Kasaala Location).....	32
4.1.1 Diversity and relative abundance of tree species in Kasaala Location.....	32
4.1.2. Mean number of <i>Melia volkensii</i> trees in the study area (Kasaala Location)	34
4.1.3: Height of <i>Melia volkensii</i> in Kasaala Location	35
4.1.4 Mean DBH (diameter at breast height) of <i>M. volkensii</i> in Kasaala Location ...	36
4.1.5 DBH size-class distribution in Kasaala Location	37
4.1.6 Natural regeneration of <i>M. volkensii</i> in Kasaala Location.....	38
4.1.7 Diversity indices of tree species in Kasaala Location	38
4.2.1 Initial germination percentage of <i>M. volkensii</i> nuts and seeds.....	39

4.2.2 Germination percentage of <i>M. volkensii</i> nuts and seeds after storage	40
CHAPTER FIVE.....	44
DISCUSSION, CONCLUSIONS AND RECOMMENDATIONS.....	44
5.1 Discussion	44
5.1.1 Status and Population structure of <i>Melia volkensii</i> in the Study Area	44
5.1.2 Effect of seed extraction and storage conditions on viability of <i>M. Volkensii</i>	46
5.2 Conclusions	50
5.3 Recommendations	50
5.3.1. Recommendations from the study	50
5.3.2 Areas for further research.....	51
REFERENCES.....	52
APPENDICES	66
Appendix 1: Composition of tree species in transect 1	66
Appendix 2: Composition of tree species in transect 2	67
Appendix 3: Composition of tree species in transect 3	68
Appendix 4: Composition of tree species in transect 4	69
Appendix 5: Composition of tree species in transect 5	70
Appendix 6: Composition of tree species in transect 6	71
Appendix 7: Composition of tree species in transect 7	72
Appendix 8: DBH size-class distribution of <i>Melia volkensii</i> in the study area (Kasaala Location).....	73
Appendix 9: Diversity of <i>M. volkensii</i> and associated tree species in Kasaala Location	74
Appendix 10: NACOSTI Research License	75

LIST OF TABLES

Table 4.1: Diversity of <i>M. volkensii</i> and associated tree species in Kasaala Location ...	32
Table 4.2: Mean number of <i>Melia volkensii</i> trees in Kasaala Location.	34
Table 4.3 Mean number of coppices and seedlings in Kasaala Location.....	38
Table 4.4: Diversity indices of tree species in the study area (Kasaala Location)	39
Table 4.5: Initial Germination percentage of <i>M. volkensii</i> nuts and seeds.....	40
Table 4.6: Germination means of <i>M. volkensii</i> during storage.....	43

LIST OF FIGURES

Figure 3.1: Map showing Tiva Forest within Kasaala Location in Kitui County.....	21
Figure 3.2: Diagram (not drawn to scale) showing line transects established in Kasaala Location.....	24
Figure 3.3: Flow diagram showing storage of <i>Melia volkensii</i> seeds and nuts	27
Figure 4.1: Relative abundance of <i>M. volkensii</i> and other tree species in Kasaala Location.....	35
Figure 4.2: Mean height of <i>M. volkensii</i> in Kasaala Location	36
Figure 4.3: Mean DBH (cm) of <i>M. volkensii</i> in Kasaala Location	36
Figure 4.4: DBH size-class distributions of <i>Melia volkensii</i> in Kasaala Location	37
Figure 4.5: Mean germination percentage of <i>M. volkensii</i> under cold temperature (-20°C) over a period of 2-6 months.....	40
Figure 4.6: Mean germination percentage of <i>M. volkensii</i> in room temperature (30°C) over a period of 2-6 months	41

LIST OF PLATES

Plate 2.1: <i>Melia volkensii</i> seedlings	10
Plate 3.1: Dried nuts and extracted seeds of <i>Melia volkensii</i>	25
Plate 3.2: Collection of <i>Melia volkensii</i> fruits from Tiva forest	28
Plate 3.3: Seed extraction and commencement of <i>M. volkensii</i> germination tests	28
Plate 3.4: <i>Melia volkensii</i> fruits.....	29

ABBREVIATIONS AND ACRONYMS

ANOVA	Analysis of variance
ASALs	Arid and Semi-Arid Lands
DBH	Diameter at breast height
GPS	Global Positioning System
ICRAF	International Centre for Research in Agroforestry
ISTA	International Seed Testing Association
JICA	Japan International Cooperation Agency
KEFRI	Kenya Forestry Research Institute
MC	Moisture Content
NACOSTI	National Commission for Science, Technology and Innovation
NMK	National Museums of Kenya

ABSTRACT

Melia volkensii Gürke is a native tree species that grows in the arid and semi-arid areas of East Africa. There has been overutilization of this tree which may lead to risk of rapid decrease in its community and genetic diversity. Additionally, planning and implementing planting programmes and conservation plans of *M. volkensii* have been hindered by insufficient knowledge on its seed storage behavior. This study sought to develop optimal protocols for seed germination and assess population status of *Melia volkensii* to enhance conservation and supply chain management in Kitui County. *Melia volkensii* fruits were collected from Tiva Forest and the germination experiment was carried out in KEFRI Kitui within nursery conditions. Seed germination experiments were arranged in a completely randomized design with three factors – extraction state, storage container and storage temperature. For extraction, seeds were divided into two lots – extracted seeds and unextracted seed (nut). Two types of storage containers were used namely sealed and unsealed. The storage temperatures were room temperature (30°C) and cold storage (-20°C). Germination tests were done before storage, and later after 2, 4 and 6 months in storage from each of the storage conditions. A hundred seeds and nuts each were sown for each experimental unit. The rate of seedling sprouting was recorded for a duration of 10 days as a percentage of seeds sown. Variation in germination percentages was tested using two-way analysis of variance (ANOVA) at 95 % confidence interval and means separated using fisher's protected least significant difference in R and SAS statistical software. The highest percentage (89 %) was scored in nuts kept in unsealed containers at 30°C at 6 months and the lowest (14 %) in seeds stored in similar conditions for 2 months. The differences were significant. These results indicate that the germination rates of *M. volkensii* seeds increased with increase in period of storage. Additionally, plant population was determined *in situ* across seven transect lines measuring 1050 meters in length. Sampling was carried out in five plots measuring 50 x 50 m established at intervals of 200 meters. Transect 1 and 7 were laid in farmland while the transects between these two were laid in bushland. All tree species represented in each plot were counted. Height and diameter at breast height (DBH) were measured for all *M. volkensii* trees within the plots. *Melia volkensii* seedlings and coppices were counted in 5 x 5 m nested plots within the larger plots. One-way analysis of variance (ANOVA) at 95 % confidence interval was used to test for variance in parameters measured and means separated using Tukey's HSD. There were significant differences in height and DBH means of *M. volkensii* among transects. Transect 2 had the lowest mean height and DBH while the tallest trees were in transects 1, 3, 4, and 6. Transect 1 had highest mean DBH. Means of seedlings were not significantly different among transects. Coppicing was significantly highest in transects 1 and 4. Moreover, relative abundance, species richness and diversity of plant species observed were analyzed using Simpson's index of biodiversity and Shannon-Wiener species diversity index. *Melia volkensii* was the most frequently occurring species with 12.2% (n=67) followed by *Acacia tortilis* 10.0 % (n=55). *Calotropis strophela*, *Commiphora capensis*, *Maerua crassifolia* were the least represented relative abundance of 0.18 %. Values of indices used varied among transects and did not follow the location of transect implying that diversity was not determined by land disturbance. In conclusion, nuts stored unsealed at room temperature have the highest longevity. Based on diversity and relative abundance of *M. volkensii* and associated tree species in the study area, tree exploitation in Kasaala Location does not cause depletion of *M. volkensii*.

CHAPTER ONE

INTRODUCTION

1.1 Background information

The genus *Melia* belongs to plant family *Meliaceae* (mahogany family) and has less than nine species, most of which are native to Australia and Asia. *Melia volkensis* (Gurke) is the only tree species that is native to East Africa (Mabberley, 1984). *Melia volkensis* is mostly found in regions that lie at an altitude of between 400 and 1600 meters on dry and wooded grasslands. Distribution of the species extends from the dry areas of Somalia to Ethiopia through Tanzania and Kenya (Orwa *et al.*, 2009). This species is known to shed leaves during the dry season and is 6-20 m tall with a diameter of 25 cm at maturity. It is open crowned and laxly branched.

The wood is durable, decay and termite resistant; mostly used in paneling, making frames and rafters. The wood of *M. volkensis* is comparable to Elgon teak or camphor (Maundu and Tengas, 2005) which makes the tree to be over-exploited and thus a great need to replenish it for conservation. *Melia volkensis* does well in soils that drain water properly, for instance the sandy loam. *Melia volkensis* has proved economically and environmentally viable as it cushions against climate change and provides profitable timber. Lateral branches develop from the tree which are usually pruned and the tree develops a bole that is straight. The bark is grey in color and the tree has bright green compound leaves which are 35 cm in length; bearing less than ten deeply lobed leaflets. The flowers are white and small sized while the fruit is a drupe about 4 cm long. The

fruit goes through the process of depulping in order to obtain nuts/ hard kernels which enclose the brown black seeds (Kamondo *et al.*, 2016).

The economic importance of *M. volkensii* are diverse, for instance it is used for timber, fodder, herbal medicine fuel and making pesticides. Overexploitation of the products of this species has led to depletion of this tree species especially for medicinal use. For instance, dried fruit is used as ruminant feed supplement, leaf extract is applied externally on boils and blisters and it also used for treatment of stomach upsets, eczema, skin rashes and relieving body ache (Joshi, 2007). In the recent past, *M. volkensii* is known as an agroforestry tree which has other desirable properties including coppicing ability (ICRAF, 1992). Researchers, farmers, agricultural extension and education officers have found it to be a tree of high economic value, though it has difficulties in seed germination (Kimondo and Kiamba, 2005). Current dissemination programmes on *M. volkensii* have to utilize plants propagated from stem cuttings and roots, rather than seedlings (Mulatya *et al.*, 2000).

Additionally, seed germination is determined partly by storage conditions (Kamondo *et al.*, 2016). Unsuitable storage methods regularly result in low germination of seeds and total failure in germination Nasreen, (2000). Domestication of plants is an important aspect which has broadened the understanding of seed storage. Naturally, seeds are categorized in three different ways namely: orthodox, recalcitrant and intermediate seeds (Hong and Ellis, 1996). Orthodox seeds are as well-known as the drying-tolerant seeds. Seed storage behavior of *M. volkensii* is termed as orthodox, and viability is enhanced for many years in airtight containers at a temperature of -20°C. Over a wide

variety of storage conditions and temperatures, the lifetime of these materials rises as the amount of moisture and temperature in the environment decreases. This leads to delayed ageing of seeds thus prolonged viability (Liu, *et al.*, 2020). Recalcitrant seeds do not withstand drying well, and as a result, they are not well suited for prolonged storage, even though the necessary level of moisture differs from species to species. Even while intermediate seeds are much more susceptible to drying out and dehydration than recalcitrant seeds, they are far less tolerant of desiccation than orthodox seeds, and they often lose viability much more quickly at low temperatures (Tweddle *et al.*, 2003).

It has been recommended that plant species with poor regeneration and consist of low importance value indices be regarded as more important than others in matters of conservation (Maua *et al.*, 2020). Natural regeneration of *M. volkensii* enables the plant to survive in an area after a number of events eventually causing change in the composition and the structure of a woodland ecosystem. According to (Nieuwenhuis and Egan 2002), natural regeneration can be defined as establishment of a new forest from the seeds that have been sown, coppice shoots, or root suckers. Coppices arise sporadically on the stump of a tree that was cut previously. Dry land habitats consist of woody plants which have regenerated and continue to exist through alarming events like wind, flooding, fire, storms and anthropogenic activities. When leaves of trees fall and decompose, they produce a favorable habitat for seeds to sprout, consequently increasing the number of seedlings (Midgley and William, 1996).

In semi-arid eco-systems, the flora formation is diverse with plant species interrelating with each other depending on environmental conditions. These conditions include: light,

temperature and moisture. Analysis of plant associations increases the knowledge of seed dispersal, environmental heterogeneity and biotic interactions (Woods *et al.*, 2019). Plant species are known to associate randomly when interactions between species are weak. *Melia volkensii* has been reported to occur in Acacia-Commiphora vegetation in ASALs and it is commonly referred to be a natural stands' remnant (Muok, *et al.*, 2010) and (Alados *et al.*, 2006). There are positive and negative relationships between the species some of which are common phenomena in plant communities. Relationship between plants explains how interacting plants safeguard each other from stressful conditions for instance; shade of trees and extreme temperature conditions. Plant association among species may be caused by biological relationships or by the varied reactions of species to abiotic stimuli, among other things. Additionally, plant association affects the possible connections among species implying that species can interact only where they co-occur (Johnson, 2018). Nevertheless, there might be elevated associations among species, which means that the apparent lack of other species can have an effect on the link between certain species in certain circumstances (Beth *et al.*, 2020; Pickett, 1980).

The study of plant population status and association may contribute to development of comprehensive and effective conservation strategies of *M. volkensii* in the arid and semi-arid lands (ASALS) of East Africa. It is therefore significant to comprehend the population status of *M. volkensii* so as to inform if there is high erosion of their population and genetic diversity (Runo *et al.*, 2004). Additionally, the study on seed storage of this tree species will inform the best storage conditions for longevity.

Generally proper seed storage improves the survival of plants and the seed have good physiological and physical conditions from the harvesting time until planting time. The study results are anticipated to help stakeholders and policy makers to come up with strategies for optimization of germination and natural regeneration of *M. volkensii*.

1.2 Statement of the problem

Over the years, there has been over-exploitation of *M. volkensii* populations leading to depletion of the genetic diversity of this tree species. Despite its significance as a species that is of high value in the dry areas, *M. volkensii* has proved difficult to propagate adequately for extensive production and dissemination. Effort to boost growing of this tree species have not been spared by propagation challenges (Kyalo, 2004). These include seed dormancy, difficulties in extraction of seeds and inability of *Melia volkensii* seeds to survive the seedling stage. Additionally, there is inadequate knowledge on the appropriate seed storage methods which leads to limited supply of quality seeds.

Moreover, mass production of seedlings has not been achieved due to challenges in the establishment of *M. volkensii* plantations in the arid and semi-arid lands of East Africa. Effort to raise *M. volkensii* through various methods like tissue culture and stem cuttings have been carried out by researchers with little success (Malinda *et al.*, 2013). Anthropogenic activities such as clear-cut logging, forest fires, grazing and conversion of forests into farmlands have however greatly affected the populations of *M. volkensii* in the drylands of Kenya. Besides the anthropogenic influence it is not clear how plant species combinations and their biological interactions influence population status and natural regeneration of *M. volkensii* in Kasaala location of Kitui County.

1.3 Justification

The act of storing and supplying *M. volkensii* seeds is presently founded on dealing with nuts rather than seeds that have been processed/extracted. Seeds of *M. volkensii* kept as nuts take up much space making charges of transportation and storage high. One kilogram of *M. volkensii* nuts yields about 200 seeds, it is interesting to note that 4,000 to 4,500 seeds per kilogram of extracted seeds is equivalent to one kilogram of *M. volkensii*. Effective handling and storage of processed/extracted *M. volkensii* seeds is rarely practiced since there is limited information on the best storage methods. However, factors responsible for low germination of *Melia volkensii* after storage are not properly comprehended. Findings from this research will advise on the acceptable practice in handling and storage of *M. volkensii* seeds. Additionally, this study will avail data on post and pre-storage extraction which increases the germination rate of *M. volkensii* and will develop improved protocols to boost propagation of *M. volkensii* (Kamondo *et al.*, 2016). According to Nturibi (2010), *M. volkensii* despite its popularity has challenges in germination due to unreliable propagation protocol and poor access of quality seeds. Moreover, optimization of seed storage conditions and assessment of *M. volkensii* population status will help in improving the conservation strategies of *M. volkensii* in the dry lands of Kenya. Findings on appropriate storage of seeds of this tree would assist seed suppliers on coming up with a stock of seeds to guarantee supply of seeds at every time even in times when fruiting is irregular.

1.4 Research Hypotheses

- i. The risk of tree depletion has no significant effects on the population status of *Melia volkensii* in Kasaala Location of Kitui County.

- ii. Storage conditions of seeds and nuts of *Melia volkensii* have no significant effects on germination.

1.5 Objectives

1.5.1 General objective

To develop optimal protocols for seed germination and assessment of population status of *Melia volkensii* to enhance conservation and supply chain management in Kitui County.

1.5.2 Specific objectives

- i. To determine the effect of storage conditions on germination of *M. volkensii* seeds stored as extracted seeds and dried nuts.
- ii. To evaluate the population status of *M. volkensii* and associated species under natural conditions in Kasaala Location in Kitui County.

1.6 Significance of the Study

Rise in population, land cultivation, economics and urbanization are the conceivable determinants of deforestation in the countries in East Africa like Kenya, Uganda, Tanzania and Ethiopia (Lemma *et al.*, 2021). Seed quality plays an important role in the production of agricultural crops. Comprehending behavior of seed storage is indispensable in planning and conducting programmes of planting which enhance sustainable utilization of plant species. The study will also inform the government organizations and non-governmental organizations (NGOs) to change from using poor methods of seed storage and instead focus on how to assist private organizations to afford best seed storage devices. As a result, there will be affordable quantities and

qualities of *M. volkensii* seeds; this will eventually raise the economic revenue for people living in the dry lands of Kenya. Moreover, research on evenness of distribution and the population status of *M. volkensii* will inform on conservation of genes and the environmental diversification of *M. volkensii*. Improved propagation of *M. volkensii* for Kenya's ASALs will place Kenya a notch higher in its efforts to increase its forest cover now standing at 7 % compared to the recommended 10 % (FAO and UNEP, 2020). Findings from this research will help improve the growth of Melia leading to the improvement of propagation and preservation of this tree species. Overall, the study will be a huge milestone in attaining environmental sustainability.

CHAPTER TWO

LITERATURE REVIEW

2.1 Description of *Melia volkensii*

Melia volkensii, also commonly known as Melia or Mukau (Kamba), belongs to the family Meliaceae. The tree has been reported to have straight and clean boles approximately 4 meters high without whorls and has no forking below 4 meters in height (Esilaba *et al.*, 2011). Trees that belong to Meliaceae family have male and female flowers or only one type of both and are either dioecious or monoecious. In rare occasions, some species of *M. volkensii* have functional hermaphrodite flowers and may be accompanied by male flowers (Styles, 1972). The fruit of *Melia volkensii* is a pulp that contains approximately 10 % crude fat and over 12 % crude protein. The mature leaves contain more than 5 % crude fat and 21 % crude protein (Muok *et al.*, 2010). Melia fruits are depulped to produce nuts which are later cracked for seed extraction. Brown black seeds are found within the hard kernels and they are propagated in a seedbed filled with sterilized river sand. Seedlings of *M. volkensii* develop a carrot-like root which is full of oils and nutrients that allow them to tolerate dry circumstances (Kamondo *et al.*, 2016).



Plate 2.1: *Melia volkensii* seedlings

2.2 Economic importance of *Melia volkensii*

Over the years *M. volkensii* has been identified as a highly valued commercial tree in the dry lands of Kenya. It is popular in the dry lands because it is able to tolerate drought. It produces high quality timber used for construction, household furniture, fuel wood and apiculture. The tree is used to make bee hives because the timber can be shaped without difficulty and its flowers produce high quality nectar which is converted into honey (Maundu and Tegas, 2005; Muok *et al.* 2010). The wood is long-lasting and could be utilized for interior paneling, as well as for floor tiles, rafters, and frame construction (Luvanda *et al.*, 2015). In addition, its fruits and leaves are used as fodder for goats and

cattle. The flowers provide excellent bee forage and the tree also has medicinal properties (Orwa *et al.*, 2009). Additionally, *Melia volkensii* is used for mulching, providing shade and acts as a wind breaker (Tedd, 1997). It is evident that leave extracts of *M. volkensii* are used by some people to control ticks and fleas on goats (Jaoko *et al.*, 2020). The tree is also known in herbal preparations where it is being used as an insecticide (Wilps *et al.*, 1993). *Melia volkensii* is used in litter accumulation whereby fallen leaves help to improve the soil fertility consequently, increasing the crop yields (Maundu and Tenge, 2005).

2.3 Distribution and ecology of *Melia volkensii*

The growth of *Melia volkensii* is restricted to the dry lands of Kenya, Tanzania, Ethiopia and Somalia (Kimondo and Kiamba, 2005). The tree is commonly found in Machakos, Mwingi, Kitui and Makueni. The tree is also found growing in the bushland and wooded grasslands in Counties like Embu, Isiolo, Taita- Taveta, Samburu and Marsabit (Crossland, *et al.*, 2021). The species is normally found growing in relation with Acacia-Commiphora vegetation (Mutiga, 2021). It emanates in deciduous bushland, sometimes bordering seasonal canals or appearing on rock outcrops (Maundu and Bo, 2005). *Melia volkensii* trees grow well in sandy, soils but prefers sandy-loamy soils with good drainage. The tree can be found growing at altitudes varying from 350 to 1700 meters above sea level, in regions with average yearly rainfall ranging from 300 to 800 millimeters. *Melia volkensii* does not tolerate black cotton soils or areas prone to water logging (Muok *et al.*, 2010).

2.4 Population status and natural regeneration of trees

Comprehension of plant diversity and natural status of woody species is vital in preserving the forests and restoration of degraded areas (Ghanbari *et al.*, 2021). Moreover, natural regeneration can be relied upon to affect the forest stands' dynamics as well as to preserve the genetic features of local populations of trees. It is popular in large scale forest restoration in African regions (Sugimoto *et al.*, 2011). The processes that bring about naturally regenerating forests promote genetic adaptation and sustain local biodiversity and biological interactions. Most of the indigenous tree species undergo natural regeneration, for instance *M. volkensis*, *Vitellaria paradoxa*, *Vitex keniensis*, *Parkia biglobosa*, *Tilia americana* and *Adansonia digitata* (Ky-Dembele, *et al.*, 2010). *Melia volkensis* sprouts from stumps as well as growing from seeds. In areas where the tree occurs coppicing is also a common mode of regeneration. Coppicing occurs in trees that have been cut from the base, leading to a strong vegetative response and the regeneration of new shoots (Wang *et al.*, 2008). Coppice sprouts are largely formed by latent buds that emerge from the trunk of a tree after it has been cut down or otherwise removed. It is evident that trees can grow from stumps which mostly produce a number of stems. The ability to re-sprouting is a common mechanism of *M. volkensis* rejuvenation in dry forests and it should be deliberated as a method of forest rehabilitation and improvement of agroforestry. (Kruger *et al.*, 1997) Therefore, understanding how abiotic and biotic factors influence natural regeneration is significant for studies of ecology (Khaine, 2018).

2.4.1 Biotic and abiotic factors affecting natural regeneration

Natural regeneration of plant communities is interrelated with the differences in conditions of environment. Saplings that grow in old-growth tropical and subtropical forests are more vulnerable to biological and non-biological factors than mature trees that have grown for a long time (Queenborough *et al.*, 2009). Majority of the tree species might have specific light, nutrient and topography requirements for regeneration (Jin *et al.*, 2018) although regeneration of species is not very sensitive to the available resources (Dechnik-Vázquez *et al.*, 2016). When there are biotic factors, like the number of adults and seedlings that are of the same species or different species, they can also affect how many trees grow from seed. Tree species go through niche splitting in conjunction with certain abiotic variables in order to facilitate the co-existence and preservation of a diverse range of species. The abiotic factors include water, light, topography and soil nutrients (Chesson *et al.*, 2000).

Barriers that deter plants' regeneration such as steep slopes of the land, inadequate source of seeds, residual trees and poor drainage which lead to mortality of the seedlings that are not able to compete vigorously. Areas with insufficient source of seeds could not be replenished normally and mountainous areas with extremely erodible soils, might not be appropriate to planting or fertilization. The productivity of forest cover can significantly reduce due to accelerated erosion processes. For instance, the organic top soil is subject to displacement through erosion following disturbances found in areas with steep slope (Ross and Dykes, 1996). In some cases, natural regeneration is adequate to re-establish forest cover however, in other cases, seeding directly or seedlings' planting would be required if re-forestation is occurring in a given time (Yang *et al.*, 2019).

2.5 Plant species associations in nature and its relationship to the environment

Plant species association can be defined as the relationship of various species found in an environment. It is also described as a motionless organic link created by the interrelationship of species. However, niche of species regulates how species relate with the environment surrounding them; by having different niches species are able to coexist (Albrecht *et al.*, 2001). Species diversity is defined as the number and relative abundance of species found in a given biological organization (Whittaker *et al.*, 2001).

In plant communities, positive and negative relationships amongst species are very common phenomenon. The concept of plant associations informs on the status of conditions of environment like light, temperatures as well as humidity (Dale, 1999). The relationship among species appears from biological associations or diverse reaction of species to abiotic factors and on the other hand, it affects the possible association among species. This implies that species can interact only where they coexist in the same environment (Kershaw and Looney, 1985). Species association in a virgin forest provides sufficient information on how to manage forests. For instance, *M. volkensii* and *Cassuarina equisetifolia* improved fertility of soil through augmented nitrogen and phosphorous which was credited to fixation of nitrogen by *C. equisetifolia* through Frankia bacteria and nutrient recycling by *M. volkensii* (Mwadalu *et al.*, 2021). Additionally, species relationship between dominant species of trees like saplings and mature trees was examined in a National Nature Reserve in China. The findings indicated that species relationship showed a remarkable net positive relationship,

showing a compelling equilibrium of a steady formation and species composition in old-growth forest (Ziwei *et al.*, 2019).

The growth rate of native forests and other crops decelerates as intraspecific competition becomes higher making it a negatively density dependent process. As a result, plant characteristics linked to resource use end up being significant determinants of aggressive relations to community of plants (Duffy, 2006). Previous study on factors affecting multipurpose trees and shrubs utilization in ASALS in Kenya showed that there was an important association amongst the frequencies of woody tree species and the inclusive ultra violet radiation index (Kisangau, 2021).

2.6 Seed storage conditions as determinants of longevity

Seed germination is defined as hydration of the seed leading to emergence of the embryonic radicle and shoot (Bewley *et al.*, 2013). Maintaining seed viability over extended periods in storage is necessary for preserving the genetic integrity of plants. Successful seed storage conditions help in preserving the genetic integrity of *Melia volkensii* for long periods. Seed longevity in storage exemplified by initial viability, freedom from fungal and insect attack, seed maturity, physiological deterioration, and mechanical damage is determined by factors like moisture content of seeds, nature of seeds, temperature and relative humidity (Rajjou *et al.*, 2008). On the basis of storage behavior, three kinds of seed are recognized which are orthodox, recalcitrant and intermediate seeds (Berjak and Pammenter, 1997). Recalcitrant seeds are sensitive to chilling, are highly hydrated and they cannot withstand intensive desiccation (Bustam,

et al., 2021). *Melia volkensii* seed behavior is orthodox thus can retain its viability for several years under appropriate storage conditions (Orwa *et al.*, 2009).

2.6.1 Seed moisture content

Seed moisture is the most crucial component in sustaining germination rate during storage, because it is the principal factor influencing all cellular processes (Bonner *et al.*, 2008). The metabolism rates of both orthodox and recalcitrant seeds could be decreased by maintaining the seedlings in a dry condition at cold temperatures. True orthodox seeds maintained at low moisture content of between 5-10 % can be stored comfortably at nearly any temperature compared to recalcitrant seeds. Therefore, seed moisture content at the time of storage determines how low the temperature can be set for seed storage. Orthodox seeds can be exposed to sub-zero temperatures, while recalcitrant seeds cannot. Orthodox seeds can be stored for a few years at temperatures as low as -20°C (Ellis *et al.*, 1992) and (Ellis and Hong, 2006).

2.6.2 Seed storage temperature and relative humidity

The ideal storage temperature for seeds is approximately 14°C with a relative humidity of slightly less than 40 %. Most refrigerators have high relative humidity and hold a temperature of approximately 14.2°C (Bharat *et al.*, 2012). Seeds stored in a refrigerator should be kept in containers that are tightly closed to keep the humidity levels low. Orthodox seeds will sustain survivability better when hardened to low content of moisture of roughly 4–8 percent and then preserved in an environment in which humidity is regulated, than when preserved in equilibrium with ambient air humidity (Zhang *et al.*, 2021). Cool condition is mostly favorable for storage of orthodox seeds. Long-term conservation of gene resources of orthodox agricultural seeds is at a

temperature of less than -20°C and a moisture content of approximately -6°C to 16°C . Previous study shows that half-life is doubled for seeds dried to equilibrium and thereafter to approximately -20°C in waterproof containers under long-term gene bank (Ellis, 1996).

2.6.3 Factors responsible for germination of *Melia volkensii*

The longevity of orthodox seeds, also referred to as drying-tolerant seeds is thus determined by content of moisture (MC) and temperature. Previous research indicates that factors responsible for poor germination of *Melia volkensii* during storage have been attributed to both physiological and physical factors (Milimo and Hellum, 1989). It is documented that farmers have used various traditional technologies to break seed dormancy to enhance propagation of *Melia volkensii*. These methods include: scarification procedure which is done by burning of the nuts. Nuts that are regurgitated by goats are usually used by farmers to raise seedlings (Mwamburi, 2005). Additionally, there are other methods that are used by farmers for instance splitting or cracking of the nuts and sowing of them in troughs (Kidundo 1997).

CHAPTER THREE

MATERIALS AND METHODS

3.1 Study area

Field study was undertaken in two sites of Kitui County, namely Kasaala and Tiva forests. Kasaala is a Location in Ikutha Sub-County of Kitui South constituency about 440 km from Kenya's capital city Nairobi, and approximately 21 km from Tsavo East National Park (Figure 3.1). Kasaala is known to host *Melia volkensii* trees that naturally occur unlike in Tiva forest where the trees are planted. Natural regeneration of *M. volkensii* is likely to occur in Kasaala because most land area is undisturbed. Tiva forest is 194 km East of Nairobi along Kitui-Machakos road (Figure 3.1). The experiment on germination of *M. volkensii* seeds was established in the nursery beds of KEFRI Kitui, 15 km away from Tiva Forest.


The soils of the study area consist of metamorphic parent material and the dominant soil groups are alfisols, ultisols, oxisols, and lithic soils. These soils are highly erodible and are of low fertility. The ultisols and alfisols are also susceptible to capping, which increases runoff and makes the clay soils hard to cultivate by the end of the dry season (Barber *et al.*, 1981). Approximately, less than 20 % of Kitui County has well-drained, deep, friable red and brown clays of good fertility. More than 60 % of the region has relatively shallow, sticky, red, black, and brown clays of variable fertility (Jaetzold and Schmidt, 1983). The study area, Kitui County, shares a similar geographical and vegetation structure with the Tsavo East National park. This environment favors the growth of the naturally growing *M. volkensii* being adapted to the dry weather conditions (Gachathi, 1994).

Kitui County is among the semi-arid Counties in Kenya that is faced with serious water scarcity. The numerous droughts encountered have resulted to a decline in water supplies which has likely contributed to the drying up of most of the seasonal rivers. The County is mostly dry and hot with temperatures ranging 14°C during the coldest months (July-August) and 34°C between January and March which are known to be the hotter months. The County receives between 500 mm and 1050 mm of rainfall annually, with an average rainfall of 900 mm a year (Gladys, 2017). The study area has an elevation between 400 m to 1830 m above sea level (Jaetzold, 2006) and the most widespread vegetation type in Kitui, is a thicket and bushland, especially *Acacia-Commiphora* associations (Ndegwa *et al.*, 2016).




3.1.1 Population and livelihood activities in Kitui County

Agriculture contributes more than 80 % of the income generated by the remote population and directly hires more than 35 % of the population in Kitui County. More than 80 % of the population is engaged in farming activities especially in the production of crops such as cereals, horticultural and industrial crops (Yohannis, 2019). The total annual average of cereal production in the County is 80,680 metric tonnes (Mt), valued at KSh 4.24 billion and industrial crops (sunflowers, cotton and sisal are valued at KSh 29.04 million for an average production of 771 Mt. The average production of horticultural crops is approximately 36,950 Mt; at a cost of nine hundred and ninety million (Mbiti and Misango, 2021). Rearing livestock is also a major economic activity in Kitui County (Kivunzya, 2019). Households relies on livestock for food security and for economic production. The major components of livestock in Kitui County include indigenous goats, cows, fowl, and donkeys. The farmers prefer these animals because they are resilient to the hot and dry

climatic conditions that occur during drought. Of the total farming households, more than 40 % rear indigenous cattle and approximately 70 % raise chicken and goats (Kivunzya, *et al.*, 2018). More than 40 % of households raise donkeys which are used as a means of transport especially when fetching water across long distances during dry seasons. Donkeys are sometimes included in dowry negotiations (Mbiti and Misango 2021). Annual production of honey in the County is 960 tonnes with 14 % of households engaged in the enterprise (Warui *et al.*, 2018; Musinguzi, *et al.*, 2018).



Legend

-  Roads
-  Tiva forest
-  Tiva forest

Scale 1:500,000

Figure 3.1: Map showing Tiva Forest within Kasaala Location in Kitui County

3.2 Research design

3.2.1 Establishment of sampling plots for study of population status of *M. volkensii* and associated species in Kasaala location

Plant sampling was done using a stratified design; this is a sampling method which encompasses the population division into minor sub-groups called strata. The entire population of trees was distributed into two different sites namely bushland and farmland. A bushland can be defined as an area covered with shrubs and trees or any other natural vegetation while a farmland is a land used for farming (Sisay *et al.*, 2021). The bushland in the study area consisted of a secondary vegetation of Acacia- Commiphora while the farmland was used for production of sorghum, cowpeas, maize and pigeon peas which were intercropped with *M. volkensii*. Broad vegetation formations of the study area consist of natural and manmade features. The natural features include seasonal rivers and valleys, while the man-made features include dams, urban settlements and roads. Seven transects were established to cover the whole forest using guidelines by Plumptre (1994). Transects 1 and 7 were laid on the disturbed sites where farming, grazing and cutting down of trees was rampant in search of timber and fuel. Transects 2,3,4,5 and 6 were laid on the bushland where natural regeneration was taking place (figure 3.2). Along each transect, 5 plots were established by using a geographic coordinate system to locate latitude and longitude of the study area. The number of trees, coppices and seedlings that were found along the length of the line and inside the plots were counted and recorded in data sheets.

3.2.2 Sampling of *M. volkensii* and associated tree species

Fifty by fifty-meter plots were established across transects measuring 1050 meters. Five plots were put in every transect at a range of 200 meters space between each plot (Manning

et al., 2020). All the trees of *M. volkensii* and associated tree species were sampled within the plots. Diameter at breast height (DBH) and height of *M. volkensii* trees were determined in each plot. A diameter tape was used to determine DBH at 1.5 meters from the ground. Suunto clinometer was adopted in measuring the tree height from the base of the tree and a linear tape was used to measure 20 meters. A nail was fixed on the tree's bark where the linear tape was hooked and placed horizontally from the eye to the tree. The linear tape and the clinometer were held by both hands and the bottom measurement of the clinometer was subtracted from the top reading. Additionally, coppices and mature trees were sampled within a fifty by fifty meters plot, while survival counts of seedlings was done within a 5 by 5 sub-plot nested within and at the corners of the 50 by 50 m plot.

Voucher specimens of *M. volkensii* and associated tree species were collected and identified in the field using identification guides (reference books, plant identification manuals and identification keys to plants). The voucher specimens were prepared by pressing the fresh materials on herbarium press boards and later drying the specimens using a plant drier (Miller, 1999). They were mounted on labelled herbarium sheets and deposited in the Kenya Forestry Research Institute local herbarium at Muguga. The specimens that could not be easily identified in the field were taken to the herbarium of the National Museums of Kenya for identification.




Figure 3.2: Diagram (not drawn to scale) showing line transects established in Kasaala Location

3.3 Germination and viability experiments

Mature ripe fruits of *M. volkensii* were collected from Tiva forest and depulped. *Melia volkensii* fruits were depulped by removing the outer fleshy part using a plank of wood to while placing them on stone. After depulping the fruits, the nuts obtained were dried and

were put on a groove carved out on a wooden plank and cracked open using a knife and hammer (Kamondo *et al.*, 2016). The germination materials were divided into two lots the first lot being *M. volkensii* nuts and the second lot extracted seeds. The nuts were cleaned in water and sun-dried for 5 days in preparation for seed extraction. The dry seed sub lots were also sun-dried for three days to a moisture content (MC) of 6 % that was determined by oven drying at 103°C for 24 hours (Ellias *et al.*, 1990). The moisture content was calculated by subtracting the dry weight of seeds from the initial/wet weight of seeds multiplied by one hundred (100).



Plate 3.1: Dried nuts and extracted seeds of *Melia volkensii*

Four hundred dry seeds were used for initial germination in 4 replicates of 100 seeds each as per International Seed Testing Association germinationn protocol (ISTA, 2012). During

test for germination, seeds were compressed and decontaminated in a solution of 5 grams of a fungicide Ridomil Gold fungicide (Metalaxyl-M and S-isomer) in 1 litre of distilled water. The seeds were later soaked in cold water (4°C) for 12 hours and finally rinsed with 1 % sodium hypochlorite solution. The coat of seeds was split longitudinally utilizing a razor blade that was decontaminated. Seeds were germinated in plastic containers using fine sterilized clean river sand as the substrate (Dolor, 2009). The seeds were covered with a layer of sand drenched with fungicide (Muok *et al.*, 2010). The number of germinated seedlings was daily recorded for a period of 10 days. Once scored, the sprouted seedlings were retrieved using a pair of tweezers to confirm the daily counts were for newly propagules.

In order to test the storage conditions' effect on germination of *M. volkensii*, seeds and nuts were stored under different conditions and tested for germination at an interval of 2 months after storage. The experimental conditions were room temperature (30°C) and in refrigerator at -20°C; under each temperature condition seeds and nuts were stored in sealed or plastic containers that are uncapped. The germination materials were tested for their germination at a 2-month interval for the next 6 months (Bharat, 2012; Probert, 2007). Each germination test consisted 4 replicates each with 25 seeds or nuts. Thus, the experimental design was three factorials arranged in a completely randomized design (figure 3.3). Germination was recorded for a period of 10 days.




Figure 3.3: Flow diagram showing storage of *Melia volkensii* seeds and nuts



Plate 3.2: Collection of *Melia volkensii* fruits from Tiva forest



Plate 3.3: Seed extraction and commencement of *M. volkensii* germination tests



Plate 3.4: *Melia volkensii* fruits

3.4 Data Analysis

Data collected were entered into Microsoft Excel spreadsheets (2019) for subsequent statistical procedures. Statistical Analysis Software (SAS), version 9.4 was used to analyze germination, while R software version 2.1.3 was used to analyze the population status of *M. volkensii*.

3.4.1 Population status of *M. volkensii* and diversity of associated tree species

One-way analysis of variance (ANOVA) at 5 % significance level was used to test whether there were any statistically significant differences between the means of diameter at breast height (DBH) and height of the trees in the sampled plots. The statistical test was also used

to determine if the means of mature trees, coppices and seedlings were different. Tukey honestly significant difference (HSD test) was utilised to separate means for all the parameters. Data obtained from counting trees in Kasaala forest were used to derive total density of plant species, species richness, relative abundance, and diversity of plants in the study area. Relative species abundance was used to compute the occurrence of each species in comparison with the other species recorded. Shannon wiener diversity index was used to estimate the tree species richness and evenness of individual distribution in the tree species. Additionally, Simpson's diversity index was instrumental in estimating diversity of trees and estimating the probability of two individual trees that belonged to different species (Spellerberg, 2003).

3.4.2 Formulae used for determining tree parameters were as follows:

- i) **Simpson's diversity index** was computed as follows:

$$D = \frac{\sum_{i=1}^R (ni(ni-1))}{N(N-1)}$$

$$N(N-1)$$

Where N = the total number of organisms of a particular species, while n = the total number of organisms of all species while R symbolizes richness which is a measure of how many different types the dataset of interest contains.

- ii) **Shannon-Wiener** was computed as follows:

$$H' = \frac{N \ln N - \sum (ni \ln ni)}{N}$$

- Where ni is the number of individuals in species i , and N is the total number of species in the sample.
- Species richness = number of species per specified number of individuals.

Since the datasets were small, the number of coppices, seedlings and mature trees were counted manually.

c) Species density = number of species per unit area. Species density per m² is usually used to measure species richness.

d) Relative species abundance = Number of plant species from one group

Total number of species from all groups

e) Plant diversity = (ni/N) . N is total number of individuals in S species, while ni is the number of individuals in the ith species.

3.4.3 Germination tests

Germination data for stored nuts and extracted seeds were compared using the t test at 5 % level of significance using a two-way ANOVA to analyze the effects of seed storage treatments and seed storage time. The treatments used were seeds stored either as extracted seeds or nuts at a temperature of 30° C and -20°C in closed and open containers. In this study seed storage time was 2, 4 and 6 months. At P <0.05, germination means were separated using Fisher's least significant difference test. At confidence interval of 95 %, the germination trend of extracted seeds and nuts that were stored at different environmental conditions and in different containers was tested using a t test.

3.5 NACOSTI research license

The license is an ethical approval to conduct research on optimization of germination and assessment of population status of *Melia volkensii* Gürke under natural conditions in Kitui County, Kenya for a period ending: 20/April/2022 (Appendix 10).

CHAPTER FOUR

RESULTS

4.1 Population status of *M. volkensii* in study area (Kasaala Location)

4.1.1 Diversity and relative abundance of tree species in Kasaala Location

In this study, a total of 20 families were represented in the sampled plots. These included Fabaceae (9 species), Euphorbiaceae (3 species), Bursereaceae (5 species) and Tiliaceae (6 species). There were 67 plants of which *M. volkensii* represented 12.2 % relative abundance and *Acacia tortilis* accounted for 10.0 %. *Adansonia digitata* represented a relative abundance of 9.5 % while *Commiphora baluensis* accounted for 8.5 % (Table 4.1).

Table 4.1: Diversity of *M. volkensii* and associated tree species in Kasaala Location

Family name	Species name	Number of trees	Relative abundance
Acanthaceae	<i>Duosperma cooperi</i>	4	0.73
Aizoaceae	<i>Delosperma cooperi</i>	2	0.36
Anacardiaceae	<i>Lannea schimperi</i>	15	2.73
Asclepiadaceae	<i>Calotropis strophela</i>	1	0.18
Bursereaceae	<i>Commiphora africana</i>	11	2.00
	<i>Commiphora baluensis</i>	8	8.46
	<i>Commiphora capensis</i>	1	0.18
	<i>Commiphora edulis</i>	17	3.10
	<i>Commiphora eminii</i>	7	1.28
Sterculiaceae/Malvaceae	<i>Sterculia rhynchorcarpa</i>	14	2.55
Capparaceae	<i>Boscia angustifolia</i>	12	2.19
	<i>Maerua crassifolia</i>	1	0.18
	<i>Maerua decumbens</i>	7	1.28
	<i>Maerua holstii</i>	3	0.55

	<i>Thylachium thomasii</i>	2	0.36
Rubiaceae	<i>Tennantia sennii</i>	7	1.28
Combretaceae	<i>Combretum aculeatum</i>	17	3.10
	<i>Terminalia brownii</i>	2	0.36
	<i>Terminalia prunioides</i>	7	1.28
Euphorbiaceae	<i>Euphorbia ingens</i>	1	0.18
	<i>Euphorbia tirucalli</i>	1	0.18
	<i>Euphorbia scheffleri</i>	1	0.18
Fabaceae	<i>Albizia anthelmintica</i>	13	2.37
	<i>Bauhinia coriacea</i>	1	0.18
	<i>Bauhinia natalensis</i>	15	2.73
	<i>Cassia abbreviata</i>	17	3.10
	<i>Dalbergia melanoxyton</i>	1	0.18
	<i>Delonix alata</i>	15	2.73
	<i>Erythrina melanacantha</i>	8	1.46
	<i>Tamarindus indica</i>	8	1.46
	<i>Tephrosia villosa</i>	2	0.36
Malvaceae	<i>Adansonia digitata</i>	52	9.47
Tiliceae	<i>Grewia africana</i>	2	0.36
	<i>Grewia bicolor</i>	2	0.36
	<i>Grewia calymmatosepala</i>	12	2.19
	<i>Grewia flava</i>	4	0.73
	<i>Grewia tephrodermis</i>	32	5.28
	<i>Grewia villosa</i>	18	3.28
Meliaceae	<i>Melia volkensii</i>	67	12.20
Mimosaceae	<i>Acacia busciae</i>	3	0.55
	<i>Acacia mellifera</i>	7	1.28
	<i>Acacia nilotica</i>	29	5.28
	<i>Acacia senegal</i>	15	2.73
	<i>Acacia tortilis</i>	55	10.02

Portulacaceae	<i>Calyptrorhiza taitensis</i>	4	0.73
Rhamnaceae	<i>Berchemia discolor</i>	9	1.64
Solanaceae	<i>Solanum campylacanthum</i>	4	0.73
	<i>Solanum taitensis</i>	2	0.36
Zygophyllaceae/ Balanitaceae	<i>Balanites aegyptiaca</i>	12	2.19
Phyllanthaceae	<i>Bridelia taitensis</i>	2	0.36
Grand total		552	100 %

4.1.2. Mean number of *Melia volkensii* trees in the study area (Kasaala Location)

There were significant differences in the number of *M. volkensii* trees that were found in the transects ($P < 0.05$) [table 4.2]. The highest number of *M. volkensii* trees were recorded in transect 2 while the lowest values were obtained in transects 1, 5 and 6 ($P < 0.05$). The differences were not however consistent with where transects were laid meaning that disturbance of land did not influence the number of trees regenerating.

Table 4.2: Mean number of *Melia volkensii* trees in Kasaala Location

Transect number	Mean number of <i>M. volkensii</i> trees
1.	1.20±0.84 ^b
2.	4.80±1.40 ^a
3.	2.40±1.12 ^{ab}
4.	2.20±0.86 ^{ab}
5.	0.60±0.25 ^b
6.	0.80±0.37 ^b
7.	1.40±0.51 ^{ab}
P value	< 0.05

The number of *M. volkensii* trees observed varied among transects (Figure 4.1). *Melia volkensii* trees found in the bushland had the highest relative abundance (24%, 12 % and

11 %) in transects 2, 3 and 4 respectively. It was least represented in transects 5 and 6 at 3 % and 4 % respectively. The trees that were found on the farmland had a moderate representation of relative abundance (6 % and 7 %) in transects 1 and 7 respectively.




Figure 4.1: Relative abundance of *M. volkensii* and other tree species in Kasaala Location

4.1.3: Height of *Melia volkensii* in Kasaala Location

The mean height of *M. volkensii* trees were highest in transects 1, 3, 4 and 6 and lowest in transect 2. The differences were significant at ($P < 0.05$) as shown in Figure 4.2.




Figure 4.2: Mean height of *M. volkensis* in Kasaala Location

4.1.4 Mean DBH (diameter at breast height) of *M. volkensis* in Kasaala Location

At $P < 0.05$, the DBH of trees within transects showed significant differences. The mean DBH of *M. volkensis* trees in transect 1 (40.8 cm) was significantly highest compared to the other transects at $p \leq 0.05$. Transects 3, 4, 5, 6 and 7 had mean DBH of between 24.0 to 31.4 cm, while transect 2 had the lowest mean DBH of 19.2 cm (figure 4.3).




Figure 4.3: Mean DBH (cm) of *M. volkensis* in Kasaala Location

4.1.5 DBH size-class distribution in Kasaala Location

Distribution of diameter at breast height of *M. volkensii* trees was computed for the all transects combined. The highest DBH size class was in the cluster ranging between 20.5 and 30 cm and the lowest was in the cluster that ranging between 70.5 and 80 cm with the latter having only 1 tree. The size class distribution ranging between 40.5-50 cm and between 50.5- 60.0 cm had the second lowest representation of 3 trees each (Figure 4.4).




Figure 4.4: DBH size-class distributions of *Melia volkensii* in Kasaala Location

4.1.6 Natural regeneration of *M. volkensii* in Kasaala Location

Coppicing and natural seed germination of *M. volkensii* were recorded in the study area. The mean number of coppices found in the study area were significantly different between transect 1 and 2 at $P < 0.05$. At $P > 0.05$ there were no significant differences in the mean number of seedlings that were found in Kasaala Location (Table 4.3).

Table 4.3 Mean number of coppices and seedlings in Kasaala Location

Transect Number	Mean number of coppices	Mean number of seedlings
1	0.60±0.4 ^b	0.40±0.4
2	9.20±1.3 ^a	2.40±0.9
3	1.20±0.5 ^b	1.80±0.2
4	8.20±1.0 ^a	1.00±0.6
5	2.20±0.9 ^b	1.60±0.5
6	1.00±0.5 ^b	1.40±0.7
7	1.40±0.7 ^b	0.60±0.6
P value	<0.05	> 0.05

4.1.7 Diversity indices of tree species in Kasaala Location

The highest Shannon diversity index of 2.23 was recorded in transect 7. The lowest Shannon diversity index was recorded in transect 3 (1.76). Shannon Wiener values as a measure of species evenness indicates that there was greater diversity in transect 7 compared to transect 3. Other (transects 1,2,4,5 and 6) had Shannon diversity indices ranging from 1.98 to 2.2. The highest Simpson's index was recorded in transect 7 (0.74),

while the lowest Simpson's index was recorded in transect 3 (0.69), other transects (transects 1,2,4,5 and 6) had Simpson's diversity indices ranging from 0.69 to 0.73 (Table 4.4). Simpson's index calculates the species evenness and richness as a measure of diversity. In this study thus transect 7 recorded the highest diversity and transect 3 the lowest. Diversity was therefore not determined by disturbance of the study site since transects 7 and 1 were established in farmland while transect 3 was in bushland.

Table 4.4: Diversity indices of tree species in Kasaala Location

Transects	Shannon-Wiener Index (H')	Simpson's Index (λ)
1	2.03	0.70
2	2.02	0.72
3	1.76	0.69
4	1.98	0.71
5	2.13	0.71
6	2.20	0.73
7	2.23	0.74

4.2: Effect of storage conditions on germination of *M. volkensii* seeds

4.2.1 Initial germination percentage of *M. volkensii* nuts and seeds

Initial germination test was done before storage of nuts and seeds. Nuts and seeds were dried to a moisture content of 6 %. At $p \leq 0.05$, there was no significant difference in initial germination percentage between dried extracted seeds and dried nuts (Table 4.5).

Table 4.5: Initial germination percentage of *M. volkensii* nuts and seeds

Seed type	Germination %
Nuts	57.7
Seeds	30.5
P value	>0.05

4.2.2 Germination percentage of *M. volkensii* nuts and seeds after storage

Nuts and extracted seeds were stored in different conditions (30°C and -20°C) and in closed and open containers for 6 months and germination tests done at 2, 4 and 6 months interval. There was a gradual increase in germination percentage over the 6 months experimental period (figure 4.5 and 4.6).




Figure 4.5: Mean germination percentage of *M. volkensii* under cold temperature (-20°C) over a period of 2-6 months.




Figure 4.6: Mean germination percentage of *M. volkensis* in room temperature (30°C) over a period of 2-6 months

At two months storage, seeds stored in open containers in cold room (-20°C) had the highest mean germination percentage of 60.58 whilst seeds in open room showed the least mean germination percentage of 14.00. The mean germination percentage under the different storage conditions after two months were in the following decreasing order: seed open cold room > nuts closed room > nuts open room > seeds closed cold room > seed closed room > nuts open cold room > nuts closed cold room > seed open room.

For the two-month period, significant differences at ($P < 0.05$) were generally observed between seeds stored in open containers under cold room (-20°C) and nuts stored in closed containers under room temperature (30°C). This was also observed between nuts in closed containers under room temperature and nuts stored in open containers under room temperature (30°C). There was also significant difference between nuts stored in closed containers under cold room and seed open room. There was no significant difference (P

>0.05) between nuts stored in open containers in ambient temperature and seeds kept in closed containers under cold room. Seeds stored in closed containers in room temperature and nuts stored in open containers under cold room had the intermediate germination means (table 4.6).

The highest mean germination percentage after a four-month period of storage was for seed that were stored in open containers in cold room (71.5 %) and the least mean germination percentage was for seed that was stored in open containers in room temperature (32 %). The mean germination percentage after four months were in the following decreasing order: seed open cold room > nuts closed room > seeds closed cold room > nuts closed room > seed closed room > nuts open room > nuts closed cold room > nuts open cold room > seed open room. At $P < 0.05$, there were significant differences between seed kept in unsealed containers at ambient temperature and nuts stored in room temperature in closed containers. The significant differences were also noted in nuts stored in room temperature in closed containers and those stored in open containers in room temperature (table 4.6).

The highest mean germination percentage after a six-month period of storage was recorded in nuts stored at room temperature (30°C) in open containers (89 %), while the lowest was in seeds stored in closed containers at room temperatures (58.5 %). Nuts stored in cold room (-20°C) in closed containers and those stored in open containers at room temperatures (30°C) had the intermediate mean germination percentages. At $P > 0.05$, there was no significant difference between seeds that were stored in closed containers in

room temperature (30°C) and nuts that were kept in unsealed containers in room temperature (table 4.6).

Table 4.6: Germination means of *M. volkensii* during storage

Storage conditions			Months in storage		
Extraction	Temperature	Container	2	4	6
Seeds	Room (30°C)	Open	14.0±1.37 ^f	32.0±2.7 ^e	58.5±4.1 ^{bc}
		Closed	49.0±0.58 ^{bc}	56.5±0.36 ^{bc}	65.1±1.8 ^c
	Cold ((20°C)	Open	60.58±2.4 ^a	71.5±1.9 ^a	79.8±3.7 ^{ab}
		Closed	43±9.6 ^d	61.0±2.3 ^b	74.5±3.1 ^{ab}
Nuts	Room (30°C)	Open	41.7±2.6 ^d	50.0±3.1 ^{cd}	89.0±3.2 ^a
		Closed	51.8±3.2 ^b	60.0±2.7 ^b	78.6±3.5 ^{ab}
	Cold (20°C)	Open	45.1±3.3 ^{bcd}	45±1.0 ^d	74.4±5.1 ^{ab}
		Closed	34.0±1.8 ^e	49.0±2.3 ^{cd}	73.1±5.2 ^{ab}
P value			<0.05	<0.05	<0.05

CHAPTER FIVE

DISCUSSION, CONCLUSIONS AND RECOMMENDATIONS

5.1 Discussion

5.1.1 Status and Population structure of *Melia volkensii* in the Study Area

In this study, *M. volkensii* was the most abundant species in the study area, followed by *Acacia tortilis*. These results are related to findings of Anjarwalla *et al.* (2016) who observed that *M. volkensii* is found growing alongside the Acacia-Commiphora vegetation, along uncultivated land which is covered with natural vegetation. The study also agrees with Kamondo *et al.* (2016), who reported that the tree species are drought tolerant and adaptable to the semi-arid weather conditions in Kitui County. Additionally, there were significant differences in the mean number of *M. volkensii* trees found in the study area, though the differences were not however consistent with where transects were established. It was expected that transects laid in the farmland would register lower mean number of *M. volkensii* trees than those in the bushland which wasn't the case. This can be attributed to the fact that the ecological patterns of a plant population are characterized by its population dynamics, as well as the correlation between each individual plant and the external environments (Odum, 1971).

In the study area, there existed vast farmlands with tall well-groomed cultivated *Melia volkensii* trees which were found growing on both the farmlands and the natural bushland vegetation. The bushland possessed a high H^1 index of tree species and supported a range of vegetation cover unlike the farmland that exhibited a lower H^1 index of tree species and a limited range of vegetation cover. This is probably because of anthropogenic activities

that were carried out in the farmland for instance cultivation, logging, overgrazing, and other habitat degradation activities thus causing impact on the trees and undercover biological species. These results are related to the observation of Caviedes and Ibarra (2017) who stated that environmental change caused by human beings have the ability to degenerate and simplify forest stands by accelerating the loss of trees, coppices and seedlings. These results also agree with the study carried out by Tsechoe (2014) that species richness increases at a lower elevation but decreases at a higher elevation. He also stated that the slope and aspect affect plant diversity in many ways, consequently affecting the formation of soils from parent rock. This is due to the soil physicochemical properties and microbial activities that are usually abundant at the topsoil leading to an abundance of tree species on flat bushland topography.

According to Borda-Nino *et al.* (2021) fluctuations in the sloppiness influence the vegetation composition and structure, leading to a lower diversity. *Melia volkensii* trees had high DBH and height indicator of the high rate of trees' growth, which were probably controlled by combined effects of biotic and abiotic factors, particularly soil microorganisms, soil nutrients, water, weather patterns, sunlight, and competition among trees (Stephenson, *et al.*, 2014). These results are also in agreement with Pang *et al.* (2017) who observed that the total tree height averages for assessment that was carried out on indicator species of a natural forest was between 0.2 -0.5 meters.

From this study, it was evident that, the steep slope farmlands carried a few species adapted to a narrow range of prevailing conditions. This can be attributed to the fact that soil erosion was prevalent on the farmland, leading to soil structure loss, nutrient

degradation, and soil compaction. Similar results were confirmed by Yang Liu *et al.* (2019), who observed that the topographical features were essential for soil formation. As well, they affect the soil condition and bacterial communities. Congruently, the steep soils may be eroded and thus lose their topsoil during the formation process. Therefore, they become thinner than the more nearly level grounds that receive deposits from areas upslope Flores *et al.* (2020).

A considerable portion of the bushland had a big number of mature trees, coppices, and seedlings of *Melia volkensii*. Thus, the positive association between mature individuals and juveniles of this tree species in the bushland suggests that *M.volkensii* juveniles are successfully establishing under the canopy of other associating trees that were found growing in the study area. This shows the role of trees in generating vegetation cover that plays a vital role in the maintenance of plant diversity in the study area. These results are congruent with Xu *et al.* (2020) who found evidence of positive association between tree over story-layer composition and the understory layer diversity in different forest types.

5.1.2 Effect of seed extraction and storage conditions on viability of *M. Volkensii*

The extracted seeds and those enclosed in nuts and stored for a period of two, four and six months significantly improved under all storage conditions from the initial germination of 30.5% to 89%. *Melia volkensii* seeds stored in open containers under cold storage recorded a higher mean germination percentage as compared to seeds stored in closed containers under cold storage. These results are akin to findings according to a *Melia volkensii* growing guide in Kenya's dryland regions (2018). From the study, it is evident that storing extracted seeds under air-tight conditions damages the seeds, leading to a

lower germinability. These results are related to Kamondo *et al.* (2016) who observed that extracted seeds are vulnerable to external conditions and lose viability in one month's period. As observed, *M. volkensii* seeds that were held in the form of nuts (un-extracted) were highly viable compared to those that were stored in form of seeds. This trend was ascribed to the stony layer covering *M. volkensii* seeds, enabling them to preserve the right moisture content, light and temperature for a longer time.

The nuts stored for a period of 2, 4 and 6 months in open containers under room temperature had a higher mean germination percentage as compared to seeds stored under similar storage conditions. These results are in agreement with the findings of Jaoko *et al.* (2020) who reported that *M. volkensii* extracted seeds possess a high oil content, which comes from health delicate unsaturated fats. Upon thermal shock, sunlight, and air, the seeds degrade, turning rancid. In reaction, the seeds lose germination and consequently have a limited shelf life that results to a reduced germination rate. In my study, *M. volkensii* nuts that are protected by the outer hardcover exhibited prolonged viability. This is probably because the stony layer that covers the *M. volkensii* seeds protects the seeds from adverse conditions for instance very high temperatures and humidity. These results are akin to findings of Chen *et al.* (2013) that seeds which are enclosed by shell are shielded from exposure to oxygen and light. The findings confirmed that nuts are free from damage caused by pests, mold and excess external moisture; hard shell provides greater protection against microbial contamination. Considering that *M. volkensii* thrives in arid and semi-arid habitats, the pulp and the hard seed coat enables this tree species to remain viable until the conditions are right for germination.

Seeds stored in open containers under cold room temperature had a significance difference and a higher mean germination percentage as compared to seeds stored in open containers under room temperature. This can be attributed to the ability of a cold room to store seeds at the correct temperature while controlling the moisture level unlike the case of ambient temperature. The results agree with the findings by Hirano, *et al.* (2009) who reported that seeds should be dried and stored at cold temperatures for long-term preservation. Cryopreservation, or freezing in liquid nitrogen at -180°C can also be used for extremely long-term storage. The results are entirely in line with what Vasques *et al.* (2014) observed in seeds of three Mediterranean shrub species stored at a low temperature. The stored seeds in their study maintained a high viability and vigor for a couple of years but ambient temperature storage led to a marked decline after one year ending in almost complete mortality after 4 years of shelf storage. By contrast, Croft *et al.* (2012) stated that very cold storage for instance refrigerators consists of quite high relative humidity which can damage seeds due to high moisture content, therefore according to their study, storing seeds in cold storage is not fit for seed viability.

Nonetheless, seeds stored at room temperature in sealed containers for 2, 4, and 6 months showed a higher germination percentage than those kept in open containers but the same temperature conditions. This is probably due to lack of a barrier upon the open containers (between the seed and the outside atmosphere), thus permitting movement of oxygen and water leading to irreversible seed deterioration. The low germination agrees with findings of Jour *et al.* (2018) that open containers are unable to create a barrier between the seed and the outside atmosphere, which allows movement of oxygen and water between the outside atmosphere and the stored seed. Consequently, this causes a rise of metabolic

activity, high respiration and fungal attack causing the seeds to die and as a result low germination is observed. The results are related to the observation of Alemayehu *et al.* (2020), who observed that agriculture seeds placed in atmospheric temperatures in open bins sometimes result in limited storage life and seed degeneration in hot, humid locations. These findings are more or less similar to the study of Steven *et al.* (2014) who reported that air is made up of 21 % by volume of oxygen, which reacts with essential materials found in the seeds. The reaction of oxygen with seeds decreases the grade of seeds and their germination rate.

Similar findings were also observed by Jour *et al.* (2018) that when seeds are stored at 7-8% MC in closed containers, they maintain over 50 % germination for a considerable number of months. There is some indication that a further reduction of MC will conserve viability even better. Sealed or hermetic storage systems are a very effective means of controlling seed moisture and insect activity, especially in tropical conditions (Pan *et al.*, 2019). According to Mattana, *et al.* (2019), reduction of moisture content of seeds up to 10 percent after seeds are dry, make the seeds to achieve moderate germination of seeds under diverse conditions of storage. The study as well agrees with the findings of Fenolossa *et al.* (2020), who stated that viability and germination tested in two different populations of *Carissa edulis* revealed germination of 70–90% for seeds which have led to reduction of moisture content. In comparison, percentage of germination condensed to 20 percent in population two, which had higher moisture content. This study is related to findings of (Himstedt, 2002) who stated that Macadamia kernels rapidly develop rancidity when stored at room temperature at higher moisture content.

5.2 Conclusions

- i. The seed storage study showed that the storage conditions of seeds and nuts significantly affected the seed germination capacity of *M. volkensii*, therefore seed germination depends on storage conditions. The study concludes that *M. volkensii* seeds and nuts stored for six months had the highest germination levels as compared to storage at 2 and 4 months in regardless of the storage conditions.
- ii. From the study of assessing the population status of *Melia volkensii*, it is evident that the risk of tree depletion has no significant effects on the population status of *Melia volkensii* and associated tree species in Kasaala Location of Kitui County. Findings from this study will help improve the production of *M. volkensii* and associated tree species thus promote its propagation and conservation.

5.3 Recommendations

5.3.1. Recommendations from the study

- i. *Melia volkensii* seeds should be stored and distributed as extracted seeds as opposed to current practice of storing and distributing these seeds when enclosed in nuts. This will reduce cost of storage and transportation due to bulkiness of *M. volkensii* nuts.
- ii. There is a great need to conduct more extensive research on the population status of *M. volkensii* and associated tree species in Kitui County. Tree species with poor regeneration should be prioritized during rehabilitation and conservation of the tree species in the study area.

5.3.2 Areas for further research

- i. Detailed research should be done to evaluate the biotic and abiotic factors that affect seed germination and growth of *Melia volkensii*.
- ii. Further research should be done to assess the dependence between germination and seed reserves as well as their mobilization during germination of different species of *Melia* tree.
- iii. Further research is required to quantitatively assess the impacts of storing *M. volkensii* seeds for more than 6 months in order to determine whether germination of this tree species will rise above the 89% that was recorded at 6 months storage.
- iv. Detailed regeneration studies should be done in Kitui County so as to fully understand the mechanisms that promote natural regeneration.
- v. Conservation stakeholders should device alternative source of livelihood for local community so as to discourage anthropogenic activities that lead to overconsumption, pollution and overexploitation of forest products. This will help to reduce the pressure on the forest and hence conserving the environment.

REFERENCES

- Addo-Fordjour, P., Obeng, S., Anning, A. and Addo, M. (2009).** Floristic composition, structure and natural regeneration in a moist semi-deciduous forest following anthropogenic disturbances and plant invasion. *International journal of biodiversity and conservation* **1** (2), 021-037.
- Alados, C. L., Gotor, P., Ballester, P., Navas, D., Escos, J. M., Navarro, T. and Cabezudo, B. (2006).** Association Between Competition and Facilitation Processes and Vegetation Spatial Patterns in Alpha Steppes. *Biological Journal of the Linnean Society* **87** (1), 103-113.
- Albrecht, M. and Gotelli, N.J.(2001).** ‘Spatial and Temporal Niche Partitioning in Grassland Aunts’. *Oecological* 126 (1): 134-141. Bibcode:2001.Oecol 126. 134A.doi: 10.10.007/s00420. Oecol. 126 C.
- Alemayehu, S., Abay, F., Ayimut, K. M., Assefa, D., Chala, A., Mahroof, R., Harvey, J. and Subramanyam, B. (2020)** Evaluating Different Hermetic Storage Technologies to Arrest Mold Growth, Prevent Mycotoxin Accumulation and Preserve Germination Quality of Stored Chickpea in Ethiopia. *Journal of Stored Products Research* 85, doi:10.1016/j. jspr.2019.101526.
- Al-Maliki, S., Al-Taey, D. K. and Al-Mammori, H. Z. (2021).** Earthworms and eco-consequences: Considerations to soil biological indicators and plant function: A review. *Acta Ecologica Sinica*, 41(6), 512-523.
- Angalaaera, L.C.N., Cobbina, J. and Sinclair, F.L (2011).** The effect of land use systems on tree diversity: Farmer preference and Species Composition of Cocoa-based Agroecosystems in Ghana. *Agroforest Sys* 81-249-265.https://doi.org/10.1007/s10457-010-9366-z.
- Anjarwalla P, Belmain S, Sola P, Jamnadass R and Stevenson PC. (2016.)** Handbook on Pesticidal Plants. World Agroforestry Centre (ICRAF), Nairobi, Kenya).
- Badola, H. K. and Pal, M (2002).** “Endangered Medicinal Plant Species in Himachal Pradesh,” *Current Science*83 (7):797– 798.
- Baldos, O., J. DeFrank and G.S. Sakamoto (2011).** Improving Germination of Piligrass (*heteropogon contortus*) seeds using liquid smoke flavoring. *HortScience* 46: S312 (abstr.)
- Bandyopadhyay, P., Nelson, D., Greenwood, M., Brittan, G. and Berwald, J. (2011).** The logic of Simpson’s paradox. *Synthese* 181 (2).185-208.

- Barber RG, Thomas DB. and Moore TR (1981).** Studies on soil erosion and runoff and proposed design procedures for terraces in the cultivated semi-arid areas of Machakos District, Kenya. In *Soil Conservation: Problems and Prospects* (Morgan R.P.C. ed.). John Wiley and Sons, Chichester, England.
- Baskin J.M. and Baskin C.C. (2004).** "A classification system for seed dormancy". *Seed Science Research*. **14**: 1–16.
- Begon, M., J.L. Harper, and Townsend C.R. (1996).** *Ecology: Individuals, Populations, and Communities*. Blackwell Science Ltd., Cambridge, MA.
- Berjak P. and Pammenter NW (1997).** Progress in the understanding and manipulation of desiccation-sensitive (recalcitrant) seeds. In *Basic and applied aspects of seed biology* (pp. 689-703). Springer, Dordrecht.
- Beth M.L. Morrison and Chase D. Mendenhall (2020).** Hummingbird-Plant Interactions are More Specialized in Forest Compared to Coffee Plantations, Diversity, 10.3390/d12040126, 12, 4, 9126).
- Bewley, J. D., Bradford, K. J., Hilhorst, H. W. and Nonogaki, H. (2013).** Germination. In *Seeds* (pp. 133-181). Springer, New York, NY.
- Bharat K. Pradhan and Hemant K. Batola (2012)** Effect of Storage Conditions and Storage Periods on Seed Germination in Eleven Populations of *Swertia Chirayita*: A critically Endangered Medicinal Herb in Himalaya'' The Scientific World Journal, vol.2012, Article ID 128105, 9 Pages, 2012.<https://doi.org/10.1100/2012/128105>.
- Blanc, L., Maury-Lechon, G. and Pascal, J. P. (2000).** Structure, floristic composition and natural regeneration in the forests of Cat Tien National Park, Vietnam: an analysis of the successional trends. *Journal of biogeography* **27** (1): 141-157.
- Blomley, T. (1994).** Indigenous Agroforestry: *Melia volkensii* in Kenya. *Agroforestry Today* 6 (4): 10-11.
- Bonner, FT (2008).** Storage of Seed. In Bonner FT, Karrfalt. RP, editors. *The Woody Plant Seed Manual*. United States. Department of Agriculture, Forest Service, Agriculture Handbook 727.pp85-95.
- Borda-Nino, M., Ceccon, E., Meli, P., Hernandez-Mucino, D., Mas, J. F. and Brancalion, P. H. (2021).** Integrating farmers' decisions on the assessment of forest regeneration drivers in a rural landscape of Southeastern Brazil. *Perspectives in Ecology and Conservation*, 19(3), 338-344.

- Breman, E., Ballesteros, D., Castillo-Lorenzo, E., Cockel, C., Dickie, J., Faruk, A. and Ulian, T. (2021).** Plant Diversity Conservation Challenges and Prospects—the Perspective of Botanic Gardens and the Millennium Seed Bank. *Plants*, 10(11), 2371.
- Bustam, S., Ali, M. S. M., Sinniah, U. R., Shamsuddin, N. A. and Kadir, A. M. A. (2021).** Short-term storage of seeds and cryopreservation of embryonic axes of *Lepisanthes fruticosa*.
- Butola, J. S. and Badola, H. K. (2004).** “Effect of pre-sowing treatment on seed germination and seedling vigour in *Angelica glauca*, The Scientific World Journal 9 threatened medicinal herb,” *Current Science* 87(6):796–799.
- Capers, R. S. (2000).** A comparison of two sampling techniques in the study of submersed macrophyte richness and abundance. *Aquatic Botany*, 68(1), 87-92.
- Caviedes J. and Ibarra JT (2017)** Influence of Anthropogenic Disturbances on Stand Structural Complexity in Andean Temperate Forests: Implications for Managing Key Habitat for Biodiversity.
- Chen, A. H., Robinson-Mosher, A., Savage, D. F., Silver, P. A. and Polka, J. K. (2013).** The bacterial carbon-fixing organelle is formed by shell envelopment of preassembled cargo. *PLoS one*, 8(9), e76127.
- Chesson P. (2000)** Mechanisms of maintenance of species diversity. *Annual review of Ecology and Systematics* (1):343-66.
- Croft, M., Bicksler, A., Manson, J. and Burnette, R. (2012)** Vacuum Sealing vs. Refrigeration: Which is the Most Effective Way to Store Seeds? *ECHO Asia Notes*.14: 1-6. Available: <https://www.echocommunity.org/en/resources/5262cfab-ddcf-42ec-843a-82f8468f0829>.
- Crossland, M., Valencia, A. M. P., Pagella, T., Magaju, C., Kiura, E., Winoweicki, L., and Sinclair, F. (2021).** Onto the farm, into the home: how intrahousehold gender dynamics shape land restoration in Eastern Kenya. *Ecological Restoration*, 39(1-2), 90-107.
- Dale, M.R.T. (1999):** Spatial pattern analysis in plant ecology. – Cambridge University Press, Cambridge.
- Dechnik-Vázquez YA, Meave JA, Pérez-García EA, Gallardo-Cruz JA. and Romero-Romero MA. (2016).** The Effect of Treefall Gaps on the Understorey Structure and Composition of the Tropical Dry Forest of Nizanda, Oaxaca, Mexico: Implications for Forest Regeneration. *Journal of Tropical Ecology*. 2(2):89-106.

- Dickie, J. B. and Smith, R. D. (2008).** “Observations on the Survival of Seeds of *Agathis* spp. Stored at Low Moisture Contents and Temperatures’. *Seed Science Research*, **5** (1), 5-14.
- Duffy, J. Emmett (2006)** “Biodiversity and the Functioning of Seagrass Ecosystems.” *Marine Ecology Progress Series*, vol. 311, Inter-Research Science Center, pp. 233–50. <http://www.jstor.org/stable/24870048>.
- Ellis, R. H., Hong, T. D., and Roberts, E. H. (1992).** The low-moisture-content limit to the negative logarithmic relation between seed longevity and moisture content in three subspecies of rice. *Annals of Botany*, **69**(1), 53-58.
- Esilaba, A. O., Okoti, M., Nyariki, D. M., Keya, G. A., Miriti, J. M., Kigomo, J. N. and Cheruiyot, H. K. (2011).** The desert margins programme approaches in upscaling best-bet technologies in arid and semi-arid lands in Kenya. In *Innovations as Key to the Green Revolution in Africa* (pp. 1177-1191). Springer, Dordrecht.
- FAO and UNEP. (2020).** The State of the World’s Forests 2020. Forests, biodiversity and people. Rome. <https://doi.org/10.4060/ca8642en>.
- Faust, Mildred E (2020).** “Germination of *Populus Grandidentata* and *P. Tremuloides*, with Particular Reference to Oxygen Consumption. “*Botanical Gazette* **97**, no.4 (1936): 808-21. <http://www.jstor.Org/stable/2471529>.
- Fenollosa, E., Jené, L. and Munné-Bosch, S. (2020).** A rapid and sensitive method to assess seed longevity through accelerated aging in an invasive plant species. *Plant Methods*, **16**(1), 1-11.
- Ferlian, O., Cesarz, S., Craven, D., Hines, J., Barry, K.E., Bruelheide, H., Buscot, F., Haider, S., Heklau, H., Herrmann, S., Kühn, P., Pruschitzki, U., Schadler, M., Wagg, C., Weigelt, A., Wubet, T. and Fisenhauer, N. (2018).** Mycorrhiza in tree diversity-ecosystem function relationships: conceptual framework and experimental implementation. *Ecosphere* (Washington, DC). **9**(5). e 02226.
- Flores, B.M., Staal, A., Jakovac, C.C. et al (2020).** Soil erosion as a resilience drain in Disturbed Tropical Forests. ,” *Journal of Environmental Studies and Sciences’* **Plant Soil** **450**, 11-25 <https://doi.Org/10.1007/s11104-019-04097-8>.
- Food and Agriculture Organization of the United Nation (FAO) (2010).** Global Forest Assessment, FAO: Rome Italy.

- Gachathi, F.N., Johansson S.G and Alakoski- Johansson G.M (1994).** A Check-List of Indigenous Trees and Shrubs of Bura, Tana River District, Kenya with Malakote, Orma and Somali Names," *Journal of East African Natural History* 83(2), 117-141.
- Ghanbari, S., Sefidi, K., Kern, C. C. and Álvarez-Álvarez, P. (2021).** Population structure and regeneration status of woody plants in relation to the human interventions, Arasbaran Biosphere Reserve, Iran. *Forests*, 12(2), 191.
- Githae E.W and Mutiga I.M. (2021).** Ecological Restoration of Pastoral Landscapes in the Drylands of East Africa. *Journal of Dryland Agriculture* 7 no. 3: 34-41.
- Gladys, K.V. (2017).** "Rainfall and temperature variability and its effect on food security in Kitui County, Kenya", *International Journal of Development and Sustainability*, Vol. 6 No. 8, pp. 924-939.
- Guillon, T., Dumont, Y. and Fourcaud, T. (2012).** Numerical methods for the biomechanics of growing trees. *Computers and Mathematics with Applications*, 64 (3), 289-309.
- Himstedt, SR (2002).** Oil Content and other Components as Indicators of Quality and Shelf life of Macadamia Kernels (Maiden and Betche).
- Hirano, T., Godo, T., Miyoshi, K., Ishikawa, K., Ishikawa, M. and Mii, M. (2009).** Cryopreservation and low-temperature storage of seeds of *Phaius tankervilleae*. *Plant Biotechnology Reports*, 3(1), 103-109.
- Hong, T.D. and R.H. Ellis. (1996).** A protocol to determine seed storage behavior. IPGRI
- Hong, T.D., S. Linington and R.H. Ellis. 1996.** Seed Storage Behaviour: a Compendium. Handbooks for Genebanks: No. 4. International Plant Genetic Resources Institute, Rome.
- ICRAF, A Selection of Useful Trees and Shrubs for Kenya (1992).** Notes on Their Identification, Propagation and Management for Use by Farming and Pastoral Communities, International Centre for Research in Agroforestry (ICRAF), Nairobi, Kenya.
- International Journal of Development and Sustainability Vol. 6, No. 8 (2017):** 924-939 926 ISDS www.isdsnet.com.
- Jaetzold, R., Schmidt, H., Hornetz, B., and Shisanya, C. (2006).** Farm Management Handbook of Kenya (Vol. 2/C1). Nairobi: Ministry of Agriculture.

- Jaetzold, R. and Schmidt, H. (1983).** Natural conditions and farm management: Farm management handbook of Kenya. Ministry of Livestock and Development, Nairobi Vol. 2.
- Jaoko, V., Nji Tizi Taning, C., Backx, S., Mulatya, J., Van den Abeele, J., Magomere, T., Olubayo, F., Mangelinckx, S., Werbrouck, S. and Smagghe, G. (2020).** The Phytochemical Composition of *Melia volkensii* and Its Potential for Insect Pest Management. *Plants (Basel, Switzerland)*, 9(2),143.
- Jiang, Z. et al. (2016)** Phytochrome B and RELLE1/2-mediated signaling controls seed dormancy and germination in Arabidopsis. *Nat. Commun.*
- Jin, Yi, Sabrina E. Russo, and Mingjian Yu. (2018):** "Effects of light and topography on regeneration and coexistence of evergreen and deciduous tree species in a Chinese subtropical forest." *Journal of ecology* 106, no. 4 1634-1645.
- Johnson, J. (2018).** *Mechanisms of coexistence between native and exotic plant species* (Doctoral dissertation, University of British Columbia).
- Jones, M.M., Tuomisto, H., Borcard, D., Legendre, P., Clark, D.B. and Olivas, P.C., (2008).** Explaining variation in tropical plant community composition: influence of environmental and spatial data quality.
- Joshi AR. and Joshi K (2007).** Ethnomedicinal plants used against skin diseases in some villages of Kali Gandaki Bagmati and Tadi Likhu watersheds of Nepal. *Ethnobotanical Leaflet*; 11:235–46.
- Jour, T.Y., Haque, A.U and Saleh, P.Y (2018).** Effect of Moisture on Germination and Pathogen Contamination in Different Containers for Storability of Jute Seed.
- Kamondo, B.M., Kariuki, J. G., Luvanda, A. M., Muturi, G. M. and Ochieng, D. (2016).** *Guideline on Production, Distribution and Use of Improved Melia Seed and Seedlings in the Drylands of Kenya* (2016 ed.). KEFRI, Kenya: Kenya Forestry Research Institute.
- Kayombo, C., Rubanza, C. and Giliba, R. (2020).** Effect of Human Disturbances on Mahungu Green Belt Forest Reserve (MGFR) in Dodoma City, Central Tanzania. *East African Journal of Forestry and Agroforestry*, 2(2), 1-15.
- Kershaw, K.A. and Looney, J.H.H. (1985):** Quantitative and Dynamic Plant Ecology. – Edward Arnold, Baltimore. 3rd. ed.
- Khaine, I., Woo, S., Kwak, M., Seong, H., Mije, S., Hana, Y., Park, Euddeum, L., Li, Yang., Haenaem, K., Kyu, J., and Jieun k. (2018).** Factors affecting Natural Regeneration of Tropical Forests across a Precipitation Gradient in Myanmar. *Forests*. 9,143. <https://doi.org/10.3390/f9030143>.

- Kidundo M. (1997).** Participatory technology development and Nursery Propagation of *Melia volkensii*: A potential Agroforestry tree species for semi-arid Mbeere. MSc Thesis, University of Wales, Bangor.
- Kigomo N.B. (2001).** State of Forest genetic Resources in Kenya. Sub-Regional Workshop FAO/IPGRI/ICRAF on the Conservation, management, sustainable utilisation and Enhancement of Forest Genetic Resources in Sahelian and North-Sudanian Africa (Ouagadougou, Burkina Faso, 22-24 September 1998). Forest Genetic Resources Working Papers, Working Paper FGR/18E. Forestry Department, FAO, Rome, Italy.
- Kimondo, J. and Kiamba, K. (2005).** Overview of Natural Distribution, Propagation and Management of *Melia volkensii*: Recent Mukau.
- Kisangau, P. D., Kimiti, J. M., Mburu, M. W. and Snelder, D. J. (2021).** Factors influencing use of multipurpose trees and shrubs in arid and semiarid lands of Kenya. *Rangeland Ecology and Management*, 74, 125-134.
- Kivunzya, A. N. (2019).** *Characterisation of livestock production systems and its contribution to the food security in Kitui County, Kenya* (Doctoral dissertation).
- Kivunzya, A. N., Kanui, I. T., Amwata, A. D. and Ndungwa, K. A. (2018).** The status of livestock production systems in semi-arid farming and arid pastoral agro-ecological zones in Kitui County, Kenya. *International Journal of Education and Research*, 6(2), 141-148.
- Kruger, L.M., Midgley, J.J. and Cowling, R.M. (1997)** Resprouters vs reseeders in South African forest trees: a model based on forest canopy height. *Functional Ecology*. 11, pgs 101 – 105.
- Kyalo, E. (2004).** An Overview of *Melia volkensii* Propagation at Tiva Nursery, Kitui District. *Research and Teachnology Development of Mukau (Melia Volkensii Gurke)*, 23-24.
- Ky-Dembele C., Tigabu M., Bayala J., Savadogo P., Boussim I.J. and Odén P.C. (2010).** Clonal Propagation of *Detariummicrocarpum* from Root Cuttings. *Silva Fennica*, 44(5), 775-786.
- Lampert, A (2019).** Over-exploitation of Natural Resources is followed by Inevitable Declines in Economic Growth and Discount Rate. *Nat- Commun* 10, 1419-<https://doi.org/10.1038/s41467-019-09246-2>.
- Lemma, D. T. and Megersa, H. G. (2021).** Impact of climate change on East African coffee production and its mitigation strategies. *World Journal of Agricultural Sciences*, 17(2), 81-89.

- Liu, U., Cossu, T. A., Davies, R. M., Forest, F., Dickie, J. B. and Breman, E. (2020).** Conserving orthodox seeds of globally threatened plants ex situ in the Millennium Seed Bank, Royal Botanic Gardens, and Kew, UK: the status of seed collections. *Biodiversity and Conservation*, 29(9), 2901-2949.
- Lohani, D., Joshi, R. and Dwivedi, B. (1980).** Vegetative Propagation of Forest Species. *UP Forest Department Bulletin, Uttar Pradesh* (41).
- Lugadiru J. and Wafula A (2016).** *Journal of Natural Sciences Research* www.iiste.org ISSN 2224-3186 (Paper) ISSN 2225-0921 (Online) Vol.6, No.2.
- Luvanda, A., Musyoki, J., Cheboiwo, J., Wekesa, L. and Ozawa, M. (2015).** An Assessment of the Socio-Economic Importance of *Melia volkensii* Based Enterprises in Kenya.
- Mabberley, D. J. (1984).** A Monograph of *Melia* in Asia and the Pacific. The History of White Cedar and Persian lilac. *Gardens' Bulletin* **37** (1), 49-64.
- Macdonald, D. W., Chiaverini, L., Bothwell, H. M., Kaszta, Ĺ., Ash, E., Bolongon, G. and Cushman, S. A. (2020).** Predicting biodiversity richness in rapidly changing landscapes: climate, low human pressure or protection as salvation? *Biodiversity and Conservation*, 29(14), 4035-4057.
- Mackie E. D. and Matthews R. W. Forest Mensuration (2006),** A Handbook For Practitioners (HMSO).
- Magurran, A. E. (1988).** *Ecological diversity and its measurement*. Princeton university press.
- Manning, A. D., Cunningham, R. B., Tongway, D. and Lindenmayer, D. B. (2020).** Woodlands and woody debris: Understanding structure and composition to inform restoration. *Plos one*, 15(3), e0224258.
- Mattana, E.,Peguero, B., Di Sacco, A. et. Al. (2019).** Assessing Seed Dessication responses of Native trees in the Caribbean. *New Forests*.
- Matula, R., Svátek, M., Kůrová, J., Úradníček, L., Kadavý, J. and Kneifl, M. (2012).** The sprouting ability of the main tree species in Central European coppices: implications for coppice restoration. *European Journal of Forest Research*, 131(5), 1501-1511.
- Matz, S. A. (1984).** Nuts and fruits. In *Snack Food Technology* (pp. 83-99). Springer, Dordrecht.

- Maua, J. O., Mugatsia Tsingalia, H., Cheboiwo, J. and Odee, D. (2020).** Population structure and regeneration status of woody species in a remnant tropical forest: A case study of South Nandi forest, Kenya. *Global Ecology and Conservation*, 21, e00820.
- Maundu, P and Bo Tengas (2005).** Useful trees and shrubs for Kenya, World Agroforestry Centre. ISBN-9966-896-70-8. www.worldagroforestrycentre.org.
- Mbiti, J. and Misango, S. (2021).** Organizational Culture and Structure as Determinants of Service Delivery in the County Government of Kitui. *European Journal of Business and Strategic Management*, 6(1), 73-88.
- Mbora, A. and Simons, A. (2003).** Annotated inventory of agroforestry related work around Mt. Kenya (1983-2002). *World Agroforestry Centre, Nairobi Kenya*.
- Midgley, J.J and William J Bond, (1996).** Why the World's Vegetation is not totally dominated by resprouting Plants; because resprouters are shorter than seeders.
- Milimo, P. and Hellum, A. (1989).** Dormancy in seeds of *Melia volkensii*. *East African Agricultural and Forestry Journal*, 54 (3), 111-122.
- Milimo, P. B. (1989).** *Preliminary Studies on Vegetative Propagation of Melia volkensii by Cuttings*. Paper presented at the Proceedings of IFS meeting, Stockholm, Sweden.
- Milimo, P. B. (2013).** *The control of germination in Melia volkensii seeds* (Doctoral dissertation).
- Miller, J. H. and K. V. Miller. (1999).** Forest Plants of the Southeast and Their Wildlife Uses. Auburn, AL: Southern Weed Science Society.
- Mulanda S.E., Adero O.M., Amugane O.N., Akunda E. and Kinyamario I.J. (2013).** Prospects for a rapid in vitro regeneration system for propagation of the pesticidal tree *Melia volkensii*, Gurke.
- Mulatya J.M. (2000).** Tree root development and interactions in drylands: focusing on *Melia volkensii* with socio-economic evaluation. PhD Dissertation, University of Dundee, Dundee, UK.
- Muok, B., Mwamburi, A., Kyalo, E. and Auka, S. (2010).** Growing *Melia volkensii*. *A Guide for Farmers and Tree Growers in the Drylands*, 3.
- Musinguzi, P., Bosselmann, A. S. and Pouliot, M. (2018).** Livelihoods-conservation initiatives: Evidence of socio-economic impacts from organic honey production in Mwingi, Eastern Kenya. *Forest Policy and Economics*, 97, 132-145.

- Muthike G and Githiomi J (2020).** Value Chain Analysis of Farm Grown *Melia volkensii* (Gurke) Timber in the South Eastern Dry Lands of Kenya. *Journal of Horticulture and Forestry* (ISSN: 2006-9782) Vol.12 (4), pp.115-121. DOI: 10.5897/JHF2020.0641.
- Mutiga, I. M. (2021).** Ecological restoration of pastoral landscapes in the drylands of East Africa. *Journal of Dryland Agriculture*, 7(3), 34-41.
- Mwadalu, R., Gathara, M., Muturi, G. and Mbuvi, M. T. E. (2021).** Potential of *Casuarina equisetifolia* and *Melia Volkensii* Tree Species in Improving Soil Fertility in Kwale and Kilifi Counties, Kenya.
- Mwamburi, A., Kimondo, J. M. and Kyalo, E. (2005).** Traditional methods used by farmers to break seed dormancy in *Melia volkensii* in eastern and coastal provinces of Kenya. In *Recent Mukau (Melia volkensii Gurke) Research and Development. Proceedings of the First National Workshop, KEFRI Kitui Regional Research Center. November 16 to* (Vol. 19, pp. 28-31).
- Nasreen, S. (2000).**"Effects of storage temperature, storage period and seed moisture content on seed viability of mash bean (*Vigna mungo*)." *Pakistan Journal of Biological Sciences (Pakistan)*.
- Ndegwa, G.M., Nehren, U., Grüninger, F. et al (2016).** Charcoal Production through Selective Logging leads to Degradation of dry woodlands: a Case study of Mutomo District, Kenya. *J.Arid Land* 8, 618-631 [https:// doi.org/10.1007/s40333-016-0124-6](https://doi.org/10.1007/s40333-016-0124-6).
- Nicolescu, V., Hochbichler, E., and Bruckman, V. (2016).** Sustainable Biomass Potentials from Coppice Forests for Pyrolysis: Chances and Limitations. In V. Bruckman, E. Apaydin Varol, B. Uzun, and J.Liu (EDs.), *Biochar: A Regional Supply Chain Approach in View of Climate Change Mitigation* (pp. 139-161). Cambridge University University Press.doi: 10.1017/9781316337974.008
- Nieuwenhuis, M. and Egan, D. (2002),** An Evaluation and Comparison of Mechanized and Manual Planting on Afforestation and Reforestation Sites in Ireland. *International Journal of Forest Engineering*, 13(2): 11-23.
- Ntubiri C. Kajuju (2010).** Clonal Propagation of *Melia volkensii* (Gürke) Trees for Seed Orchard Establishment in Kiambeere, Embu County, Kenya.
- Odum, E.P., 1971.** Fundamentals of Ecology (3rd edn.) WB Saunders Co. *Philadelphia, Pennsylvania*.
- Omondi, W. (2004).** Ecological correlates of seed desiccation tolerance in tropical African dryland trees. *American Journal of Botany*, **91** (6), 863-870.

- Orwa, C., Mutua, A., Kindt, R., Jamnadass, R. and Anthony, S (2009).** Agroforestry Database: A Tree Reference and Selection Guide Version 4.0.
- Pang, X., Nordström, E. M., Böttcher, H., Trubins, R. and Mörtberg, U. (2017).** Trade-offs and synergies among ecosystem services under different forest management scenarios–The LEcA tool. *Ecosystem Services*, 28, 67-79.
- Parthasarathy, N. (2001).** Changes in Forest Composition and Structure in Three Sites of Tropical Evergreen Forest Around Sengaltheri, Western Ghats. *Current Science*, 389-393.
- Pickett, S. T. (1980).** Non-equilibrium coexistence of plants. *Bulletin of the Torrey Botanical Club*, 238-248.
- Plumptre, A. J. and Reynolds, V. (1994).** The Effect of Selective Logging on the Primate Populations in the Budongo Forest Reserve, Uganda. *Journal of Applied Ecology*, 31(4), 631–641. <https://doi.org/10.2307/2404154>.
- Poorter, L. (2004).** *Biodiversity of West African forests: an ecological atlas of woody plant species*: CABI.
- Pradhan, B. K and Badola, H. K (2008).** Seed Germination Response of Populations of *Swertiachirayita* following Periodical Storage, *Seed Technology*, Vol. 30, no. 1.
- Probert, R. J. (2003).** Seed viability under ambient conditions, and the importance of drying. *Seed conservation: turning science into practice*, 337-365.
- Qinggang Wang, Dachuan Bao, Yili Guo, Junmeng Lu, Zhijun Lu, Yaozhan Xu, Kuihan Zhang, Haibo Liu, Hongjie Meng, Mingxi Jiang, Xiujuan Qiao and Handong Huang (2014).** Species Associations in a Species-Rich Subtropical Forest Explained by Stochastic Geometry of Biodiversity.
- Queenborough SA, Burslem DF, Garwood NC and Valencia R. (2009)** Taxonomic scale-dependence of habitat niche partitioning and biotic neighborhood on survival of tropical tree seedlings. *Proceedings of the Royal Society B: Biological Sciences*. 276(1676):4197-205.
- Rajjou, L. and Debeaujon, I. (2008).** Seed longevity: survival and maintenance of high germination ability of dry seeds. *Comptes rendus biologiques*, 331(10), 796–805. <https://doi.org/10.1016/j.crv.2008.07.021>.
- Ranal, M. A., Santana, D. G. d., Ferreira, W. R. and Mendes-Rodrigues, C. (2009).** Calculating germination measurements and organizing spreadsheets. *Brazilian Journal of Botany*, 32 (4), 849-855.

- Ross, S.M. and Dykes, A. (1996).** Soil Conditions, erosion and Nutrient Loss on Steep Slopes under Mixed Dipterocarp Forest In Brunei Darussalam. In: Edwards, D.s., et al. (Eds.)Tropical Rainforest Research- Current issues. Kluwar, The Netherlands, PP.259-270.
- Rumpf, S. B., Hülber, K., Klonner, G., Moser, D., Schütz, M., Wessely, J., and Dullinger, S. (2018).** Range dynamics of mountain plants decrease with elevation. *Proceedings of the National Academy of Sciences*, 115(8), 1848-1853.
- Runo, M. S., Muluvi, G. M. and Odee, D. W. (2004).** Analysis of genetic structure in *Melia volkensii* (Gurke.) populations using random amplified polymorphic DNA. *African Journal of Biotechnology*, 3(8), 421-425.
- Shuruti Singh and Kanu Priy (2020)** Impact of Human Activities on Forest. Journal of Critical Reviews (ISSN-2394-5125) VOL 7, ISSUE 9.
- Sisay, G., Gitima, G., Mersha, M. and Alemu, W. G. (2021).** Assessment of land use land cover dynamics and its drivers in Bechet Watershed Upper Blue Nile Basin, Ethiopia. *Remote Sensing Applications: Society and Environment*, 24, 100648.
- Śliwińska, E. (2000).** Analysis of the Cell Cycle in Sugarbeet Seed duringDevelopment, Maturation and Germination. pp. 133–139. in Black, M.;Bradford, K.J.;Vázquez-Ramos, J., (Eds) *Seed biology: Advances and applications*. Wallingford, CABI Publishing.
- Spellerberg, I.F. (1991)** Monitoring Ecological Change. Cambridge University Press, Cambridge, UK.
- Spellerberg, Ian F., and Peter J. Fedor. (2003)** A tribute to Claude Shannon (1916–2001) and a plea for more rigorous use of species richness, species diversity and the ‘Shannon–Wiener’Index. *Global Ecology and Biogeography* 12.3, 177-179.
- Stephenson N. L., Das A. J., Condit R., Russo S. E., Baker P. J., Beckman N. G., et al. 2014.** Rate of Tree Carbon Accumulation Increases Continuously with Tree Size. *Nature* 507:90–93.
- Steven P.C Groot., Liesbeth de Groot. and Jankoddeand Rob Van Treuren (2014).** Prolonging the Longevity of ex situ Conserved Seeds by Storage under Anoxia”. *Plant Genetic Resources* Doi: <https://doi.Org/10.1017/s147926114000586>.
- Stewart, M. and Blomley, T. (1994).** Use of *Melia volkensii* in a Semi-Arid Agroforestry System in Kenya. *The Commonwealth Forestry Review*, 73 (2), 128-131.
- Styles, B.T. (1972).** The flower biology of the Meliaceae and its bearing on tree breeding. *Sivlae Genetica*, nr. 21, pp. 175-182.

- Sugimoto, K., Gordon, S. P. and Meyerowitz, E. M. (2011).** Regeneration in plants and animals: dedifferentiation, transdifferentiation, or just differentiation?. *Trends in cell biology*, 21(4), 212-218.
- Tedd, J. (1997).** *Perception, management and usage of Melia volkensii by farmers: A case study from the Kibwezi district, Kenya* (Doctoral dissertation, University of Nottingham).
- Tirado, R., Pugnaire, F. I. and Eriksson, O. (2005).** Community Structure and Positive Interactions in Constraining Environments. *Oikos*, 111 (3), 437-444.
- Tweddle, J.C. Dickie, J.B., Baskin, C.C. and Baskin, J.M. (2003).** The Ecological Aspects of Seeds Desiccation Sensitivity (2003). *Journal of Ecology* 91(2), 294-304.
- UNESCO- World heritage Centre (2010).** Tentative lists of State parties to the Convention., *A Guideline for on Tsavo park and Chyulu hills complex., Ref 5515.* KWS, Kenya Wildlife services.
- Van Treuren R, Bas N, Kodde J, Groot SPC. and Kik C. (2018);** Rapid loss of seed viability in *ex-situ* conserved wheat and barley at 4°C as compared to -20°C storage. *Conserv Physiol.* 6(1):coy033. doi: 10.1093/conphys/coy033. PMID: 29977565; PMCID: PMC6016650.
- Vasques A., Ramon V., Santos, M. C., and Keizer J. J (2014).** The Role of Cold Storage and Seed Source in the Germination of Three Mediterranean Shrub Species with Contrasting Dormancy Types.
- Vernooy, Ronnie (2017).** “The roles of community seed banks in climate change adaptation”. *Development in Practice*, 27(3): 316-327. Doi: 10.1080/09614524.2017.1294653. S2.
- Vertucci, C.W and Roos, E. E (1991).** Theoretical Basis of Protocols for Seed Storage, *Plant Physiology*, vol. 94, no. 3, pp. 1019– 1023, 1990. “Seed Moisture Content, Storage, Viability and vigour,” *Seed Science Research*, vol. 1, pp. 277–279.
- Wang, Hexin, Genzhu Li, Dongmei Yu, and Ying-min Chen. (2008):** "Barrier effect of litter layer on natural regeneration of forests: a review." *Chinese Journal of Ecology* 27, no. 1,83-88.
- Warui, M. W., Gikungu, M., Bosselmann, A. S. and Hansted, L. (2018).** Pollination of Acacia woodlands and honey production by honey bees in Kitui, Kenya. *Future of Food: Journal on Food, Agriculture and Society*, 6(1), 40-50.
- Xu, Y., Zhang, Y., Shi, M. and Lu, Z. (2020).** Environmental variation, functional diversity and identity predicting community biomass in an old-growth subtropical broad-leaved forest. *Global Ecology and Conservation*, 23, e01093.

- Wekesa L, Muturi G, Mulatya J, Esilaba AO, Keya GA. and Ihure S. (2012).** Economic Viability of *Melia volkensii* (Gurke) Production on Smallholdings in Drylands of Kenya. *International Research Journal of Agricultural Science and Soil Science* (ISSN: 2251-0044)2(8): 364-369.
- Whittaker, R. J., Willis, K. J. and Field, R. (2001).** Scale and species richness: towards a general, hierarchical theory of species diversity. *Journal of biogeography*, 28(4), 453-470.
- Wilcove, D. S., Rothstein, D., Dubow, J., Phillips, A. and Losos, E. (1998).** Quantifying Threats to Imperiled Species in the United States. *BioScience*, 48 (8), 607-615.
- Wilps, H., Nasseh, O., Rembold, H and Krall, S (1993).** The Effect of Extracts of *Melia volkensii* (Gurke) on Mortality and Fitness of Adualt *Schistocerca Gregaria* (Forsk.) Orth., Cyrtacanthacrinae). *J. Appl ENT.* 116: 12-19.
- Woods, N. N., Dows, B. L., Goldstein, E. B., Moore, L. J., Young, D. R. and Zinnert, J. C. (2019).** Interaction of seed dispersal and environmental filtering affects woody encroachment patterns in coastal grassland. *Ecosphere*, 10(7), e02818.
- Yang Liu, Liping Zhang, Jike Lu, Weimin Chen, Gehong Wei. and Yanbing Lin (2019).** Topography Affects the Soil Condition and Bacteria Communities along a restoration Gradient on Loess Plateau. *Journal on Applied Soil Ecology* Volume 150, June, 103471.
- Yili Guo., Bin Wang., Azim, U., Mallik, Fuzhao Huang., Wusheng Xiang., Tao, Ding., Shujun, Wen., Shuhua, Lu, Dongxing, Li and Yunlin He(2016).** *Journal of Plant Ecology*, Volume 10, Issue 3, 1 June 2017, Pages 450-460, <https://doi.org/10.1093/jpe/rtw057>.
- Yohannis, M. A. (2019).** *ICTS As A Bridge Between Climate Information And Livelihood Strategies Among Rural Women In Kitui County, Kenya* (Doctoral dissertation, UoN).
- Zhang, K., Zhang, Y., Sun, J., Meng, J. and Tao, J. (2021).** Deterioration of orthodox seeds during ageing: Influencing factors, physiological alterations and the role of reactive oxygen species. *Plant Physiology and Biochemistry*, 158, 475-485.
- Ziwei Liu.,Yan Zhu., Jianjun Wang., Wu Ma and Jinghui Meng (2019).** Species Association of the Dominant Tree Species in an Old-Growth Forest and Implications for Enrichment Planting for the Restoration of Natural Degraded Forest in Subtropical China.

APPENDICES

Appendix 1: Composition of tree species in transect 1

Species	Number	Percentage
<i>Grewia tephrodermis</i>	12	13.48
<i>Acacia tortilis</i>	10	11.24
<i>Adansonia digitata</i>	7	7.87
<i>Melia volkensii</i>	6	6.74
<i>Tennantia sennii</i>	6	6.74
<i>Albizia anthelmintica</i>	5	5.62
<i>Cassia abbreviate</i>	5	5.62
<i>Acacia nilotica</i>	4	4.49
<i>Combretum aculeatum</i>	4	4.49
<i>Acacia Senegal</i>	3	3.37
<i>Balanites aegyptiaca</i>	3	3.37
<i>Commiphora Africana</i>	3	3.37
<i>Commiphora baluensis</i>	3	3.37
<i>Delonix alata</i>	3	3.37
<i>Tamarindus indica</i>	3	3.37
<i>Delosperma cooperi</i>	2	2.25
<i>Grewia bicolor</i>	2	2.25
<i>Grewia villosa</i>	2	2.25
<i>Acacia mellifera</i>	1	1.12
<i>Bauhinia coriacea</i>	1	1.12
<i>Bauhinia taitensis</i>	1	1.12
<i>Boscia angustifolia</i>	1	1.12
<i>Commiphora edulis</i>	1	1.12
<i>Tephrosia villosa</i>	1	1.12
Total	89	100

Appendix 2: Composition of tree species in transect 2

Species	Number	Percentage
<i>Melia volkensii</i>	24	24.24
<i>Acacia tortilis</i>	11	11.11
<i>Sterculia rhynchorcarpa</i>	10	10.10
<i>Adansonia digitata</i>	7	7.07
<i>Acacia nilotica</i>	4	4.04
<i>Cassia abbreviate</i>	4	4.04
<i>Combretum aculeatum</i>	4	4.04
<i>Grewia villosa</i>	4	4.04
<i>Tamarindus indica</i>	4	4.04
<i>Bauhinia nataliensis</i>	3	3.03
<i>Grewia tephrodermis</i>	3	3.03
<i>Berchemia discolor</i>	2	2.02
<i>Commiphora Africana</i>	2	2.02
<i>Commiphora baluensis</i>	2	2.02
<i>Commiphora edulis</i>	2	2.02
<i>Erythrina melanacantha</i>	2	2.02
<i>Grewia flava</i>	2	2.02
<i>Terminalia prunioides</i>	2	2.02
<i>Acacia Senegal</i>	1	1.01
<i>Calyptrorhiza taitensis</i>	1	1.01
<i>Commiphora eminii</i>	1	1.01
<i>Euphorbia ingens</i>	1	1.01
<i>Euphorbia tirucalli</i>	1	1.01
<i>Lanea schimperi</i>	1	1.01
<i>Thylachium thomasii</i>	1	1.01
Total	99	100

In transect 2, *Melia volkensii* was most represented tree species 24.24% (n=24), followed by *Acacia tortilis* 11.11% (n=11). There was a total of 99 tree species.

Appendix 3: Composition of tree species in transect 3

Species	Number	Percentage
<i>Melia volkensii</i>	12	20.69
<i>Acacia tortilis</i>	10	17.24
<i>Adansonia digitata</i>	8	13.79
<i>Delonix alata</i>	4	6.90
<i>Grewia tephrodermis</i>	4	6.90
<i>Acacia busciae</i>	3	5.17
<i>Grewiavillosa</i>	3	5.17
<i>Acacia nilotica</i>	2	3.45
<i>Berchemia discolor</i>	2	3.45
<i>Lannea schimperi</i>	2	3.45
<i>Terminalia brownie</i>	2	3.45
<i>Boscia angustifolia</i>	1	1.72
<i>Commiphora africana</i>	1	1.72
<i>Commiphora baluensis</i>	1	1.72
<i>Grewia calymmatosepala</i>	1	1.72
<i>Grewia flava</i>	1	1.72
<i>Maerua crassifolia</i>	1	1.72
Total	58	100

In transect 3, *Melia volkensii* was most represented tree species 20.69% (n=12), followed by *Acacia tortilis* 17.24% (n=10) and *Adansonia digitata* 13.79% (n=8).

There was a total of 58 tree species.

Appendix 4: Composition of tree species in transect 4

Species	Number	Percentage
<i>Acacia tortilis</i>	11	17.46
<i>Melia volkensii</i>	11	17.46
<i>Adansonia digitata</i>	10	15.87
<i>Acacia nilotica</i>	3	4.76
<i>Balanites aegyptiaca</i>	3	4.76
<i>Lannea schimperi</i>	3	4.76
<i>Bauhinia nataliensis</i>	2	3.17
<i>Berchemia discolor</i>	2	3.17
<i>Boscia angustifolia</i>	2	3.17
<i>Commiphora africana</i>	2	3.17
<i>Grewia villosa</i>	2	3.17
<i>Cassia abbreviate</i>	1	1.59
<i>Combretum aculeatum</i>	1	1.59
<i>Commiphora baluensis</i>	1	1.59
<i>Duosperma cooperi</i>	1	1.59
<i>Erythrina melanacantha</i>	1	1.59
<i>Grewia calymmatosepala</i>	1	1.59
<i>Grewia tephrodermis</i>	4	6.06
<i>Maerua decumbens</i>	1	1.59
<i>Solanum campylacanthum</i>	1	1.59
<i>Sterculia rhynchorcarpa</i>	1	1.59
<i>Tamarindus indica</i>	1	1.59
<i>Tephrosia villosa</i>	1	1.59
Total	66	100

In transect 4, both *Acacia tortilis* and *Melia volkensii* were equally represented amongst tree species 17.46% (n=11) being the highest in the population, followed by *Adansoniadigitata* 15.87% (n=10). There was a total of 63 tree species.

Appendix 5: Composition of tree species in transect 5

Species	Number	Percentage
<i>Grewia tephrodermis</i>	8	10.39
<i>Commiphora edulis</i>	7	9.09
<i>Grewia calymmatosepala</i>	7	9.09
<i>Acacia tortilis</i>	6	7.79
<i>Adansonia digitata</i>	6	7.79
<i>Combretum aculeatum</i>	6	7.79
<i>Bauhinia nataliensis</i>	3	3.90
<i>Commiphora africana</i>	3	3.90
<i>Melia volkensii</i>	3	3.90
<i>Acacia Senegal</i>	2	2.60
<i>Albizia anthelmintica</i>	2	2.60
<i>Boscia angustifolia</i>	2	2.60
<i>Calyptrorhiza taitensis</i>	2	2.60
<i>Commiphora eminii</i>	2	2.60
<i>Delonix alata</i>	2	2.60
<i>Duosperma cooperi</i>	2	2.60
<i>Lannea schimperi</i>	2	2.60
<i>Maerua decumbens</i>	2	2.60
<i>Cassia abbreviate</i>	1	1.30
<i>Commiphora baluensis</i>	1	1.30
<i>Commiphora campensis</i>	1	1.30
<i>Grewia africana</i>	1	1.30
<i>Grewia flava</i>	1	1.30
<i>Grewia villosa</i>	1	1.30
<i>Maerua holstii</i>	1	1.30
<i>Solanum taitensis</i>	1	1.30
<i>Sterculia rhynchorcarpa</i>	1	1.30
<i>Tennantia sennii</i>	1	1.30
Total	77	100

In transect 5, both *Grewiatephrodermis* 10.39% (n=8), *Commiphoraedulis* 9.09% (n=7) and *Grewia calymmatosepala*9.09% (n=7) were most represented amongst the population. *Melia volkensii* was 3.90% (n=3) of the population. There was a total of 77 tree species.

Appendix 6: Composition of tree species in transect 6

Species	Number	Percentage
<i>Acacia nilotica</i>	10	11.90
<i>Acacia mellifera</i>	6	7.14
<i>Acacia Senegal</i>	6	7.14
<i>Acacia tortilis</i>	6	7.14
<i>Adansonia digitata</i>	6	7.14
<i>Lannea schimperi</i>	5	5.95
<i>Albizia anthelmintica</i>	4	4.76
<i>Boscia angustifolia</i>	4	4.76
<i>Melia volkensii</i>	4	4.76
<i>Balanites aegyptiaca</i>	3	3.57
<i>Berchemia discolor</i>	3	3.57
<i>Cassia abbreviate</i>	3	3.57
<i>Grewia villosa</i>	3	3.57
<i>Solanum campylacanthum</i>	3	3.57
<i>Commiphora edulis</i>	2	2.38
<i>Delonix alata</i>	2	2.38
<i>Erythrina melanacantha</i>	2	2.38
<i>Grewia calymmatosepala</i>	2	2.38
<i>Maerua decumbens</i>	2	2.38
<i>Terminalia prunioides</i>	2	2.38
<i>Calotropis strophela</i>	1	1.19
<i>Calyptrorhiza taitensis</i>	1	1.19
<i>Dalbergia melanoxylon</i>	1	1.19
<i>Duosperma cooperi</i>	1	1.19
<i>Grewia tephrodermis</i>	1	1.19
<i>Thylachium thomasii</i>	1	1.19
Total	84	100

In transect 6, *Acacia nilotica* 11.90% (n=10) had the highest representation in the population, followed by *Acacia mellifera* 7.14% (n=6), *Acacia senegal* 7.14% (n=6), *Acacia tortilis* 7.14% (n=6) and *Adansoniadigitata* 7.14% (n=6). *Melia volkensii* was 4.76% (n=4) of the population. There was a total of 84 tree species.

Appendix 7: Composition of tree species in transect 7

Species	Number	Percentage
<i>Adansonia digitata</i>	8	10.53
<i>Melia volkensii</i>	7	9.21
<i>Acacia nilotica</i>	6	7.89
<i>Bauhinia nataliensis</i>	6	7.89
<i>Commiphora edulis</i>	5	6.58
<i>Commiphora eminii</i>	4	5.26
<i>Delonix alata</i>	4	5.26
<i>Acacia senegal</i>	3	3.95
<i>Balanite aegyptiaca</i>	3	3.95
<i>Cassia abbreviate</i>	3	3.95
<i>Erythrina melanacantha</i>	3	3.95
<i>Grewia villosa</i>	3	3.95
<i>Terminalia prunioides</i>	3	3.95
<i>Albizia anthelmintica</i>	2	2.63
<i>Boscia angustifolia</i>	2	2.63
<i>Combretum aculeatum</i>	2	2.63
<i>Lannea schimperi</i>	2	2.63
<i>Maerua decumbens</i>	2	2.63
<i>Maerua holstii</i>	2	2.63
<i>Sterculia rhynchorcarpa</i>	2	2.63
<i>Acacia tortilis</i>	1	1.32
<i>Grewia africana</i>	1	1.32
<i>Grewia calymmatosepala</i>	1	1.32
<i>Solanum taitensis</i>	1	1.32
<i>Eurphobia schefferi</i>	1	1.32
<i>Bridelia taitensis</i>	2	2.63
Total	79	100

In transect 7, *Adansonia digitata* (n=8) had the highest numbers in the population (10.53%) and *Melia volkensii* 9.21% (n=7) had the second highest numbers in the population. There was a total of 76 tree species.

Appendix 8: DBH size-class distribution of *Melia volkensii* in the study area (Kasaala**Location)**

DBH clusters	Transects							Total
	1	2	3	4	5	6	7	
0.5 – 10	0		1	0	0	0	0	4
10.5 – 20	1	10	3	2	0	0	1	17
20.5 – 30	1	3	4	7	2	2	4	23
30.5 – 40	1	6	3	1	1	2	1	15
40.5 – 50	0	0	1	1	0	0	1	3
50.5 – 60	2	1	0	0	0	0	0	3
60.5 – 70	1	0	0	0	0	0	0	1
70.5 – 80	0	1	0	0	0	0	0	1
Total	6	24	12	11	3	4	7	67

Appendix 9: Diversity of *M. volkensii* and associated tree species in Kasaaala

Location

Tree species	Species richness (mean values)	Groups
<i>Adansonia digitata</i>	0.51	a
<i>Acacia busciae</i>	0.48	ab
<i>Acacia tortilis</i>	0.44	ab
<i>Albizia anthelmintica</i>	0.38	ab
<i>Acacia nilotica</i>	0.37	ab
<i>Acacia senegal</i>	0.37	ab
<i>Terminalia prunioides</i>	0.36	ab
<i>Tamarindus indica</i>	0.36	ab
<i>Balanite aegyptiaca</i>	0.35	ab
<i>Delosperma cooperi</i>	0.30	ab
<i>Grewia bicolor</i>	0.30	ab
<i>Terminalia brownii</i>	0.30	ab
<i>Acacia mellifera</i>	0.30	ab
<i>Tennantia sennii</i>	0.30	ab
<i>Erythrina melanacantha</i>	0.27	ab
<i>Delonix alata</i>	0.26	ab
<i>Grewia tephrodermis</i>	0.26	b
<i>Bauhinia taitensis</i>	0.20	b
<i>Maerua holstii</i>	0.20	b
<i>Sterculia rhynchorcarpa</i>	0.18	b
<i>Commiphora edulis</i>	0.16	b
<i>Boscia angustifolia</i>	0.15	b
<i>Grewia calymmatosepala</i>	0.15	b
<i>Lannea schimperi</i>	0.15	b
<i>Commiphora eminii</i>	0.15	b
<i>Combretum aculeatum</i>	0.14	b
<i>Cassia abbreviata</i>	0.13	b
<i>Commiphora africana</i>	0.11	b
<i>Grewia villosa</i>	0.11	b
<i>Commiphora baluensis</i>	0.10	b
<i>Duosperma cooperi</i>	0.10	b
<i>Grewia flava</i>	0.10	b
<i>Berchemia discolor</i>	0.09	b
<i>Maerua decumbens</i>	0.05	b
<i>Bauhinia coriacea</i>	0.01	b
<i>Calotropis strophela</i>	0.01	b
<i>Calyptrotheca taitensis</i>	0.01	b
<i>Commiphora campensis</i>	0.01	b
<i>Dalbergia melanoxyton</i>	0.01	b
<i>Euphorbia ingens</i>	0.01	b
<i>Euphorbia tirucalli</i>	0.01	b
<i>Euphorbia scheffleri</i>	0.01	b
<i>Grewia africana</i>	0.01	b
<i>Maerua crassifolia</i>	0.01	b
<i>Melia volkensii</i>	0.01	b
<i>Solanum campylacanthum</i>	0.01	b
<i>Solanum taitensis</i>	0.01	b
<i>Tephrosia villosa</i>	0.01	b
<i>Thylachium thomasii</i>	0.01	b

